

ACKNOWLEDGEMENTS

The author expresses his deep sense of gratitude to Dr.R.Ramakrishna Reddy Lecturer in Physics, Sri Krishnadevaraya University Anantapur, for his inspiring guidance and constant encouragement through out the progress of this work.

The author is much indebted to Prof.S.V.Subrahmanyam for having created the opportunity of prosecuting his higher studies.

The author expresses his sincere gratitude to Prof.E.Rajagopal, Head of Department of Physics, S.K.University, Anantapur for having provided the facilities to complete this work.

The author feels much elated in expressing his thanks to Prof.T.V.Ramakrishna Rao, Sri Krishnadevaraya University, Anantapur for his valuable suggestions throughout the progress of this work.

The author's thanks are due to his research colleague Sri.R.Viswanath for his assistance in bringing the thesis to the present form.

The author records his deepest sense of gratitude for his parents to whom the present work has been dedicated to.

The author expresses his sincere thanks to Smt.R.Ramatulsamma wife of Dr.R.Ramakrishna Reddy for having extended her co-operation in the present thesis.

The author seizes the opportunity to extend a warm feeling of gratitude to his wife Smt.C.Praneela for having spurred his idea to acquire the degree of Master of Philosophy.

The author's sincere thanks are due to Sri.P.Raghunatha Reddy, Grade I Lecturer in Chemistry.