

LIST OF FIGURES

S. No.	Fig. Legends	PAGE No.
1.	Fig 1.1.1: Worldwide mortality rate of cancer in both female and males	2
2.	Fig 1.2.1: Acquired Capabilities of Cancer	6
3.	Fig 1.3.1: Over all NF- κ B activation pathway in response to various signals	14
4.	Fig 1.3.2: Role of NF- κ B in expression and activation of different pathway proteins	18
5.	Fig.1.4.1: Apoptosis - Mechanisms of Initiation and Regulation. Refer to text for descriptions. DISC, death-inducing signaling complex; FADD, Fas-associated death domain; tBid, truncated Bid; Apaf-1, apoptotic protease-activating factor 1; Cyt c, Cytochrome c; ATP, adenosine triphosphate; PARP, poly (ADP-ribose) polymerase.	22
6.	Fig. 1.7.1: Chemical structure of Resveratrol	26
7.	Fig.1.7.2: Schematic representation of mechanisms of anticancer and anti-inflammatory properties of Resveratrol	30
8.	Fig.1.8.1: Steps involved in structure based drug design through <i>in silico</i> approach.	36
9.	Fig.1.8.2: Pathway of transport of drugs across biological membranes (Lehninger <i>et al.</i> , 1993)	42
10.	Fig.1.9.1: Thermodynamic cycle for Enzyme, Inhibitor at various solvent parameters and change in free energy (ΔG) under vacuo and water conditions under E-I complex condition.	52
11.	Fig 3.2.1.1: Minimum acceptable solubility in $\mu\text{g/ml}$. Bars show the minimum solubility for low, medium and high permeability (K_a) at different clinical dose	76
12.	Fig. 3.2.3.1: Chemical Conversation of MTT to Formazan	79
13.	Fig. 3.2.3.2: The biochemistry of DNA fragmentation and the appearance of the DNA ladder	81
14.	Fig.3.3.4.1 Parameters of grid log file used in docking simulation	92
15.	Fig.3.3.4.1 Parameters of docking log file used in docking simulation	93
16.	Fig.3.3.5.1 Parameters of Energy minimization file used in Molecular dynamic simulations	94
17.	Fig.3.3.5.2 Parameters of Equilibrium Phase-I file used in Molecular dynamic simulations	95
18.	Fig.3.3.5.3 Parameters of Equilibrium Phase-II file used in Molecular dynamic simulations	96
19.	Fig. 4.2.1: Schematic representation of RESL-11 and RESL-12 synthesis.	107

20.	Fig. 4.2.2 Assigned Spectra for RESL11 400 MHz ¹ H Spectrum	108
21.	Fig. 4.2.3: Chemical ionization Mass spectrum of RESL11	109
22.	Fig. 4.2.4: Assigned Spectra for RESL12 400 MHz ¹ H Spectrum	110
23.	Fig.4.2.5: Chemical ionization Mass spectrum of RESL12	111
24.	Fig. 4.2.6: Schematic representation of RESL43	112
25.	Fig. 4.2.7 Assigned Spectra for RESL43 400 MHz ¹ H Spectrum	113
26.	Fig.4.2.8: Chemical ionization Mass spectrum of RESL43	114
27.	Fig 4.3.1: MTT uptake % of death of U937 cell with different concentrations of RESL and its analogs	116
28.	Fig 4.3.2: PI staining of U937 with RESL analogs (A. Untreated, B. Oleyandrin C. RESL D. RESL11, RESL12 and RESL43)	117
29.	Fig 4.3.3: Fragmentation of genomic DNA in cells after treatment for 72h with 10µm concentration of RESL and its analogs	118
30.	Fig. 4.3.4: a) Effect of RESL and its analogs on cleavage of PARP b-e) TNF, IL-8, Fas and FasL levels after treatment with RESL and its analogs	120
31.	Fig. 4.3.5: U937 cells were pretreated with TNF and RESL analogs and incubated for 72h, the nuclear extract were prepared and NF-κB DNA-protein binding activity was determined by EMSA	122
32.	Fig 4.4.1: Overall view of the NF-κB p50/p65 heterodimer-DNA complex structure along the DNA helical axis with the approximate dyad vertical. The DNA duplex is in blue. Color coding for the protein is as in Secondary structure elements is labeled.	124
33.	Fig. 4.4.2: Right zoom view is NF-κB DNA binding amino acids (Pink color, p50 and green color, p65), Left zoom view shown the Electrostatic surface with active site residues and the extreme left with electrostatic potentials proposed binding site (Blue +ve charge and Red -ve charge). These figures were generated with Pymol (DeLano, 2002).	125
34.	Fig. 4.4.3: Schematic representation of DNA interaction with NF-κB [chain A (p65); Chain B (P50)]	126
35.	Fig. 4.5.1: RESL binding interaction with DNA binding residues (red color sticks) of p65 chain of NF-κB	128
36.	Fig. 4.5.2: RESL11 binding interaction with both chains of NF-κB DNA (Yellow color, p50; Red color p65) binding residues.	128
37.	Fig. 4.5.3: RESL12 binding mode with NF-κB DNA binding residues (Yellow color, p50; Red color p65)	129
38.	Fig. 4.5.4: Tight binding mode of RESL43 with the DNA binding region of NF-κB (Yellow color, p50; Red color p65)	129
39.	Fig. 4.6.1: A-D) Root Mean Square Deviation of native NF-κB and NF-κB-RESL complexes at 299K temp for 10000ps	133
40.	Fig. 4.6.2: Time evolution of the radius of gyration (Rg)	133

	during 6000 ps of MD simulation native NF- κ B and NF- κ B-RESL complexes	
41.	Fig. 4.6.3: A-C) The rmsf values of unliganded NF- κ B and RESL complexes were plotted against residue numbers.	136
42.	Fig. 4.6.4: The profile of atomic fluctuations of DNA binding residues before and after binding of RESL analogs with NF- κ B.	137
43.	Fig. 4.6.5: Time evolution of number hydrogen bonds formed in between the NF- κ B and RESL analogs	137
44.	Fig. 4.6.6: Time evolution of number of hydrogen bond index in between the NF- κ B and RESL analogs	138
45.	Fig. 4.7.1: The DNA interaction with NF- κ B RESL complex; Zoom top view represents the interaction of RESL at DNA binding region (Yellow color stick (p65) Cyan color stick (p50)) of NF- κ B. The Zoom bottom view shows the binding interaction of DNA (Black color ribbons) with the complex of NF- κ B-RESL	141
46.	Fig. 4.7.2: The DNA interaction with NF- κ B RESL11 complex; Zoom top view represents the interaction of RESL11 at DNA binding region (Yellow color stick (p65) Cyan color stick (p50)) of NF- κ B. The Zoom bottom view shows the binding interaction of DNA (Black color ribbons) with the complex of NF- κ B-RESL11	142
47.	Fig. 4.7.3: The DNA interaction with NF- κ B RESL12 complex; Zoom top view represents the interaction of RESL12 at DNA binding region (Yellow color stick (p65) Cyan color stick (p50)) of NF- κ B. The Zoom bottom view shows the binding interaction of DNA (Black color ribbons) with the complex of NF- κ B-RESL12	143
48.	Fig. 4.7.4: The DNA interaction with NF- κ B RESL43 complex; Zoom bottom view represents the interaction of RESL43 at DNA binding region (Yellow color stick (p65) Cyan color stick (p50)) of NF- κ B. The Zoom top view shows the binding interaction of DNA (Black color ribbons) with the complex of NF- κ B-RESL43	144