

- Abdallah, F.B., Kallel, H., Bakhrouf, A., 2009. Enzymatic, outer membrane proteins and plasmid alterations of starved *Vibrio parahaemolyticus* and *Vibrio alginolyticus* cells in sea water. *Arch. Microbiol.* 191, 493-500.
- Abraham, T.J., Manley, R., 1995. Luminous and non-luminous *Vibrio harveyi* associated with shell disease in cultured *Penaeus indicus*. *Journal of Aquaculture in Tropics.* 10, 273-276.
- Abraham, W.R., 2006. Controlling biofilms of gram-positive pathogenic bacteria. *Curr. Med. Chem.* 13, 1509-1524.
- Adams, A., 1991. Response of penaeid shrimp to exposure to *Vibrio* species. *Fish Shellfish Immunol.* 1, 59-70.
- Aguerre-Guzman, G., Vazquez-Juarez, R., Ascencis, F., 2001. Difference in the susceptibility of American white shrimp larval substages (*Litopenaeus vannamei*) to four *Vibrio* sp. *J. Invert. Pathol.* 78, 215-219.
- Alam, M., Miyoshi, S.I., Yamamoto, S., Tomochika, K.I., Shinoda, S., 1996. Expression of virulence-related properties by intestinal adhesiveness of *V.mimicus* strains isolated from aquatic environment. *Appl. Environ. Microbiol.* 62, 3871-3874.
- Alavandi, S.V., Manoranjita, V., Vijayan, K.K., Kalaimani, N., Santiago, T.C., 2006. Phenotypic and molecular typing of *Vibrio harveyi* isolates and their pathogenicity to tiger shrimp larvae. *Letts. Appl. Microbiol.* 43, 566-570.
- Alcaide, E., 2003. Numerical Taxonomy of Vibrionaceae isolated from cultured Amberjack (*Seriola dumerili*) and surrounding water. *Curr. Microbiol.* 46 184–189.
- Alekshun, M.N., Levy, S.B., 2007. Molecular mechanisms of antibacterial multidrug resistance. *Cell.* 128, 1037-1050.
- Allen, A.E., Booth, M.G., Frischer, M.E., Verity, P.G., Zehr, J.P., Zani, S., 2001. Diversity and detection of nitrate assimilation genes in marine bacteria. *Appl. Environ. Microbiol.* 67, 5343-5348.
- Alsina, M., Blanch, A.R., 1994a. A set of keys for biochemical identification of *Vibrio* species. *J. Appl. Bacteriol.* 76, 79-85.

References

- Alsina M, Blanch A, 1994b. Improvement and update of a set of keys for biochemical identification of *Vibrio* species. *J. Appl. Bacteriol.* 77, 719-721.
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J., 1990. Basic local alignment search tool. *J. Mol. Biol.* 215, 403-410.
- Aly, S.M., Mohamed, M.F., John, G. 2008b. Effect of probiotics on the survival, growth and challenge infection in *Tilapia nilotica* (*Oreochromis nilotica*). *Aquacult. Res.* 39, 647-656.
- Anderson, D.P., 1992. Immunostimulants, adjuvants, and vaccine carriers in fish: applications to aquaculture. *Annu. Rev. Fish. Dis.* 2, 281-307.
- Angulo, F.J., Nargund, V.N., Chiller, T.C., 2004. Evidence of an association between use of anti-microbial agents in food animals and anti-microbial resistance among bacteria isolated from humans and the human health consequences of such resistance. *J. Vet. Medi. Series B.* 51, 374-379.
- Anzar, R., Ludwig, W., Schleifer, H.H., 1993. Ribotyping and randomly amplified polymorphic DNA analysis of *Vibrio vulnificus* biotypes. *Syst. Appl. Microbiol.* 16, 303-309.
- Applegate, B., Kelly, C., Lackey, L., McPherson, J., Kehrmeier, S., Menn, F.M., Bienkowski, P., Salyer, G., 1997. Pseudomonas putida B2: a tod-lux bioluminescent reporter for toluene and trichloroethylene co-metabolism. *J. Ind. Microbiol. Biotechnol.* 18, 4-9.
- Applegate, B. M., Kehrmeier, S. R., Salyer, G. S., 1997. A chromosomally based tod-luxCDABE whole-cell receptor for benzene, toluene, ethylbenzene and toluene (BTEX) sensing. *Appl. Environ. Microbiol.* 64, 2730-2735.
- Arias, C.R., Verdonck, L., Swings, J., Aznar, R., Garay, E., 1997a. A polyphasic approach to study the intraspecific diversity amongst *Vibrio vulnificus* isolates. *Syst. Appl. Microbiol.* 20, 622-633.
- Arias, C.R., Verdonck, L., Swings, J., Aznar, R., Garay, E., 1997b. Intraspecific differentiation of *Vibrio vulnificus* biotypes by amplified fragment length polymorphism and ribotyping. *Appl. Environ. Microbiol.* 63, 2600-2606.

References

- Arijo, S., Rico, R., Chabrillon, M., Diaz-Rosales, P., Marti nez-Manzanares, E., Balebona, M, C., MagariÇos, B., Toranzo, A, E., MoriÇigo, M, A., 2005. Effectiveness of a divalent vaccine for sole, *Solea senegalensis* (Kaup), against *Vibrio harveyi* and *Photobacterium damsela* subsp. *piscicida*. *J. Fish Dis.* 28, 33-38.
- Armada, S.P., Farto, R., Perez, M.J., Nieto, T.P., 2003. Effect of temperature, salinity and nutrient content on the survival responses of *Vibrio splendidus* biotype I. *Microbiology.* 149, 369-375.
- Austin, B., Baudet, E., Stobie, M., 1992. Inhibition of bacterial fish pathogens *Tetrasetmis suezica*. *J. Fish Dis.* 15, 55-61.
- Austin, B., Stuckey, L.F., Robertson, P.A.W., Effendi, I., Griffith, D.R.W., 1995. A probiotic strain of *Vibrio alginolyticus* effective in reducing diseases caused by *Aeromonas salmonicida*, *Vibrio anguillarum* and *Vibrio ordalli*. *J. Fish Dis.* 18, 93-96.
- Austin, B., Austin, D.A., Falconer, V.M., Pedersen, K., Larsen, J.L., Swings, J., Verdonck, L., 1996. Dissociation of *Vibrio anguillarum* and *V.ordalii* cultures into two or three discrete colony types. *Bull. Eur. Fish. Pathol.* 16, 101-103.
- Austin, B., Austin, D.A., 1999. Bacterial fish pathogens: Diseases of farmed and wild fish, 3rd (Ed). Springer-Praxis, Godalming, England.
- Austin, B., Pride, A.C., Rhodie, G.A., 2003. Association of a bacteriophage with virulence in *Vibrio harveyi*. *J. Fish. Dis.* 26, 55-58.
- Austin, B., Zhang, X.H., 2006. *Vibrio harveyi*: a significant pathogen of marine vertebrates and invertebrates. *Lett. Appl. Microbiol.* 43, 119 – 124.
- Azam. F., 2001. Introduction, history, and overview: the methods to our madness. *Methods Microbiol.* 30, 1-12.
- Aznar, R., Amaro, C., Alcaide, E., Lemos, M., 1989. Siderophore production by environmental strains of *Salmonella* species. *FEMS Microbiol. Lett.* 57, 7-12.
- Bache`re, E., 2000. Shrimp immunity and disease control. *Aquaculture.* 191, 3-11.

References

- Baffone, W., Vittoria, E., Campana, R., Citterio, B., Casaroli, A., Pierfelici, L., 2005. Occurrence and expression of virulence related properties by environmental *Vibrio* spp. In *in vitro* and *in vivo* systems. *Food Control*. 16, 451-457.
- Baffone, W., Tarsi, R., Pane, L., Campana, R., Repetto, B., Mariottini, G.L., Pruzzo C., 2006. Detection of freeliving and plankton-bound vibrios in coastal waters of the Adriatic Sea (Italy) and study of their pathogenicity-associated properties. *Environ. Microbiol.* 8, 1299-1305.
- Bai, F., Pang, L., Qi, Z., Chen, J., Austin, B., Zhang, X.H., 2008. Distribution of five *Vibrio* virulence-related genes among *Vibrio harveyi* isolates. *J. Gen. Appl. Microbiol.* 54, 71-78.
- Balcazar, J.L., Blas, I.D., Ruiz-Zarzuola, I., Cunningham, D., Vendrell, D., Muzquiz, J.L., 2006. The role of probiotics in aquaculture. *Vet. Microbiol.* 114, 173-186.
- Balcazar, J.L., Rojas-Luna, T., Cunningham, D.P., 2007. Effect of the addition of four potential probiotic strains on the survival of pacific white shrimp (*Litopenaeus vannamei*) following immersion challenge with *Vibrio parahaemolyticus*. *J. Invert. Pathol.* 96, 147-150.
- Bang, S.S., Baumann, P., Nealson, K.H., 1978. Phenotypic characterization of *Photobacterium logei* (sp. nov.), a species related to *P.fischeri*. *Curr. Microbiol.* 1, 285-288.
- Banin, E., Khare, S.K., Naider, F., Rosenberg, E., 2001. Proline-rich peptide from the coral pathogen *Vibrio shiloi* that inhibits photosynthesis of zooxanthellae. *Appl. Environ. Microbiol.* 67, 1536-1541.
- Barer, M.R., Gribbon, L.T., Harwood, C.R., Nwoguh, C.E., 1993. The viable but non-culturable hypothesis and medical microbiology. *Rev. Med. Microbiol.* 4, 183-191.
- Bassler, B.L., Gibbons, P.J., Yu, C., Roseman, S., 1991. Chitin utilization by marine bacteria. Chemotaxis to chitin oligosaccharides by *Vibrio furnissii*. *J. Biol. Chem.* 266, 24268-24275.
- Bassler, B.L., Wright, M., Showalter, R.E., Silverman, M.R., 1993. Intercellular signalling in *Vibrio harveyi*: sequence and function of genes regulating expression of luminescence. *Mol. Microbiol.* 9, 773-786.

References

- Bassler, B.L., Wright, M., Silverman, M.R., 1994a. Sequence and function of LuxO, a negative regulator of luminescence in *Vibrio harveyi*. *Mol. Microbiol.* 12 403-412.
- Bassler, B.L., Wright, M., Silverman, M.R., 1994b. Multiple signalling systems controlling expression of luminescence in *Vibrio harveyi*: sequence and function of genes encoding a second sensory pathway. *Mol. Microbiol.* 13, 273-286.
- Bassler, B.L., Greenberg, E.P., Stevens, A.M., 1997. Cross-species induction of luminescence in the quorum-sensing bacterium *Vibrio harveyi*. *J. Bacteriol.* 179, 4043-4045.
- Bauer, A., Kirby, W., Sherris, J., Turck, M., 1966. Antibiotic susceptibility testing by a standardized single disk method. *Am. J. Clin. Pathol.* 45, 493-496.
- Baumann, P., Schubert, R.H.M., 1983. Vibrionaceae. In: Bergey's Manual of Systematic Bacteriology. (ed. by N.R. Krieg & J.G. Holt). The Williams & Wilkins Co., Baltimore, MD, USA 1, 516-550.
- Baumann, P., Baumann, L., 1981. The marine Gram-negative eubacteria: genera *Photobacterium*, *Beneckea*, *Alteromonas* and *Alcaligenes*. In M.Starr, H.Stolp, H.Truper, A.Balows and H.Schlegel (ed.), the Prokaryotes. Springer-Verlag, New York, N.Y., 1302-1331.
- Baumann, P., Baumann, L., Reichelt, J.L., 1973. Taxonomy of marine bacteria: *Beneckea parahaemolytica* and *Beneckea alginolytica*. *J. Bacteriol.* 113, 1144-1155.
- Baumann, P., Baumann, L., 1977. Biology of the marine enterobacteria: genera *Beneckea* and *Photobacterium*. *Annu. Rev. Microbiol.* 31, 39-61.
- Baumann, P., Schubert, R.H.W., 1984. *Vibrionaceae*, In N. R. Krieg and G. J. Holt (Eds.), Bergey's manual of systematic bacteriology, vol.1. The Williams & Wilkins Co., Baltimore, Md. 516-550.
- Beaber, J.W., Burrus, V., Hochhut, B., Waldor, M.K., 2002a. Comparison of SXT and R391, two conjugative integrating elements: definition of a genetic backbone for the mobilization of resistance determinants. *Cell. Mol. Life Sci.* 59, 2065-2070.

References

- Beachey, E.H., 1981. Bacterial Adherence: Adhesin-Receptor Interactions Mediating the Attachment of Bacteria to Mucosal Surfaces. *J. Infect Dis.* 143, (3): 325-345.
- Ben-Haim, Y., Thompson, F.L., Thompson, C., Cnockaert, M., Hoste, B., Swings, J., Rosenberg, E., 2003. *Vibrio coralliilyticus* sp. nov., a temperature dependent pathogen of the coral *Pocillopora damicornis*. *Int. J. Syst. Evol. Microbiol.* 53, 309–315.
- Benitez, J.A., Silva, A.J., Finkelstein, R.A., 2001. Environmental signals controlling production of hemagglutinin/protease in *Vibrio cholerae*. *Infect. Immun.* 69, 6549-6553.
- Benitez-Nelson, C.R., 2000. The biogeochemical cycling of phosphorus in marine systems. *Earth. Sci. Rev.* 51, 109-135.
- Bonin, P., 1996. Anaerobic nitrate reduction to ammonium in two strains isolated from a coastal marine sediment: a dissimilatory pathway. *FEMS Microbol. Ecol.* 19, 27-38.
- Bordas, A., Balebona, M.C., Rodriguez-Maroto, J.M., Borrego, J.J., Moriñigo, M.A., 1998. Chemotaxis of Pathogenic *Vibrio* Strains towards Mucus Surfaces of Gilt-Head Sea Bream (*Sparus aurata* L.). *Appl. Environ. Microbiol.* 64, 1573-1575.
- Bourne, D.G., Høj, L., Webster, N.S., Swan, J., Hall, M.R., 2006. Biofilm development within a larval rearing tank of the tropical rock lobster, *Panulirus ornatus*. *Aquaculture.* 260, 27–38.
- Bowser, P.R., R.Rosemar, C.R.Reiner, 1981. A preliminary report of vibriosis in cultured American lobster, *Homarus americanus*. *J. Invertebr. Path.* 37, 80-85.
- Boyd, C.D., Hollerman, W.D., Plum, J.A., Saeed, M. 1984. Effect of treatment with a commercial bacterial suspension on water quality of channel catfish pond. *Prog. Fish. Cult.* 46, 36-40.
- Boyd, E.F., Davis, B.M., Hochhut, B., 2001. Bacteriophage-bacteriophage interactions in the evolution of pathogenic bacteria. *Trends Microbiol.* 9, 137-144.
- Bradford, M.M., 1976. A refined and sensitive method for the quantification of microgram quantity of proteins utilizing the principle of protein dye binding. *Ann. Biochem.* 72, 248-254.

References

- Bramhachari, P.V., Dubey, S.K., 2006. Isolation and characterization of exopolysaccharide produced by *Vibrio harveyi* strain VB23. *Letts. Appl. Microbiol.* 43, 571-577.
- Breed, R.S., Murray, E.G.D., Smith, N.R., 1957. *Bergey's manual of Determinative Bacteriology*, 7th Ed, Lippincott Williamd & Wilkins, Baltimore, Md. 229-249.
- *Brock, T.D., 1999. *Robert Koch. A life in medicine and bacteriology*. ASM Press, Washington, D.C.
- Brown, C., 1989. A study of the 2 shellfish pathogenic vibrios isolated from a long island hatchery during a recent outbreak of disease. *J. Shellfish Res.* 1, 63-87.
- Brussow, H., Canchaya, C., Hardt, W.D., 2004. Phages and the evolution of bacterial pathogens: From genomic rearrangements to lysogenic conversion. *Microbiol. Mol. Biol. Rev.* 68, 560-602.
- Bryant, T.N., Lee, J.V., West, P.A., 1986a. Numerical classification of species of *Vibrio* and related genera. *J. Appl. Bacteriol.* 61, 437-467.
- Bryant, T.N., Lee, J.V., West, P.A., Colwell, R.R., 1986b. A probability matrix for the identification of species of *Vibrio* and related genera. *J. Appl. Bacteriol.* 61, 469-480.
- Burlage, R.S., Saylor, G.S., Larimer, F., 1990. Monitoring of naphthalene catabolism by bioluminescence with nah-lux transcriptional fusions. *J. Bacteriol.* 172, 4749-4757.
- Butler, D. A., Lobregat, C. M., Gavan, T. L., 1975. Reproducibility of the Analytab (API 20E) system. *J. Clin. Microbiol.* 2, 322-326.
- Cabello, F.C., 2006. Heavy use of prophylactic antibiotics in aquaculture: a growing problem for human and animal health and for the environment. *Environ. Microbiol.* 8, 1137-1144.
- Campbell, A.C., Buswell, J.A., 1983. The intestinal microflora of farmed dover sole (*Solea solea*) at different stages of fish development. *J. Appl. Bacteriol.* 55, 215-223.
- Cano-Gomez, A., Bourne, D.G., Hall, M.R., Owens, L., Hoj, L., 2009. Molecular identification, typing and tracking of *Vibrio harveyi*

References

- in aquaculture systems: current methods and future prospects. *Aquaculture*. 287, 1-10.
- Carmignani, G.M., Bennete, J.P., 1977. Rapid start-up of a biological filter in a closed aquaculture system. *Aquaculture*. 11, 85-88.
 - Casadevall, A., Pirofski, L.A., 2000. Host-pathogen interactions: basic concepts of microbial commensalisms, colonization, infections and disease. *Infect. Immun.* 68, 6511-6518.
 - Cashel, M., Gentry, D.R., Hernandez, V.J., Vinella, D., 1996. The stringent response. In F.C.Neidhardt, R.Curtiss III, J.L.Ingraham, E.C.C.Lin, K.B.Low, B.Magasanik, W.S.Reznikoff, M.Riley, M.Schaechter and H.E.Umbarger (ed), *Escherichia coli* and *Salmonella typhimurium*. Cellular and Molecular Biology, 2nd ed, ASM Press, Washington D.C., 1458-1496.
 - Castex, M., Chim, L., Pham, D., Lemaire, P., Wabete, N., Nicolas, J., Schmidely, P., Mariojous, C., 2008. Probioic *Pediococcus acidilactici* application in shrimp *Litopenaus stylirostris* culture subject to vibriosis in New Caledonia. *Aquaculture*. 275, 182-193.
 - Castro, D., Pujalte, M.J., Lopez-Cortes, L., Garay, E., Borrego, J.J., 2002. Vibrios isolated from the cultured manila clam (*Ruditapes philippinarum*): numerical taxonomy and antibacterial activities. *J. Appl. Microbiol.* 93, 438–447.
 - Caugant, D.A., Kristiansen, B.E., Frøholm, L.O., Bøvre, K., Selander, R.K., 1988. Clonal diversity of *Neisseria meningitidis* from a population of asymptomatic carriers. *Infect. Immun.* 56 2060–2068.
 - Cerda`-Cue`llar, M., R.A.Rossello´-Mora, Lalucat, J., J.Jofre, Blanch, A.R., 1997. *Vibrio scophthalmi* sp. nov., a new species from turbot (*Scophthalmus maximus*). *Int. J. Syst. Bacteriol.* 47, 58–61.
 - Chair, M., Dehasque, M., Poucke, S.V., Nelis, H., Sorgeloos, P., Leenheer, A.P.D., 1994. An oral challenge for turbot larvae with *Vibrio anguillarum*. *Aquac. Int.* 2, 270–272.
 - Chang-Ping Yu, Yue-Hwa Yu, 2000. Evaluation of biodegradability by the reduction of Tetrazolium violet in Biolog microplates. *Biotechn. Letts.* 22, 909-913.

References

- Chang, C.I., Liu, W.Y., Shyu, C.Z., 2000. Use of prawn blood agar hemolysis to screen for bacteria pathogenic to cultured tiger prawns *Penaeus monodon*. *Dis. Aquat. Org.* 43, 153-157.
- Chang, C.I., Lee, W.C., Shyu, C.Z., Liao, I.C., 1996. The development of shrimp blood agar for testing the hemolysis of shrimp's haemocyte by bacteria. *Chinese J. Microbiol. Immun.* 29, 232-239.
- Cheetham, R.C., Katz, M.E., 1995. A role of bacteriophage in the evolution and transfer of bacterial virulence determinants. *Mol. Microbiol.* 18, 201-208.
- Chen, F.R., Liu, P.C., Lee, K.K., 1999. Purification and partial characterization of a toxic serine protease produced by pathogenic *Vibrio alginolyticus*. *Microbes.* 98, 95-111.
- *Chen, S.N., Huang, S.L., Kuo, G.H., 1992. Studies on the epizootiology and pathogenicity of bacterial infections in cultured giant tiger prawn, *P. monodon* in Taiwan. In: Fulks, W., Main, K.L. (Eds.), *Diseases of cultured penaeid shrimp in Asia and the United States*. The Oceanic Institute, Hawaii. 195-205.
- Chen, X., Schauder, S., Potier, N., Dorsselaer, A.V., Pelczer, I. 2002. Structural identification of a bacterial quorum-sensing signal containing boron. *Nature.* 415, 545-549.
- Cheng, W., Liu, C.H., Kuo, C.M., Chen, J.C., 2005. Dietary administration of sodium alginate enhances the immune ability of white shrimp *Litopenaeus vannamei* and its resistance against *Vibrio alginolyticus*. *Fish Shellfish Immunol.* 18, 1, 1-12.
- Christensen, W.B., 1946. Urea decomposition as a means of differentiating *Proteus* and *Paracolon* cultures from each other and from *Salmonella* and *Shigella* types. *J. Bacteriol.* 52, 461-466.
- Chythanya, R., Karunasagar, I., Karunasagar, I., 2002. Inhibition of shrimp pathogenic vibrios by a marine *Pseudomonas* I-2 strain. *Aquaculture.* 208, 1-10.
- Cipriano, R.C., Griffin, B.R., Lidgerding, B.C., 1981. *Aeromonas salmonicida*: relationship between extracellular growth products and isolate virulence. *Can. J. Fish. Aquat. Sci.* 38, 1322-1326.

References

- Clark, A. G., Caroline, M., Lanigan, S., 1993. Prospects for estimating nucleotide divergence with RAPDs'. *Mol. Biol. Evol.* 10, (5): 1096- 1111.
- Coenye, T., Gevers, D., Peer, Y.V.D., Vandamme, P., Swings, J., 2005. Towards a prokaryotic genomic taxonomy. *FEMS Microbiol. Rev.* 29, 147–167.
- Cognetti, G., Maltagliati, F., Saroglia, M., 2006. The risk of “genetic pollution” in Mediterranean fish populations related to aquaculture activities. *Marine Pollution Bulletin.* 52: 11, 1321-1323.
- Cohan, F.M., 2002 .What are bacterial species? *Annu. Rev.Microbiol.* 56, 457–487. Collins, C.M., Lyne, P.M., Grange, J.M., Falkinham, J, D., 2004. *Microbiology Methods.* 8th Ed. Arnold group, Oxford University Press, London. 94-100.
- Collis, C.M., Hall, R.M., 1992. Gene cassettes from the insert region of integrons are excised as covalently closed circles. *Mol. Microbiol.* 16, 2875-2885.
- Collis, C.M., Grammaticopoulos, G., Briton, J., Stokes, H.W., Hall, R.M., 1993. Site-specific insertion of gene cassettes into integrons. *Mol. Microbiol.* 9, 41-52.
- Colquhoun, D.J., Sorum, H. 2002. Cloning, characterisation and phylogenetic analysis of the fur gene in *Vibrio salmonicida* and *Vibrio logei*. *Gene.* 296, 213-220.
- Colwell, R. R., 2002. A voyage of discovery: cholera, climate and complexity. *Environ. Microbiol.* 4, 67–69.
- Colwell, R. R., 2004. Infectious disease and environment: cholera as a paradigm for waterborne disease. *Perspectives.* 7, 285-289.
- Conejero, M.J.U., Hedreyda, C.T., 2003. Isolation of partial toxR gene of *Vibrio harveyi* and design of toxR-targeted PCR primers for species detection. *J. Appl. Microbiol.* 95, 602-611.
- Conejero, M.J.U., Hedreyda, C.T., 2004. PCR detection of hemolysin (*vhh*) gene in *Vibrio harveyi*. *J. Gen. Appl. Microbiol.* 50, 137-142.

References

- *Corpe, W. A., 1980. Biofilm: A survival strategy of bacteria. In Adsorption of Microorganisms to Surfaces (eds Bitton, G. and Marshall, K. C.), John Wiley, New York. 105–144.
- Costerton, J.W., Geesey, G.G., Cheng, K.J., 1978. How bacteria stick? *Sci. Am.* 238, 86–95.
- Costerton, J.W., Irvin, R.T., Cheng, K.J., 1981. The role of bacterial surface structures in pathogenesis. *Crit. Rev. Microbiol.* 8, 303-338.
- *Cowan, S.T., Steel, K.J., (Eds). 1965. Manual for the identification of medical bacteria. 1st Ed. Cambridge University Press.
- Coyer, J., Cabello-Pasini, A., Swift, A., Alberte, H., 1996. N₂ fixation in marine heterotrophic bacteria: dynamics of environment and molecular regulation. *Proc. Natl. Acad. Sci. USA.* 93, 3575-3580.
- Croci, L., Suffredini, E., Cozzi, L., Toti, L., Ottaviani, D., Pruzzo, C., Serratore, P., Fischetti, R., Goffredo, E., Loffredo, G., Mioni, R., 2007. Comparison of different biochemical and molecular methods for the identification of *Vibrio parahaemolyticus*. *J. Appl. Microbiol.* 102, 229-237.
- Crosa, J., 1989. Genetics and molecular biology of siderophore-mediated iron transport in bacteria. *Microb. Rev.* 53, 517-530.
- Dalla, V.L., Zanella, L., Belvedere, P., Colombo, I., 2002. Use of random amplification to developed a PCR detection method for the causative agents of the fish pasteurellosis, *Photobacterium damsela* subsp. piscida (Vibrionaceae). *Aquaculture.* 207, 187-202.
- Dalmin, G., Kathiresan, K., Purushothaman, A., 2001. Effect of probiotics on bacterial population and health status of shrimp in culture pond ecosystem. *Indian. J. Exp. Biol.* 39, 939-942.
- Davies, D.G., Parsek, M.R., Pearson, J.P., Iglewski, B.H., Costerton, J.W., Greenberg, E.P., 1998. The involvement of cell-to-cell signals in the development of a bacterial biofilm. *Science.* 280, 295-298.
- Davis, B.M., Moyer, K.E., Boyd, E.F., Waldor, M.K., 2000b. CTX prophages in classical biotype *Vibrio cholerae*: functional phage genes but dysfunctional phage genomes. *J. Bacteriol.* 182, 6992-6998.

References

- de Carvalho, C.C.C.R., Fernandes, P., 2010. Production of Metabolites as Bacterial Responses to the Marine Environment. *Mar. Drugs*. 8, (3): 705-727
- de Kievit, T.R., Iglewski, B.H., 2000. Bacterial quorum sensing in pathogenic relationships. *Infect. Immun.* 68, 4839-4849.
- de la Peña, L.D., Nakai, T., Muroga, K., 1995. Dynamics of *Vibrio* sp. in organs of orally infected kuruma prawn, *Penaeus japonicus*. *Fish Pathol.* 30, 39–45.
- de la Pena, L.D., Tamaki, T., Momoyama, K., Nakai, T., Muroga, K., 1993. Characteristics of the causative bacterium of vibriosis in the kuruma prawn, *Penaeus japonicus*. *Aquaculture*. 115, 1-12.
- Defoirdt, T., Boon, N., Sorgeloos, P., Verstraete, W., Bossier, P., 2007. Alternatives to antibiotics to control bacterial infections: Luminescent Vibriosis in aquaculture as an example. *Trends Biotechnol.* 25, 472-479.
- Defoirdt, T., Boon, N., Sorgeloos, P., Verstraete, W., Bossier, P., 2008. Quorum sensing and quorum quenching in *Vibrio harveyi*: lessons learned from in vivo work. *The ISME J.* 2, 19-26.
- Denner, E.B.M., Vybiral, D., Fischer, U.R., Velimirov, B., Busse, H.J., 2002. *Vibrio calviensis* sp. nov. a halophilic, facultatively oligotrophic 0.2 µm-filterable marine bacterium. *Int. J. Syst. Evol. Microbiol.* 52, 549-553.
- Di Lorenzo, M., Strok, M., Tolmasky, M.E., Actis, L.A., Farrell, D., Welch, T.J., Corsa, L.M., Wertheimer, A.M., Chen, Q., Salinas, P., Waldbeser, L., Crosa, J.H., 2003. Complete sequence of virulence plasmid pJM1 from the marine fish pathogen *Vibrio anguillarum* strain 775. *J. Bacteriol.* 185, 5822-5830.
- Diggles, B.K., Moss, G.A., Carson, J., Anderson, C.D., 2000. Luminous vibriosis in rock lobster *Jasus verreauxi* (Decapoda: Palinuridae) phyllosoma larvae associated with infection by *Vibrio harveyi*. *Dis. Aquat. Org.* 43, 127-137.
- Dorsch, M., Lane, D., Stackebrandt, E., 1992. Towards a phylogeny of the genus *Vibrio* based on 16S rRNA sequences. *Int. J. Syst. Bacteriol.* 42, 58–63.

References

- *Douglas, J., 1975. Bacteriophages. Chapman and Hall publishers, 4th ed., London. 20-46.
- Doyle, R.J., 2000. Contribution of the hydrophobic effect to microbial infection. *Microbes and Infection*. 2, 391-400.
- Drancourt, M., Bollet, C., Carlouz, A., Martelin, R., Gayral, J. P., Raoult, D., 2000. 16S ribosomal DNA sequence analysis of a large collection of environmental and clinical unidentifiable bacterial isolates. *J. Clin. Microbiol.* 38, 3623–3630.
- *Duguid, J.P., Old, D.C., 1980. Bacterial adherence (receptors and recognition) *In Bacterial adherence*, series B ed. Beachey, E.H. vol.6. New York: Chapman & Hall.
- Dukan, S., Nystrom, T., 1999. Oxidative stress defense and deterioration of growth-arrested *Escherichia coli* cells. *J. Biol. Chem.* 274, 26027-26032.
- Dunlap, P.V., Greenberg, E.P., 1985. Control of *Vibrio fischeri* luminescence gene expression in *Escherichia coli* by cyclic AMP and cyclic AMP receptor protein. *J. Bacteriol.* 164, 45-50.
- Dunlap, P.V., Greenberg, E.P., 1988. Control of *Vibrio fischeri* lux gene transcription by a cyclic AMP receptor protein-LuxR protein regulatory circuit. *J. Bacteriol.* 170, 4040-4046.
- Dunlap, P.V., Ray, J.M., 1989. Requirement for autoinducer in transcriptional negative autoregulation of the *Vibrio fischeri* luxR gene in *Escherichia coli*. *J. Bacteriol.* 171, 3549-3552.
- Engebrecht, J., Silverman, M., 1984. Identification of genes and gene products necessary for bacterial bioluminescence. *Proc. Nat. Acad. Sci. USA* .81, 4154-4158.
- Falkow, S., 1958. Activity of lysine decarboxylase as an aid in the identification of *Salmonella* and *Shigella*. *Am. J. Clin. Pathol.* 19, 598-600.
- FAO, 2003. Health management and biosecurity maintenance in white shrimp (*Penaeus vannamei*) hatcheries in Latin America, FAO Fisheries Technical.
- FAO, 2006. State of the world aquaculture: 2006. FAO Fisheries Technical Paper No. 500. Food and Agricultural Organization, Rome, pp. 134.

References

- FAO, 2009. FAO year book: Fisheries and Aquaculture Statistics, xvi. Food and Agriculture Organization of the United Nations, Rome.
- Farmer III, J.J., 1992. The family of Vibrionaceae. In: Balows, A., Truper, H.G., Dworkin, M., Harder, W., Schleifer, K.H. (Eds.), The prokaryotes. Springer-Verlag, New York. 2938-2951.
- Farmer, J.J. III, Janda, J.M., Brenner, F.W., Cameron, D.N., Birkhead, K.M., 2005. Genus 1. *Vibrio* Pacini 1854, In Bergey's Manual of Systematic Bacteriology, 2nd Ed, Vol.2. The Proteobacteria Part B The Gammaproteobacteria. Brenner, D.J., Krieg, N.R. and Staley, J.T. (Eds) New York: Springer. 494–546.
- Faury, N., Saulnier, D., Thompson, F.L., Gay, M., Swings, J., Roux, F.L., 2004. *Vibrio crassostrea* sp. nov., isolated from the hemolymph of oysters (*Crassostrea gigas*). *Int. J. Syst. Evol. Microbiol.* 54, 2137-2140.
- Felsenstein, J., 1985. Confidence limits on phylogenies: An approach using the bootstrap. *Evolution.* 39, 783-791.
- Finney, D.J., 1952. Probit Analysis. Cambridge, England, Cambridge University Press. 1st Ed
- Finney, D.J., Stevens, W.L., 1948. A table for the calculation of working probits and weights in probit analysis. *Biometrika.* 35, 191-201.
- Flegel, T.W., 1997. Special topic review: Major viral diseases of the black tiger prawn (*Peneaus monodon*) in Thailand. *World Journal of Microbiology and Biotechnology.* 13, 433-442.
- Fluit, A.C., Schmitz, F.J., 1999. Class 1 integrons, gene cassettes, mobility, and epidemiology. *Eur. J. Clin. Microbiol. Infect. Dis.* 18, 761-770.
- Francis C. Yeh., Rong-cai Yang., Tim Boyle. 1999. PopGene Version 1.31. Microsoft Windows based freeware for Population Genetics Analysis.
- Frazier, W.C., 1926. A method for detection of changes in gelatin due to bacteria. *J. Infect. Dis.* 39, 302.

References

- Freeman, J.A., Bassler, B.L., 1999a. Sequence and function of LuxU: a two-component phosphorelay protein that regulates quorum sensing in *Vibrio harveyi*. *J. Bacteriol.* 181, 899-906.
- Freeman, J.A., Bassler, B.L., 1999b. A genetic analysis of the function of LuxO, a two-component response regulator involved in quorum sensing in *Vibrio harveyi*. *Mol. Microbiol.* 31, 665-677.
- Freeman, J.A., Lilley, B.N., Bassler, B.L., 2000. A genetic analysis of the functions of LuxN: a two-component hybrid sensor kinase that regulates quorum sensing in *Vibrio harveyi*. *Mol. Microbiol.* 35, 139-149.
- Fukasawa, S., Nakamura, K., Kamii, A., Ohyama, Y., Osumi, M., 1988a. Purification and properties of a proteinase from a marine luminous bacterium, *Vibrio harveyi* strain FLA-11. *Agric. Biol. Chem.* 52, 435-441.
- Fukasawa, S., Nakamura, K., Kamii, A., Ohyama, Y., Osumi, M., 1988b. Some properties of two proteinases from a luminous bacterium, *Vibrio harveyi* strain FLN-108. *Agric. Biol. Chem.* 52, 3009-3014.
- Fuqua, W.C., Winans, S.C., Greenberg, E.P., 1994. Quorum sensing in bacteria: the LuxR-LuxI family of cell density-responsive transcriptional regulators. *J. Bacteriol.* 176, 269-275.
- Fuqua, W.C., Winans, S.C., Greenberg, E.P., 1996. Census and consensus in bacterial ecosystems: the LuxR-LuxI family of quorum-sensing transcriptional regulators. *Annu. Rev. Microbiol.* 50, 727-751.
- Garcia, T., Otto, K., Kjelleberg, S., Nelson, D.R., 1997. Growth of *Vibrio anguillarum* in salmon intestinal mucus. *Appl. Environ. Microbiol.* 63, 1034-1039.
- Garg, P., Aydanian, A., Smith, D., Morris, J.G., Nair, G.B., Shine, O.C., 2003. Molecular epidemiology of O139 *Vibrio cholerae*: mutation, lateral gene transfer, and founder flush. *Emerg. Infect. Dis.* 9, 810-814.
- Garrity, G.M., Holt, J.G. 2001. The road map to the manual (Boone, D.R. and Castenholz, R.W., Eds.), *Bergey's Manual of Systematic Bacteriology*, Springer, New York.1.

References

- Gatesoupe, F.J., 1999. The use of probiotics in aquaculture. *Aquaculture*. 180, 147-165.
- Gauger, E.J., Gomez-Chiarri, M., 2002. 16S ribosomal DNA sequencing confirms the synonymy of *Vibrio harveyi* and *V. carchariae*. *Dis. Aquat. Org.* 52, 39-46.
- Geiselbrecht, A.D., Herwig, R.P., Deming, J.W., Staley, J.T., 1996. Enumeration and phylogenetic analysis of polycyclic aromatic hydrocarbon-degrading marine bacteria from Puget Sound sediments. *Appl. Environ. Microbiol.* 62, 3344-3349.
- Gestwicki, J.E., Lamanna, A.C., Harshey, R.M., McCarter, L.L., Kiessling, L.L., Adler, J., 2000. Evolutionary conservation of methyl-accepting chemotaxis protein location in Bacteria and Archaea. *J. Bacteriol.* 182, 6499-6502.
- Gevers, D., Vandepoele, K., Simillion, C., Peer, Y.V.D., 2004. Gene duplication and biased functional retention of paralogs in bacterial genomes. *Trends Microbiol.* 12, 148–155.
- Ghatak, S., Agarwal, R.K., Bhilegaonkar, K.N., 2006. Comparative study of cytotoxicity of *Aeromonas* spp. on four different cell lines. *Comp. Immunol. Microbiol. Infect. Dis.* 29, (4): 233-241.
- Gilda D. Lio-Po., Eduardo M. Leño., Ma. Michelle., Peñaranda, D., Annie U. Villa-Franco., Christopher D. Sombito., Nicholas G. Guanzon Jr., 2005. Anti-luminous *Vibrio* factors associated with the ‘green water’ grow-out culture of the tiger shrimp *Penaeus monodon*. *Aquaculture*. 250, 1-2, 1-7.
- Gilles, B., Defoirdt, T., Miyamoto, C., Bossier, P., Calenbergh, S.V., Nelis, H., Coenye, T., 2008. Cinnamaldehyde and cinnamaldehyde derivatives reduce virulence in *Vibrio* spp. by decreasing the DNA-binding activity of the quorum sensing response regulator LuxR. *BMC Microbiol.* 8, 149, 1-14.
- Gilles, B., Celen., S, Baruah., K, Bossier., P, Calenbergh, S.V., Nelis, H., Coenye,T., 2009. AI-2 quorum-sensing inhibitors affect the starvation response and reduce virulence in several *Vibrio* species, most likely by interfering with LuxPQ. *Microbiology*. 155, 4114–4122.
- Goarant, C., Herlin, J., Brizard, R., Marteau, A.L., Martin, C., Martin, B., 2000. Toxic factors of *Vibrio* stains pathogenic to shrimp. *Dis. Aquat. Org.* 40, 101-107.

References

- Gomez Gil, B., Tron-Mayen, L., Roque, A., Turnbull, J.F., Inglis, V., Guerra-Flores, A.L., 1998. Species of *Vibrio* isolated from hepatopancreas, hemolymph and digestive tract of a population of healthy juvenile *Penaeus vannamei*. *Aquaculture*. 163, 1-9.
- Gomez-Gil, B., Roque, A., Turnbull, J.F., 2000. The use and selection of probiotic bacteria for use in the culture of larval aquatic organisms. *Aquaculture*. 191, 259-270.
- Gomez-Gil, B., Thompson, F.L., Thompson, C.C., Swings, J., 2003. *Vibrio pacinii* sp. nov., isolated from cultured aquatic organisms. *Int. J. Syst. Evol. Microbiol.* 53, 1569-1573.
- Gomez-Gil, B., Soto-Rodríguez, S., García-Gasca, A., Roque, A., Vazquez-Juarez, R., Thompson, F.L., Swings, J., 2004a. Molecular identification of *Vibrio harveyi*-related isolates associated with diseased aquatic organisms. *Microbiology*. 150, 1769–1777.
- Gomez-Gil, B., Thompson, F.L., Thompson, C.C., García-Gasca, A., Roque, A., Swings, J., 2004b. *Vibrio hispanicus* sp. nov., isolated from *Artemia* sp. and sea water in Spain. *Int. J. Syst. Evol. Microbiol.* 54, 261-265.
- Gonzalez, S.F., Krug, M.J., Nielsen, M.E., Santos, Y., Call, D.R., 2004. Simultaneous detection of marine fish pathogens by using multiplex PCR and a DNA microarray. *J. Clin. Microbiol.* 1414-1419.
- *Gopalakrishnan, A., Mohindra, V., 2002. Molecular Methods, Tools for Genetic Variability Analysis. (Eds) Mohindra, V, Kuldeep, K.L. and Gopalakrishnan, A. National Bureau of Fish Genetic Resources, Training Manual, Lucknow, U.P, India.
- Goris, J., Suzuki, K., Vos, P.D., Nakase, T., Kersters, K., 1998. Evaluation of a microplate DNA-DNA hybridization method compared with the initial renaturation method. *Can. J. Microbiol.* 44, 1148-1153.
- Gram, L., de Nys, R., Maximilien, R., Givskov, M., Steinberg, P., Kjelleberg, S., 1996. Inhibitory effects of secondary metabolites from the red alga *Delisea pulchra* on swarming motility of *Proteus mirabilis*. *Appl. Environ. Microbiol.* 62, 4284–4287.
- Gram, L., Melchiorson, J., Spanggaard, B., Huber, I., Nielsen, T.F., 1999. Inhibition of *Vibrio anguillarum* by *Pseudomonas fluorescens*

References

- AH2, a possible probiotic treatment of fish. *Appl. Environ. Microbiol.* 65, 969-973.
- Gram, L., Lovold, T., Nielsen, J., Melchiorson, J., Spanggaard, B., 2001. In vitro antagonism of the probiont *Pseudomonas fluorescens* strain AH2 against *Aermonas salmonicida* does not confer protection of salmon against furunculosis. *Aquaculture.* 199, 1-11.
 - Greenberg, E.P., 1997. Quorum sensing in Gram-negative bacteria. *Am. Soc. Microbiol. News.* 63, 371-377.
 - Grisez, L., F.Ollevier, 1995. Comparative Serology of the Marine Fish Pathogen *Vibrio anguillarum*. *Appl. Environ. Microbiol.* 61, 4367-4373.
 - Grisez, L., Chair, M., Sorgeloos, P., Ollevier, F., 1996. Mode of infection and spread of *Vibrio anguillarum* in turbot *Scophthalmus maximus* larvae after oral challenge through live feed. *Dis. Aquat. Org.* 26, 181-187.
 - Groisman, E.A., Ochman, H., 1996. Pathogenicity islands: bacterial evolution in quantum leaps. *Cell.* 87, 791-794.
 - Gurtler, V., Mayall, B.C., 2001. Genomic approaches to typing, taxonomy and evolution of bacterial isolates. *Int. J. Syst. Evol. Microbiol.* 51, 3-16.
 - Hacker, J., Kaper, B.J., 2000. Pathogenicity islands and the evolution of microbes. *Annu. Rev. Microbiol.* 54, 641-679.
 - Hacker, J., Hentschel, U., Dobrindt, U., 2003. Prokaryotic chromosomes and disease. *Science.* 301, 790-793.
 - Hagens, S., Loessner, M.J., 2007. Application of bacteriophages for detection and control of foodborne pathogens. *Appl. Microbiol. Biotechnol.* 7, 1031-1038.
 - Hall, R.M., Stokes, H.W., 1993. Integrons: novel DNA elements which capture genes by site-specific recombination. *Gentica.* 90, 115-132.
 - Hammer, B.K., Bassler, B.L., 2003. Quorum sensing controls biofilm formation in *Vibrio cholerae*. *Mol Microbiol.* 50 101-104.
 - Harris, J.M., 1993. The presence, nature, and role of gut microflora in aquatic invertebrates: a synthesis. *Microb. Ecol.* 25, 195-231.

References

- Harris, L., Owens, L., 1999. Production of exotoxins by two luminous *Vibrio harveyi* strains known to be primary pathogens of *Penaeus monodon* larvae. *Dis. Aquat. Org.* 38, 11-22.
- Harris, J.K., Kelley, S.T., Spiegelman, G.B., Pace, N.R., 2003. The genetic core of the universal ancestor. *Genome Res.* 13, 407-412.
- Hayashi, K., Moriwaki, J., Sawabe, T., Thompson, F.L., Swings, J., Gudkovs, N., Christen, R., Ezura, Y., 2003. *Vibrio superstes* sp. nov., isolated from the gut of Australian abalones *Haliotis laevis* and *Haliotis rubra*. *Int. J. Syst. Evol. Microbiol.* 53, 1813–1817.
- Heidarich, M., Afsharnasab, M., Soltani, M., Dashtyannasab, A., Rajabifar, S., Sheikhzadeh, N., Tamimi, A.H., 2010. Effects of Ergosan and Vibromax to prevent vibriosis and WSSV in *Litopenaeus vannamei*. *J. Fish. Aqu. Sci.* 5, 120-125.
- Heidelberg, J.F., Heidelberg, K.B., Colwell, R.R., 2002a. Bacteria of the α -subclass Proteobacteria associated with zooplankton in Chesapeake Bay. *Appl. Environ. Microbiol.* 68, 5498-5507.
- Heidelberg, J.F., Heidelberg, K.B., Colwell, R.R., 2002b. Seasonality of Chesapeake Bay bacterioplankton species. *Appl. Environ. Microbiol.* 68, 5488-5497.
- Heidelberg, J.F., Eisen, J.A., Nelson, W.C., Clayton, R.A., Gwinn, M.L., Dodson, R.J., Haft, D.H., Hickey, E.K., Peterson, J.D., Richardson, D., Eermolaeva, M.D., Vamathevan, J., Bass, S., Quin, H., Dragoi, I., Seller, P., McDonald, L., Utterback, T., Smith, R.D., Colwell, R.R., Mekalanos, J.J., Venter, J.C., Fraser, C.M., 2000. DNA sequence of both chromosomes of the cholera pathogen *Vibrio cholerae*. *Nature.* 406, 477-484.
- Heithoff, D.M., Mahan, M.J., 2004. *Vibrio cholerae* biofilms: struck between a rock and a hard place. *J. Bacteriol.* 186, 4835-4837.
- Henke, J.M., Bassler, B.L., 2004a. Quorum sensing regulates type III secretion in *Vibrio harveyi* and *Vibrio parahaemolyticus*. *J. Bacteriol.* 186. 3794-3805.
- Henke, J.M., B.L.Bassler, 2004b. Three parallel quorum-sensing systems regulate gene expression in *Vibrio harveyi*. *J. Bacteriol.* 186. 6902-6914.

References

- Henke, J.M., B.LBassler, 2004c. Bacterial social engagements. *Trends Cell Biol.* 14 648-656.
- Herbert, R.A., 1999. Nitrogen cycling in coastal marine ecosystems. *FEMS Microbiol. Rev.* 23, 563-590.
- Hernandez, G., J.Olmos, 2004. Molecular identification of pathogenic and nonpathogenic strains of *Vibrio harveyi* using PCR and RAPD. *Appl. Microbiol. Biotechnol.* 63, 722–727.
- Hernandez-Zarate, G., Olmos-Soto, J., 2006. Identification of bacterial diversity in the oyster *Crassostrea gigas* by fluorescent in situ hybridization and polymerase chain reaction. *J. Appl. Microbiol.* 100, 664-672.
- Hjelm, M., Bergh, O., Riaza, A., Nielsen, J., Melchiorson, J., Jensen, S., Duncan, H., Ahrens, P., Birkbeck, H., Gram, L., 2004a. Selection and identification of autochthonous potential probiotic bacteria from turbot larvae (*Scophthalmus maximus*) rearing units. *Syst. Appl. Microbiol.* 27, 360-371.
- Hjelm, M., Riaza, A., Formoso, F., Melchiorson, J., Gram, L., 2004b. Seasonal Incidence of Autochthonous Antagonistic Roseobacter spp. and Vibrionaceae Strains in a Turbot Larva (*Scophthalmus maximus*) rearing. *System. Appl. Environ. Microbiol.* 70, 7288-7294.
- *Holding, A.J., Collee, J.G., 1971. Routine biochemical tests. In: Norris, J.R., Ribbons, D.W. (Ed.), *Methods in Microbiology* 6A. Academic Press, London, pp. 1-32.
- Holmstrom, K., Gram, L., 2003a. Elucidation of the *Vibrio anguillarum* genetic response to the potential fish probiont *Pseudomonas fluorescens* AH2, using RNA-arbitrarily primed PCR. *J. Bacteriol.* 185, 831-842.
- Holmstrom, K., Graslund, S., Wahlstrom, A., Pongshompoo, S., Bengtsson, B.E., Kautsky, N., 2003b. Antibiotic use in shrimp farming and implications for environmental impacts and human health. *Int. J. Food Sci. Technol.* 38, 255-266.
- Hood, M.A., Winter, P.A., 1997. Attachment of *Vibrio cholerae* under various environmental conditions and to selected substrates. *FEMS Microbiol. Ecol.* 22, 215-223.

References

- Hooper, S.D., Berg, O.G., 2003. On the nature of gene innovation: duplication patterns in microbial genomes. *Mol. Biol. Evol.* 20, 945–954.
- Hopkins, K. L., Desai, M., Frost, J. A., Stanley, J., Logan, J. M., 2004. Fluorescent amplified fragment length polymorphism genotyping of *Campylobacter jejuni* and *Campylobacter coli* strains and its relationship with host specificity, serotyping, and phage typing. *J. Clin. Microbiol.* 42, 229–235
- Hopkins, K.L., Davies, R.H., Threlfall, E.J., 2005. Mechanisms of quinolone resistance in *Escherichia coli* and *Salmonella*: recent developments. *Int. J. Antimicrobial Agents.* 25, 358-373.
- Hoppe, H., 2003. Phosphatase activity in the sea. *Hydrobiologia.* 493, 187-200.
- Holt, J.G., Krieg, N.R., Sneath, P.H.A., Staley, J.T., Williams, S.T., 1994. *Bergey's manual of determinative bacteriology*, 9th Ed. Williams & Wilkins, Baltimore, MD
- Hu, C. Y., Xu, Y.J., Duo, S.W., Li, W.K., Xiang, J.H., Li, M.S., Zhang, R.F., 2010. Preparation of inorganic hollow spheres based on different methods. *J. Chin. Chem. Soc.* 57, (5A), 1091-1098.
- Hueck, C.J., 1998. Type III protein secretion systems in bacterial pathogens of animals and plants. *Microbiol. Mol. Biol. Rev.* 62, 379-433.
- Huys, G., Swings, J., 1999. Evaluation of a fluorescent amplified fragment length polymorphism methodology for the genotypic discrimination of *Aeromonas* taxa. *FEMS Microbiology.* 177, 83-92.
- Hyytiä, E., Björkroth, J., Hielm, S., Korkeala, H., 1999. Characterisation of *Clostridium botulinum* groups I and II by randomly amplified polymorphic DNA analysis and repetitive element sequence-based PCR. *Int. J. Food Microbiol.* 48, 179–189.
- Irianto, A., Austin, B., 2002a. Probiotics in aquaculture. *J. Fish. Dis.* 25, 633-642.
- Irianto, A., Austin, B., 2002b. Use of probiotics to control furunculosis in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *J. Fish. Dis.* 25, 333-342.

References

- Janda, J.M., Poweres, C., Bryant, R.G., Abbott, S.L., 1988. Current perspectives on the epidemiology and pathogenesis of clinically significant *Vibrio* spp. *Clin. Microbiol. Rev.* 1, 245-267.
- Jayaprakash, N.S., Pai, S.S., Anas, A., Preetha, R., Philip, R., Singh, I.S.B., 2005. A marine *Micrococcus* MCCB104 antagonistic to vibrios in prawn larval rearing systems. *Dis. Aquat. Org.* 68, 39-45.
- Jefferies C.D., Holtman D.E., Guse D.G., 1957. Rapid method for the determining the activity of microorganisms on nucleic acids. *J. Bacteriol.* 73, 590.
- Jefferson, K.K., Pier, D.B., Goldmann, D.A., Pier, G.B., 2004. The teicoplanin-associated locus regulator (TcaR) and the intercellular adhesin locus regulator (IcaR) are transcriptional inhibitors of the *ica* locus in *Staphylococcus aureus*. *J. Bacteriol.* 186, 2449–2456.
- Jenkins, D.E., Chaisson, S.A., Martin, A., 1990. Starvation-induced cross protection against osmotic challenge in *Escherichia coli*. *J. Bacteriol.* 172, 2779-2781.
- Jiang, S., Louis, C.V., Choopun, N., Sharma, A., Huq, A., Colwell, R.R. 2000a. Genetic diversity of *Vibrio cholerae* in Chesapeake Bay determined by amplified fragment length polymorphism fingerprinting. *Appl. Environ. Microbiol.* 66, 140-147.
- Jiang, S., Matte, C.M., Matte, G., Huq, A., Colwell, R.R., 2000b. Genetic diversity of clinical and environmental isolates of *Vibrio cholerae* determined by amplified fragment length polymorphism fingerprinting. *Appl. Environ. Microbiol.* 66, 148-153.
- Johan Vandenberghe, Fabiano L. Thompson, Bruno Gomez-Gil, Jean Swings, 2003. Phenotypic diversity amongst *Vibrio* isolates from marine aquaculture systems. *Aquaculture.* 219, 9 -20.
- Johnsen, K., Andersen, S., Jacobsen, C.S., 1996. Phenotypic and genotypic characterization of phenanthrene degrading fluorescent *Pseudomonas* biovars. *Appl. Environ. Microbiol.* 62, 3818-3825.
- Johnson, F., I.Shunk, 1936. An interesting new species of luminous bacteria. *J Bacteriol.* 31, 585-592.
- Jonson, G., Svennerholm, A.M., Holmgren, J., 1989. *Vibrio cholerae* expresses cell surface antigens during intestinal infection

References

- which are not expressed during in vitro Culture. *Infect. Immun.* 57, (6): 1809- 1815.
- Joseph, S.W., Colwell, R.R., Kaper, J.B., 1982. *Vibrio parahaemolyticus* and related halophilic vibrios. *Crit. Rev. Microbiol.* 10, 77-124.
 - Kang, J.H., Lee, J.H., Park, J.H., Huh, S.H., Kong, I.S., 1998. Cloning and identification of a phospholipase gene from *Vibrio mimicus*. *Biochimica et Biophysica Acta (BBA) - Lipids and Lipid Metabolism.* 1394, (1): 85-89.
 - *Kaper, J.B., Hacker, J. 1999. Pathogenicity islands and other mobile virulence elements. ASM Press, Washington, D.C.
 - Karunasagar, I., Pai, R., Malathi, G.R., Karunasagar, I., 1994. Mass mortality of *Penaeus monodon* larvae due to antibiotic resistant *Vibrio harveyi* infection. *Aquaculture.* 128, 203-209.
 - Kaufman, M.R., Shaw, C.E., Jones, I.D., Taylor, R.K., 1993. Biogenesis and regulation of the *V.cholerae* toxin-coregulated pilus: analogies to their virulence factor secretory systems. *Gene.* 126, 43-49.
 - Kaysner, C.A., Abeyta, C., Wekell, M.M., Jr. DePaola, A., Stott, Jr. R.F., Leitch, J.M., 1987. Virulent strains of *Vibrio vulnificus* isolated from estuaries of the United States west coast. *Appl. Environ. Microbiol.* 53, 1349-1351.
 - Kelly, M.T., 1982. Effect of temperature and salinity on *Vibrio (Beneckea) vulnificus* occurrence in a Gulf Coast environment. *Appl. Environ. Microbiol.* 44, 820-824.
 - Kesarcodi-Watson, A., Kaspar, H., Lategan, M. J., 2008. Probiotics in aquaculture: The need, principles and mechanisms of action and screening processes. *Aquaculture.* 274, 1-14.
 - Khemayan, K., Pasharawipas, T., Puiprom, O., Sriurairatana, S., Suthienkul, O., Flegel, T.W., 2006. Unstable lysogeny and pseudolysogeny in *Vibrio harveyi* Siphovirus-like phage 1. *Appl. Environ. Microbiol.* 72, 1355-1363.
 - Kierek, K., Watnick, P.I., 2003a. Environmental Determinants of *Vibrio cholerae* Biofilm Development. *Appl. Environ. Microbiol.* 69, 5079-5088.

References

- Kim, S.R., Nonaka, L., Suzuki, S., 2004a. Occurrence of tetracycline resistance genes tet(M) and tet(S) in bacteria from marine aquaculture sites. *FEMS Microbiol. Letts.* 237, 147-156.
- Kim, Y.H., Pak, K., Pothuluri, J.V., Cerniglia, C.E., 2004b. Mineralization of erythromycin A in aquaculture sediments. *FEMS Microbiol. Letts.* 234, 169-175.
- Kirkup, B. C., Jr, Chang, L. A., Chang, S., Gevers, D., Martin F Polz, M. F., 2010. *Vibrio* chromosomes share common history. *BMC Microbiology.* 10, (137): 1- 13.
- Kita-Tsukamoto, K., Oyaizu, H., Nanba, K., Simidu, U., 1993. Phylogenetic relationships of marine bacteria, mainly members of the family *Vibrionaceae*, determined on the basis of 16S rRNA sequences. *Int. J. Syst. Bacteriol.* 43, 8-19.
- Klingler, J.M., Stowe, R.P., Obenhuber, D.C., Groves, T.O., Mishra, S.K., Pierson, D.L. 1992. Evaluation of the Biolog automated microbial identification system. *Appl. Environ. Microbiol.* 58, 2089-2092.
- *Koch, R., 1893. *J. Hyg. Inf.* 14, 319-333.
- Kong, R.Y., Lee, S.K., Law, T.W., Law, S.H., Wu, R.S., 2002. Rapid detection of six types of bacterial pathogens in marine wate by multiplex PCR. *Water Res.* 36, 2802-2812.
- Kolter, R., Siegele, D.A., Tormo, A. 1993. The stationary phase of the bacterial life cycle. *Annu. Rev. Microbiol.* 47, 855-874.
- Kovacs, N., 1956. Identification of *Pseudomonas pyocyanea* by the oxidase reaction. *Nature.* 128, 703.
- Krieg, N.R., Holt, J.G., 1984. *Bergey's Manual of Systematic Bacteriology.* Williams and Wilkins, Baltimore, USA, 140- 219.
- Krumperman, P.H., 1983. Multiple antibiotic resistance indexing of *Escherichia coli* to identify high-risk sources of fecal contamination of foods. *Appl. Environ. Microbiol.* 46, 165-170.
- Kühn, I., Allestam, G., Stenstrom, T.A., Mollby, R., 1991. Biochemical fingerprinting of water coliform bacteria, a new method

References

- for measuring phenotypic diversity and for comparing different bacterial populations. *Appl. Environ. Microbiol.* 57, 3171-3177.
- Kumar, R., Mukherjee, S.C., Prasad, K.P., Pal, A.K., 2006. Evaluation of *Bacillus subtilis* as a probiotic to Indian major carp *Labeo rohita* (Ham.). *Aquacult. Res.* 37, 1215-1221.
 - Kurland, C.G., Canback, B., Berg, O.G., 2003. Horizontal gene transfer a critical view. *PROC. NATL. ACAD. SCI. USA.* 100, 9658-9662.
 - Lachica, R.V., Zink, D.L., 1984. Plasmid-associated cell surface charge and hydrophobicity of *Yersinia enterocolitica*. *Infect. Immun.* 44, 540-543.
 - Lambert, C., Nicolas, J., Cilia, V., Corre, S., 1998. *Vibrio pectenecida* sp. nov., a pathogen of scallop (*Pecten maximus*) larvae. *Int. J. Syst. Bacteriol.* 48, 481-487.
 - Lambert, M. A., Hickman-Brenner, F. W., Farmer, J. J., III., Moss, C. W., 1983. Differentiation of Vibrionaceae species by their cellular fatty acid composition. *Int. J. Syst. Bacteriol.* 33, 777-792.
 - Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature.* (London). 227: 680-685.
 - Lane, D.J., Pace, B., Olsen, G.J., Stahl, D.A., Sogin, M.L., 1985. Rapid determination of 16S ribosomal RNA sequences for phylogenetic analyses. *Proc. Natl. Acad. Sci. USA.* 82, 6955-6959.
 - Lange, R., Hengge-Aronis, R., 1991. Identification of a central regulator of stationary-phase gene expression in *Escherichia coli*. *Mol. Microbiol.* 5, 49-59.
 - Larsen, M.H., Boesen, H.T., 2001. Role of flagellum and chemotactic motility of *Vibrio anguillarum* for phagocytosis by and intracellular survival in fish macrophages. *FEMS Microbiol. Letts.* 203, 149-152.
 - Larsen, M.H., Blackburn, N., Larsen, J.L., Oslen, J.E., 2004. Influences of temperature, salinity and starvation on the motility and chemotactic response of *Vibrio anguillarum*. *Microbiology.* 150, 1283-1290.

References

- Lauriano, C.M., Ghosh, C., Correa, N.E., Klose, K.E., 2004. The sodium-driven flagellar motor controls exopolysaccharide expression in *Vibrio cholerae*. *J. Bacteriol.* 186, 4846-4874.
- Lavilla-Pitogo, C.R., Baticados, M.C.L., Cruz-Lacierda, E.R., de la Pena, L.D., 1990. Occurrence of luminous bacterial disease of *Penaeus monodon* larvae in the Philippines. *Aquaculture.* 91, 1-13.
- Lavilla-Pitogo, C.R., Leanño, E.M., Paner, M.G., 1998. Mortalities of pond-cultured juvenile shrimp, *Penaeus monodon*, associated with dominance of luminescent vibrios in the rearing environment. *Aquaculture.* 164, 337-349.
- Le Chevalier, P., Boulay, C.L., Paillard, C. 2003. Characterization by restriction fragment length polymorphism and plasmid profiling of *Vibrio tapetis* strains. *J. Basic. Microbiol.* 43, 414-422.
- Le Roux, F., Gay, M., Lambert, C., Nicolas, J.I., Gouy, M., Berthe, F. 2004. Phylogenetic study and identification of *Vibrio splendidus*-related strains based on gyrB gene sequences. *Dis. Aquat. Org.* 58, 143-150.
- Lee, K.H., Ruby, E.G. 1995. Symbiotic role of the viable but nonculturable state of *Vibrio fischeri* in Hawaiian coastal seawater. *Appl. Environ. Microbiol.* 61, 278-283.
- Lee, K. K., Yii, K. C., 1996. A comparison of three methods for assaying hydrophobicity of pathogenic vibrios. *Letts. Appl. Microbiol.* 23, (5): 343- 346.
- Lee, K.K., Chen, Y.L., Liu, P.C., 1999. Hemostasis of tiger prawn *Penaeus monodon* affected by *Vibrio harveyi*, extracellular products and a toxic cysteine protease. *Blood Cells, Molecules and Diseases.* 25, 180-192.
- Lee, K.K., Chen, F.R., Yu, R.R., Yang, T.I., Liu, P.C., 1997a. Effects of extracellular products of *Vibrio alginolyticus* on penaeid shrimp plasma components. *Letst. Appl. Microbiol.* 25, 98-100.
- Lee, K.K., Liu, P.C., Kou, G.H., Chen, S.N., 1997b. Passive immunization of the tiger prawn, *Penaeus monodon*, using rabbit antisera to *Vibrio harveyi*. *Letts. Appl. Microbiol.* 25, 34-37.

References

- Lee, K.K., Yu, S.R., Chen, F.R., Yang, T.I., Liu, P.C., 1996. Virulence of *Vibrio alginolyticus* isolated from diseased tiger prawn, *Penaeus monodon*. *Curr. Microbiol.* 32, 229-231.
- Lee, M.J., Jeong, D.Y., Kim, W.S., Kim, H.D., Kim, C.H., Park, W.W., Park, Y.H., Kim, K.S., Kim, H.M., Kim, D.S., 2000. A tetrodotoxin-producing *Vibrio* strain, LM-1, from the puffer fish *Fugu vermicularis radiatus*. *Appl. Environ. Microbiol.* 66, 1698-1701.
- Lee, S.H., Camilli, A., 2000. Novel approaches to monitor bacterial gene expression in infected tissue and host. *Curr. Opin. Microbiol.* 3, 97-101.
- Lenz, D.H., Mok, K.C., Lilley, B.N., Kulkarni, R.V., Wingreen, N.S., 2004. The small RNA chaperone Hfq and multiple small RNAs control quorum sensing in *Vibrio harveyi* and *Vibrio cholerae*. *Cell.* 118 69-82.
- Leung, K.Y., Stevenson, R.M.W. 1988. Characteristics and Distribution of Extracellular Proteases from *Aeromonas hydrophila*. *J. Gen. Microbiol.* 134, 151-160.
- Leverstein-van-Hall, M.A., Paauw, A., Box, A.T.A., Blok, H.E.M., Verhoef, J., Fluit, A.C., 2002. Presence of integron-associated resistance in the community is widespread and contributes to multidrug resistance in the hospital. *J. Clin. Microbiol.* 40, 3038-3040.
- Li, N., Yang, Z., Bai, J., Fu, X., Liu, L., Shi, C., Wu, S., 2010. A shared antigen among *Vibrio* species: Outer membrane protein - OmpK as a versatile Vibriosis vaccine candidate in Orange-spotted grouper (*Epinephelus coioides*). *Fish and Shellfish Immunology.* 28, 952-956.
- Liebert, C.A., Hall, R.M., Summers, A.O., 1999. Transposon Tn21, Flagship of the Floating Genome. *Microbiology and Molecular Biology Reviews.* 63, 507-522.
- Lightner, D.V., 1996. A Handbook of Shrimp Pathology and Diagnostic Procedures for Diseases of Cultured Penaeid Shrimp. World Aquaculture Society, Baton Rouge, LA, USA. 1st Ed. 304.
- Lightner, D.V., Redman, R.M., Bell, T.A., 1983. Infectious hypodermal and hematopoietic necrosis, a newly recognized virus disease of penaeid shrimp. *J. Invert. Pathol.* 42, 62-70.

References

- Lightner, D.V., Redman R.M., 1998. Shrimp diseases and current diagnostic methods. *Aquaculture*. 164,201-220.
- Lilley, B.N., Bassler, B.L., 2000. Regulation of quorum sensing in *Vibrio harveyi* by LuxO and sigma-54. *Mol. Microbiol.* 36, 940-945.
- Lin, Z., Kumagai, K., Baba, K., Mekalanos, J.J., Nishibuchi, M., 1993. *Vibrio parahaemolyticus* has a homolog of the *Vibrio cholerae* toxRS operon that mediates environmentally induced regulation of the thermostable direct hemolysin gene. *J. Bacteriol.* 175, 3844-3855.
- Lingappa, Y., Lockwood, J.L., 1961. Chitin medium for selective isolation and culture of actinomycetes. *Nature*. 189, 158.
- Liston, J., 1954. A group of luminous and nonluminous bacteria from the intestine of flatfish. *J. Gen. Microbiol.* 12, 1.
- Liu, P.C., Lee, K.K., Yii, K.C., Kou, G.S., Chen, S.N., 1996a. Isolation of *Vibrio harveyi* from diseased kuruma prawns *Penaeus japonicus*. *Curr. Microbiol.* 33, 129-132.
- Liu, P.C., Lee, K.K., Chen, S.N., 1996b. Pathogenicity of different isolates of *Vibrio harveyi* in tiger prawn, *Penaeus monodon*. *Letts. Appl. Microbiol.* 22, 413-416.
- Liu, P.C., Lee, K.K., Yii, K.C., Kou, G.H., Chen, S.N., 1997. Purification and characterization of a cysteine protease produced by pathogenic luminous *Vibrio harveyi*. *Curr. Microbiol.* 35, 32-39.
- Liu, P.C., Lee, K.K., 1999. Cysteine protease is a major exotoxin of pathogenic luminous *Vibrio harveyi* in the tiger prawn, *Penaeus monodon*. *Letts. Appl. Microbiol.* 28, 428-430.
- Liu, P.C., Lin, J.Y., Chuang, W.H., Lee, K.K., 2004. Isolation and characterization of pathogenic *Vibrio harveyi* (*V. carchariae*) from the farmed marine cobia fish *Rachycentron canadum* L. with gastroenteritis syndrome. *World J. Microbiol. Biotechnol.* 20, 495-499.
- Logothetis, P.N., Austin, B., 1996. Variations in antigenicity of *Aeromonas hydrophila* strains in rainbow trout (*Oncorhynchus mykiss*, Walbaum): an association with surface characteristics. *Fish Shellfish Immunolo.* 6, 47-55.

References

- Lu, S., Liu, B., Cao, J., Zhou, B., E. Levin, R., 2006. Incidence and enumeration of *Vibrio parahaemolyticus* in shellfish from two retail sources and the genetic diversity of isolates as determined by RAPD-PCR analysis. *Food Biotechnology*. 20, 193-209.
- Ludwig, W., Klenk, H.P., 2001. Overview: a phylogenetic backbone and taxonomic frame work for prokaryotic systematics. *In Bergey's Manual of Systematic Bacteriology*. 2 Ed. Springer-Verlag, New York, N.Y. 49-65.
- Lunder, T., Sørum, H., Holstad, G., Steigerwalt, A.G., Mowinckel, P., Brenner, D.J., 2000. Phenotypic and genotypic characterization of *Vibrio viscosus* sp. nov. and *Vibrio wodanis* sp. nov. isolated from Atlantic salmon (*Salmo salar*) with 'winter ulcer'. *Int. J. Syst. Evol. Microbiol.* 50, 427-450.
- Macián, M.C., Garay, E., Pujalte, M.J., 1996. The arginine dihydrolase (ADH) system in the identification of some marine *Vibrio* species. *Syst. Appl. Microbiol. Biotechnol.* 19, 451-456.
- Macián, M.C., Garay, E., González-Candela, F., Pujalte, M.J., Anzar, R., 2000. Ribotyping of *Vibrio* populations associated with cultured oysters (*Ostrea edulis*). *Syst. Appl. Microbiol.* 23, 409-417.
- Macián, M.C., Ludwig, W., Aznar, R., Grimont, P.A., Schleifer, K.H., Garay, E., Pujalte, M.J., 2001a. *Vibrio lentus* sp. nov., isolated from Mediterranean oysters. *Int. J. Syst. Evol. Microbiol.* 51, 1449-1456.
- Macián, M.C., Ludwig, W., Schleifer, K.H., Pujalte, M.J., Garay, E., 2001b. *Vibrio agarivorans* sp. nov., a novel agarolytic marine bacterium. *Int. J. Syst. Evol. Microbiol.* 51, 2031-2036.
- Maeda, T., Takada, N., Furushita, M., Shiba, T. 2000. Structural variation in the 16S-23S rRNA Intergenic spacers of *Vibrio parahaemolyticus*. *FEMS Microbiology Letters*. 192, (1): 73-77.
- Magnusson, K.E., Davies J., Gundstrom, T., Kihlstrom, E., Normark, S., 1980. Surface charge and hydrophobicity of *Salmonella*, *E. coli* and *Gonococci* in relation to their tendency to associate with animal cells. *Sc and J Infect Dis* 24, 135-140.
- Magariños, B., Osorio, C. R., Toranzo, A. E., Romalde, J. L., 1997. Applicability of ribotyping for intraspecific classification and

References

- epidemiological studies of *Photobacterium damsela* subsp. *piscicida*. *Syst. Appl. Microbiol.* 20: 634- 639.
- Maiden, M.C., Bygraves, J.A., Feil, E., Morelli, G., Russell, J.E., Urwin, R., Zhang, Q., Zhou, J., Zurth, K., Caugant, D.A., Feavers, I.M., Achtman, M., Spratt, B.G., 1998. Multilocus sequence typing: a portable approach to the identification of clones within populations of pathogenic microorganisms. *Proc. Natl. Acad. Sci. USA* 95, 3140-3145.
 - Maiti, B., Shekar, M., Khushiramani, R., Karunasagar, I., Karunasagar, I., 2009. Evaluation of RAPD-PCR and protein profile analysis to differentiate *Vibrio harveyi* strains prevalent along the southwest coast of India. *J. Genetics.* 88, 273-279.
 - Majtanova, L., Majtan, V., 2006. Phage types and virulence markers of clinical isolates of *Salmonella enteritidis*. *Epidemiol. Mikrobiol. Imunol.* 55, 87-91.
 - Makino, K., Oshima, K., Kurokawa, K., Yokoyama, K., Uda, T., Tagomori, K., Iijima, Y., Najima, M., Nakano, M., Yamashita, A., Kubota, Y., Kimura, S., Yasunaga, T., Honda, T., Shinagawa, H., Hottori, M., Iida, T., 2003. Genome sequence of *Vibrio parahaemolyticus*: a pathogenic mechanism distinct from that of *V.cholerae*. *Lancet.* 361, 743-749.
 - Manefield, M., de Nys, R., Kumar, N., Read, R., Givskov, M., Steinberg, P., Kjelleberg, S., 1999. Evidence that halogenated furanones from *Delisea pulchra* inhibit acylated homoserine lactone (AHL)-mediated gene expression by displacing the AHL signal from its receptor protein. *Microbiology.* 145, 283-291.
 - Manefield, M., Harris, L., Rice, S.A., de Nys, R., Kjelleberg, S., 2000. Inhibition of luminescence and virulence in the black tiger prawn (*Penaeus monodon*) pathogen *Vibrio harveyi* by intercellular signal antagonists. *Appl. Environ. Microbiol.* 66, 2079-2084.
 - Marchesi, J., Sato, T., Weightman, A., Martin, T., Fry, J., Hiom, S., Wade, W., 1998. Design and evaluation of useful bacterium-specific PCR primers that amplify genes coding for bacterial 16S rRNA. *Appl. Environ. Microbiol.* 64, 795-799.
 - Marco-Noales, E., Milan, M., Fouz, B., Sanjuan, E., Amaro, C., 2001. Transmission to eels, portals of entry, and putative reservoirs of *Vibrio vulnificus* serovar E (biotype 2). *Appl. Environ. Microbiol.* 67, 4717-4725.

References

- Marques, A., Dinh, T., Ioakeimidis, C., Huys, G., Swings, J., Verstraete, W., Dhont, J., Sorgeloos, P., Bossier, P., 2005. Effects of bacteria on *Artemia franciscana* cultured in different gnotobiotic environments. *Appl. Environ. Microbiol.* 71, 4307-4317.
- Martí'nez-Picado, J., Alsina, M., Blanch, A.R., Cerda`, M., Jofre, J., 1996. Species-specific detection of *Vibrio anguillarum* in marine aquaculture environments by selective culture and DNA hybridization. *Appl. Environ. Microbiol.* 62, 443-449.
- Martinez, E., Cruz, F.D.L., 1990. Genetic elements involved in Tn21 site-specific integration, a novel mechanism for the dissemination of antibiotic resistance genes. *EMBO J.* 9, 1275-1281.
- Martinez-Murcia, A., Benlloch, J.S., Collins, M.D., 1992. Phylogenetic Interrelationships of Members of the Genera *Aeromonas* and *Plesiomonas* as determined by 16S Ribosomal DNA sequencing lack of congruence with results of DNA-DNA Hybridizations. *Int. J. Syst. Bacteriol.* 42, 412-421.
- Martinez-Murcia, A.J., Borrell, N., Figueras, M.J., 2000. Typing of clinical and environmental *Aeromonas veronii* strains based on the 16S-23S rDNA spacers. *FEMS Immun. Med. Microbiol.* 28, 225-232.
- Martin-Kearley, J., Gow, J.A., Peloquin, M., Green, C.W., 1994. Numerical analysis and application of Random Amplified Polymorphic DNA polymerase chain reaction to differentiation of vibrio strain from a seasonally cold ocean. *Can. J. Microbiol.* 40, 446-455.
- Marwick, J.D., Wright, P.C., Burgess, J.G., 1999. Bioprocess intensification for production of novel marine bacterial antibiotics through bioreactor operation and design. *Mar. Biotechnol.* 1, 495-507.
- Matz, C., Deines, P., Jurgens, K., 2002b. Phenotypic variation in *Pseudomonas* sp. CM10 determines microcolony formation and survival under protozoan grazing. *FEMS Microbiol. Ecol.* 39, 57-65.
- Maugeri, T.L., Caccamo, D., Gugliandolo, C., 2000. Potentially pathogenic vibrios in brackish waters and mussels. *J. Appl. Microbiol.* 89, 261-266.

References

- Mazel, D., 2006. Integrons, agents of bacterial evolution. *Nat. Rev. Microbiol.* 4, 608-620.
- Mazel, D., Dychinco, B., Webb, V.A., Davies, J., 1998. A distinctive class of integron in the *Vibrio cholerae* genome. *Science.* 280, 605-608.
- McCall, J.O., Sizemore, R.K., 1979. Description of a bacteriocinogenic plasmid in *Beneckeia harveyi*. *Appl. Environ. Microbiol.* 38, 974-979.
- McCarter, L.L., 1998. *OpaR*, a homolog of *Vibrio harveyi LuxR*, controls opacity of *Vibrio parahaemolyticus*. *J. Bacteriol.* 180, 3166–3173.
- McCarter, L.L., 2001. Polar flagellar motility of Vibrionaceae. *Microbiol. Molbio. Rev.* 65, 3, 445-462.
- McDougald, D., Rice, S.A., Kjelleberg, S., 2003. The marine pathogen *Vibrio vulnificus* encodes a putative homologue of the *Vibrio harveyi* regulatory gene, *luxR*: a genetic and phylogenetic comparison. *Gene.* 248, 213–221.
- McDougald, D., Rice, S.A., Kjelleberg, S., 2001. SmcR-dependent regulation of adaptive response in *Vibrio vulnificus*. *J. Bacteriol.* 183, 758-762.
- McFall-Ngai, M.J., 1999. Consequences of evolving with bacterial symbionts: Insights from the Squid-Vibrio associations. *Annu. Rev. Ecol. Syst.* 30, 235-256.
- McFall-Ngai, M.J., 2002. Unseen forces: the influence of bacteria on animal development. *Dev. Biol.* 242, 1–14.
- McGee, D., P.Horstedt, D.L.Milton, 1996. Identification and characterization of additional flagellin genes from *Vibrio anguillarum*. *J.Bacteriol.* 178, 5188-5198.
- McGrath, S., Fitzgerald, G.F., van Sinderen, D., 2004. The impact of bacteriophage genomics. *Curr. Opin. Biotechnol.* 15, 94- 99.
- Meighen, E.A., 1991. Molecular biology of bacterial bioluminescence. *Microbiol. Mol. Biol. Rev.* 55, (1): 123-142.
- Meighen, E.A., 1994. Genetics of bacterial bioluminescence. *Ann. Rev. Genet.* 28, 117-139.

References

- Messier, N., Roy, P.H., 2001. Integron integrases possess a unique additional domain necessary for activity. *J. Bacteriol.* 183, 6699-6706.
- Michel Janda, J., Oshiro, S., Abbott, L., Duffey, P.S., 1987. Virulence markers of Mesophilic Aeromonads: Association of the autoagglutination phenomenon with mouse pathogenicity and the presence of a Peripheral Cell-Associated Layer. *Infect. Immun.* 55, 3070-3077.
- Miller, V.L., Taylor, R.K., Mekalanos, J.J., 1987. Cholerae toxin transcriptional activator ToxR is a transmembrane DNA binding protein. *Cell.* 48, 271-279.
- Miller, V.L., Mekalanos, J.J., 1988. A novel suicide vector and its use in construction of insertion mutations: osmoregulation of outer membrane proteins and virulence determinants in *Vibrio cholerae* requires *toxR*. *J. Bacteriol.* 170, 2572-2583.
- Miller, J.M., Rhoden, D.L., 1991. Preliminary evaluation of Biolog, a carbon source utilization method for bacterial identification. *J. Clin. Microbiol.* 29, 1143-1147.
- Miller, M.B., B.L.Bassler, 2001. Quorum sensing in bacteria. *Annu. Rev. Microbiol.* 55, 165-199.
- Milton, D.L., O'Toole, R., Hörstedt, P., Wolf-Watz, H., 1996. Flagellin A is essential for the virulence of *Vibrio anguillarum*. *J. Bacteriol.*, 1310-1319.
- Milton, D.L., Hardman, A., Camara, M., Chhabra, S.R., Bycroft, B.W., Stewart, G.S.A.B., Williams, P., 1997. Quorum sensing in *Vibrio anguillarum*: characterization of the *vanI/vanR* locus and identification of the autoinducer N-(3-oxodecanoyl)-L-homoserine lactone. *J. Bacteriol.* 179, 3004-3012.
- Milton, D.L., 2006. Quorum sensing in vibrios: Complexity for diversification. *Int. J. Medi. Microbiol.* 296, 61-71.
- Miranda, C.D., Zemelman, R., 2002. Antimicrobial multiresistance in bacteria isolated from freshwater Chilean salmon farms. *Sci. Total Environ.* 293, 207-218.

References

- Miranda, C.D., Kehrenberg, C., Ulep, C., Schwarz, S., Roberts, M.C., 2003. Diversity of tetracycline resistance genes in bacteria from Chilean salmon farms. *Antimicrobial Agents & Chemotherapy*. 47, 883-888.
- Miyamoto, C.M., Boylan, M., Graham, A.F., Meighen, E.A., 1988. Organization of the lux structural genes of *Vibrio harveyi*. *J. Biol. Chem.* 263, 13393-13399.
- Mok, K.C., Wingreen, N.S., Bassler, B.L., 2003. *Vibrio harveyi* quorum sensing: a coincidence detector for two autoinducers controls gene expression. *EMBO. J.* 22, 870-881.
- Moller, V., 1955. Simplified tests for some aminoacids decarboxylases and for the arginine dihydrolase system. *Acta. Pathol. Microbiol. Scand.* 36, 2. 158-172.
- Mollet, C., Drancourt, M., Raoult, D., 1997. *rpoB* sequence analysis as a novel basis for bacterial identification. *Mol. Microbiol.* 26, 1005-1011.
- Montero, A.B., Austin, B., 1999. Characterization of extracellular products from an isolate of *Vibrio harveyi* recovered from diseased post-larval *Penaeus vannamei* (Bonne). *J. Fish. Dis.* 22, 377-386.
- Montes, M., Pe´rez, M.J., Nieto, T.P., 1999. Numerical taxonomy of Gram-negative, facultative anaerobic bacteria isolated from skin of turbot (*Scophthalmus maximus*) and surrounding water. *Syst. Appl. Microbiol.* 22, 604-618.
- Montgomery, M.T., Kirchman, D.L., 1993. Role of chitin binding proteins in the specific attachment of the marine bacterium *Vibrio harveyi* to chitin. *Appl. Environ. Microbiol.* 59, 373-379.
- Montgomery, M.T., Kirchman, D.L., 1994. Induction of chitin-binding proteins during the specific attachment of the marine bacterium *Vibrio harveyi* to chitin. *Appl. Environ. Microbiol.* 60, 4284-4288.
- Moreno, C., Romero, J., Espejo, R.T., 2002. Polymorphism in repeated 16S rRNA genes is a common property of type strains and environmental isolates of the genus *Vibrio*. *Microbiology.* 148, 1233-1239.

References

- Moriarty, D.J.W., 1997. The role of microorganisms in aquaculture ponds. *Aquaculture*. 151, 333-349.
- Moriarty, D.J.W., 1998. Control of luminous *Vibrio* species in penaeid aquaculture ponds. *Aquaculture*. 164, 351-358.
- Mosmann, T., 1983. Rapid colorimetric assay for cellular growth and survival, application to proliferation and cytotoxicity assays. *J. Immunol. Methd.* 65, 55-63.
- Munro, J., Oakey, J., Bromage, E., Owens, L., 2003. Experimental bacteriophage mediated virulence in strains of *Vibrio harveyi*. *Dis. Aquat. Org.* 54, 187-194.
- Muroga, K., M.Higashi, Keitoku, H., 1987. The isolation of intestinal microflora of farmed red seabream (*Pagrus major*) and black seabream (*Acanthopagrus schlegeli*) at larval and juvenile stages. *Aquaculture*. 65, 79-88.
- Muroga, K., 2001. Viral and bacterial diseases of marine fish and shellfish in Japanese hatcheries. *Aquaculture*. 202, 23-44.
- Mutharia, L.M., Amor, P.A., 2002. Monoclonal antibodies against *Vibrio anguillarum* O2 and *Vibrio ordalii* identify antigenic differences in lipopolysaccharide O-antigens. *FEMS Microbiol. Lett.* 123, 289-298.
- Nagpal, M.L., Fox, K.F., Fox, A., 1998. Utility of 16S-23SrRNA spacer region methodology: how similar are interspacer region within a genome and between strains of closely related organisms? *J. Microbiol. Methods.* 33, 211-219.
- Nakae, T., Nikaido, H., 1975. Outer membrane as a diffusion barrier in *Salmonella typhimurium*: penetration of oligo- and polysaccharides into isolated outer membrane vesicles and cells with degraded peptidoglycan layer. *J. Biol. Chem.* 250, 7359-7365.
- Nakae, T., 1976a. Identification of the outer membrane protein of *Escherichia coli* that produces transmembrane channels in reconstituted vesicle membranes. *Biochem. Biophys. Res. Communi.* 71, 877-884.
- Nakae, T., 1976b. Outer membrane of *Salmonella*. Isolation of protein complex that produces transmembrane channels. *J. Biol. Chemi.* 251, 2176-2178.

References

- Nakayama, T., Nomura, N., Matsumura, M., 2005. Analysis of the relationship between luminescence and toxicity of *Vibrio carchariae* pathogenic to shrimp. *Fish. Sci.* 71, 1236-1242.
- Nakayama, T., Ito, E., Nomura, N., Nomura, N., Matsumura, M., 2006a. Comparison of *Vibrio harveyi* strains isolated from shrimp farms and from culture collection in terms of toxicity and antibiotic resistance. *FEMS Microbiol. Lett.* 258, 194-199.
- Nakayama, T., Nomura, N., Matsumura, M., 2006b. Study on the relationship of protease production and luminescence in *Vibrio harveyi*. *J. Appl. Microbiol.* 101, 200-205.
- Namdari H., and Bottone, E.J., 1988. Correlation of the suicide phenomenon in *Aeromonas* species with virulence and Enteropathogenicity. *J. Clin. Microbiol.* 26, 12, 2615-2619.
- Namdari, H., Cabelli, V.J., 1989. The suicide phenomenon in motile aeromonads. *Appl. Environ. Microbiol.* 55, 543-547.
- Nandi, S., Maurer, J.J., Hofacre, C., Summers, A.O., 2004. Gram-positive bacteria are a major reservoir of Class 1 antibiotic resistance integrons in poultry litter. *PROC. NATL. ACAD. SCI.* , USA. 101, 7118-7122.
- Natrah, F.M., Ruwandeepika, H.A., Pawar, S., Karunasagar, I., Sorgeloos, P., Bossier, P., Defoirdt T., 2011. Regulation of virulence factors by quorum sensing in *Vibrio harveyi*. *Vet Microbiol.* 29: 154 (1-2): 124-129.
- Nealson, K.H., 1978. Isolation, identification and manipulation of luminous bacteria. *Methods Enzymol.* 57, 153-166.
- Nealson, K.H., Hastings, J.W., 1979. Bacterial bioluminescence: its control and ecological significance. *Microbiol. Rev.* 43, 496–518.
- Nei, M., 1973. Analysis of Gene Diversity in Subdivided Populations. *PROC. NATL. ACAD. SCI.* .USA. 70 3, 3321-3323.
- Nemecek-Marshall, M., Wojciechowski, C., Wagner, W.P., Fall, R., 1999. Acetone formation in *Vibrio* family: a new pathway for bacterial leucine catabolism. *J. Bacteriol.* 181, 7493-7499.

References

- Nesper, J., Schild, S., Lauriano, C.M., Kraiss, A., Klose, K. E., Reidl, J., 2002. Role of *Vibrio cholerae* O139 surface polysaccharide in intestinal colonization. *Infect. Immun.* 70, 5590-5996.
- Nhan, D.T., Cam, D.T.V., Wille, M., Defoirdt, T., Bossier, P., Sorgeloos, P., 2010. Quorum quenching bacteria protect *Macrobrachium rosenbergii* larvae from *Vibrio harveyi* infection. *J.Appl. Microbiol.* 109, 1007-1016.
- Nield, B.S., Holmes, A.J., Gillings, M.R., Recchia, G.D., Mabbutt, B.C., Nevalainen, K.M., Stokes, H.W., 2001. Recovery of new integron classes from environmental DNA. *FEMS. Microbiol. Lett.* 195, 59-65.
- Nikaido, H., 1988. Structure and functions of the cell envelope of gram- negative bacteria. *Microbiol. Rev. Infect. Dis.* 10, 2, 279-281.
- Nikaido, H., 2003. Molecular basis of bacterial outer membrane permeability revisited. *Microbiol. Mol. Biol. Rev.* 67, 593-656.
- Ningqiu, L., Junjie, B., Shuqin, W., Xiaozhe, F., Haihua, L., Xing, Y., Cunbin, S., 2008. An outer membrane protein, OmpK, is an effective vaccine candidate for *Vibrio harveyi* in orange-spotted grouper (*Epinephelus coioides*). *Fish and Shellfish Immunology.* 25, 829-833.
- Nishiguchi, M.K., Nair, V.S., 2003. Evolution of symbiosis in the *Vibrionaceae*: a combined approach using molecules and physiology. *Int. J. Syst. Evol. Microbiol.* 53, 2019-2026.
- Nogami, K., Maeda, M., 1992. Bacteria as biocontrol agents for rearing larvae of the crab *Portunus trituberculatus*. *Can. J. Fish. Aquat. Sci.* 49, 2373-2376.
- Noguchi, T., Hwang, D.F., Arakawa, O., Sugita, H., Deguchi, Y., Shida, Y., Hashimoto, K., 1987. *Vibrio alginolyticus*, a tetrodotoxin-producing bacterium, in the intestines of the fish *Fugu vermicularis vermicularis*. *Mar. Biol.* 94, 625-630.
- Nogueroles, I., Basslar, A.R, 2008. Identification of *Vibrio* spp. with a set of dichotomous keys. *J. Appl. Microbiol.* 105 175-185.
- Norqvist, N., Wolf-Watz, H., 1993. Characterization of a novel chromosomal virulence locus involved in expression of a major surface flagella sheath antigen of the fish pathogen *Vibrio anguillarum*. *Infect. Immun.* 61, 2434-2444.

References

- Nottage, A.S., Birkbeck, T.H., 1987. Production of proteinase during experimental infection of *Ostrea edulis* L. larvae with *Vibrio alginolyticus* NCMB 1339 and the antigenic relationship between proteinases produced by marine vibrios pathogenic for fish and shellfish. *J. Fish. Dis.* 10, 265–273.
- Nystrom, T., 2004. Stationary-phase physiology. *Annu. Rev. Microbiol.* 58, 161-181.
- Oakey, H.J., Levy, N., Bourne, D.G., Cullen, B., Thomas, A., 2003. The use of PCR to aid in the rapid identification of *Vibrio harveyi* isolates. *J. Appl. Microbiol.* 95, 1293-1303.
- Oakey, H.J., Owens, L., 2000. A new bacteriophage, VHML, isolated from a toxin-producing strain of *Vibrio harveyi* in tropical Australia. *J. Appl. Microbiol.* 89, 702-709.
- Ochman, H., Lawrence, J.G., Groisman, E., 2000. Lateral gene transfer and the nature of bacterial innovation. *Nature.* 405, 299-304.
- Okada, K., Iida, T., Kita-Tsukamoto, K., Honda, T., 2005. Vibrios commonly possess two chromosomes. *J. Bacteriol.* 187, 752-757.
- Okuda, J., T. Nakai, Chang, P. S., Oh, T., Nishino, T., Koitabashi, T., Nishibuchi, M., 2001. The *toxR* gene of *Vibrio* (*Listonella*) *anguillarum* controls expression of the major outer membrane proteins but not virulence in a natural host model. *Infect. Immun.* 69, 6091-6101.
- Olive, D. M., Bean, P., 1999. Principles and applications of methods for DNA-based typing of microbial organisms. *J. Clin. Microbiol.* 37, 1661–1669.
- Oliver, J. D., 1982. Taxonomic scheme for the identification of marine bacteria. *Deep Sea Research.* 29, 795-798.
- Oliver, J. D., 1993. Formation of viable but nonculturable cells. In S.Kjelleberg (ed), starvation in Bacteria. Plenum Press, New York, N.Y. 239-272.
- Ortigosa, M., Garay, E., Pujalte, M.J., 1994. Numerical taxonomy of Vibrionaceae isolated from oysters and seawater along an annual cycle. *Syst. Appl. Microbiol.* 17, 216-225.

References

- Osawa, R., Okitsu, T., Morozumi, H., Yamai, S., 1996. Occurrence of urease-positive *Vibrio parahaemolyticus* in Kanagawa, Japan, with specific reference to presence of thermostable direct hemolysin (TDH) and the TDH-related-hemolysin gene. *Appl. Environ. Microbiol.* 62, 725-727.
- Osorio, C.R., Klose, K.E., 2000. A region of the transmembrane regulatory protein ToxR that tethers the transcriptional activation domain to the cytoplasmic membrane displays wide divergence among *Vibrio* species. *J. Bacteriol.* 182, 526-528.
- *Ostling, J., Holmquist, L., Flardh, K., Svenblad, B., Jouper-Jann, A., Kjelleberg, S., 1993. Starvation and recovery of *Vibrio*. In S.Kjelleberg (ed), starvation in Bacteria. Plenum Press, New York, N.Y. 103-127.
- O'Toole, G.A., Kolter, R., 1998. Initiation of biofilm formation in *Pseudomonas fluorescens* WCS365 proceeds via multiple, convergent signaling pathways: a genetic analysis. *Mol. Microbiol.* 28, 449-461.
- Ottaviani, D., Bacchiocchi, I., Masini, L., Leoni, F., Carraturo, A., Giammarioli, M., Sbaraglia, G., 2001. Antimicrobial susceptibility of potentially pathogenic halophilic vibrios isolated from seafood. *Int. J. Antimicrobial Agents.* 18, 135-140.
- Ottaviani, D., Masini, L., Bacchiocchi, S., 2003. A biochemical protocol for the isolation and identification of current species of *Vibrio* in seafood. *J. Appl. Microbiol.* 95, 1277-1284.
- Owens, L., Austin, D., Austin, B., 1996. Effect of strain origin on siderophore production in *Vibrio harveyi* isolates. *Dis. Aquat. Org.* 27, 157-160.
- Owens, L., Busico-Salcedo, Nancy., 2006. *Vibrio harveyi*: pretty problems in paradise. In: *The Biology of vibrios*. ASM Press, Washington, DC, pp. 266-280.
- Oxley, A.P., Shipton, W., Owens, L., McKay, D., 2002. Bacterial flora from the gut of the wild and cultured banana prawn, *Penaeus merguensis*. *J. Appl. Microbiol.* 93, 214-223.
- *Pacini, F., 1854. Osservazioni microscopiche e deduzione patologiche sul colera asiatico. *Gaz. Med. Italiana.* 6, 405-412.

References

- Pages, J.M., James, C.E., Winterhalter, M., 2008. The porin and permeating antibiotic: a selective diffusion barrier in Gram negative bacteria. *Nat. Rev. Microbiol.* 6, 893-903.
- Pai, H., Byeon, J.H., Yu, S., KLee, B., Kim, S., 2003. *Salmonella enterica serovar typhi* strains isolated in Korea containing a multidrug resistance class 1 integron. *Antimicrob. Agents. Chemother.* 47, 2006–2008.
- Pakshirajan, P., 2002. Use of antibiotics, drugs and chemicals in shrimp farming and steps for their regulation - a report, Aquaculture Authority News. 14-15.
- Palmer, L.M., Colwell, R.R., 1991. Detection of luciferase gene sequence in nonluminescent *Vibrio cholerae* by colony hybridization and polymerase chain reaction. *Appl. Environ. Microbiol.* 57, 1286–1293.
- Pang, L., Zhang, X.H., Zhong, Y., Chen, J., Li, Y., Austin, B., 2006. Identification of *Vibrio harveyi* using PCR amplification of the *toxR* gene. *Lett. Appl. Microbiol.* 43, 249-255.
- Panicker, G., Call, D.R., Krug, M.J., Bej, A.K., 2004. Detection of pathogenic *Vibrio* spp. in shellfish by using multiplex PCR and DNA microarrays. *Appl. Environ. Microbiol.* 70, 7436-7444.
- Park, T. S., Oh, S. H., Lee, E. Y., Lee, T. K., Park, K. H., Figueras, M. J., Chang, C. L., 2003. Misidentification of *Aeromonas veronii* biovar *sorbria* as *Vibrio alginolyticus* by the Vitek system. *Lett. Appl. Microbiol.* 37, 349-353.
- Park, K., Ono, S, T., Rokuda, M., Jang, H., Okada, K., Iida, T., Honda, T., 2004. Functional characterization of two type III secretion systems of *Vibrio parahaemolyticus*. *Infect. Immun.* 72, 6659-6665.
- Parveen, S., Lukasik, J., Scott, T.M., Tamplin, M.L., Portier, K.M., Sheperd, S., Braun, K., Farrah, S.R., 2006. Geographical variation in antibiotic resistance profiles of *Escherichia coli* isolated from swine, poultry, beef and dairy cattle farm water retention ponds in Florida. *J. Appl. Microbiol.* 100, 50-57.
- Payne, M., Oakey, J., Owens, L., 2004. The ability of two different *Vibrio* spp. bacteriophages to infect *Vibrio harveyi*, *Vibrio cholerae* and *Vibrio mimicus*. *J. Appl. Microbiol.* 97, 663-672.

References

- Pazos, F., Y.Santos, B.Magariños, Bandín, I., Núñez, S., Toranzo, A.E., 1993. Phenotypic characteristics and virulence of *Vibrio anguillarum*-related organisms. *Appl. Environ. Microbiol.* 59, 2969-2976.
- Pedersen, K., Verdonck, L., Austin, B., Austin, D.A., Blanch, A.R., Grimont, P.A.D., Jofre, J., Koblavi, S., Larsen, J.L., Tiainen, T., Vigneulle, M., Swings, J., 1998. Taxonomic evidence that *Vibrio carchariae* Grimes et al. 1985 is a junior synonym of *Vibrio harveyi* (Johnson and Shunk 1936) Bauman et al. 1981. *Int. J. Syst. Bacteriol.* 48, 749-758.
- Pernthaler, A., Pernthaler, J., Eilers, H., Amann, R. (2001) Growth patterns of two marine isolates: Adaptations to substrate patchiness? *Appl. Environ. Microbiol.* 67, 4077-4083.
- Pernthaler, A., Pernthaler, J., Amann, R., 2002a. Fluorescence in situ hybridization and catalyzed reporter deposition for the identification of marine bacteria. *Appl. Environ. Microbiol.* 68, 3094-3101.
- Pernthaler, A., Amann, R., 2004. Simultaneous fluorescence in situ hybridization of mRNA and rRNA in environmental bacteria. *Appl. Environ. Microbiol.* 70, 5526-5533.
- Pereira, J., Shanmugam, S.A., Sulthana, M., Sundaraj, V., 2009. Effect of vaccination on vibriosis resistance of *Fenneropenaeus Indicus*. Tamilnadu. *J. Veterinary and Animal Sciences.* 5, 246-250.
- Phianphak, W., Rengpipat, S., Rukpratanporn, S., Longyant, S., Chaivisuthangkura, P., Sithigorngul, W., Sithigorngul, P., 2005. Production of monoclonal antibodies for detection of *Vibrio harveyi*. *Dis. Aquat. Org.* 63, 161-168.
- Pizzutto, M., Hirst, R.G., 1995. Classification of isolates of *Vibrio harveyi* virulent to *Penaeus monodon* larvae by protein profile analysis and M13 DNA fingerprinting. *Dis. Aquat. Org.* 21, 61-68.
- Pontes, D.S., Pinheiro, F.A., Lima-Bittencourt, C.I., Guedes, R.L.M., Cursino, L., Barbosa, F., Santos, F.R., Chartone-Souza, E., Nascimento, A.M.A., 2009. Multiple Antimicrobial Resistance of Gram-Negative Bacteria from Natural Oligotrophic Lakes under Distinct Anthropogenic Influence in a Tropical Region. *Microbial Ecology.* 58, 762-772.

References

- Prasad, S., Morris, P.C., Hansen, R., Meaden, P.G., Austin, B., 2005. A novel bacteriocin-like substance (BLIS) from a pathogenic strain of *Vibrio harveyi*. *Microbiology*. 151, 3051-3058.
- Priest, F.G., 1977. Extracellular enzyme synthesis in the genus *Bacillus*. *Bacteriological Reviews*. 41, 711-753.
- Proctor, L.M., Gunsalus, R.P. 2000. Anaerobic respiratory growth of *Vibrio harveyi*, *Vibrio fischeri* and *Photobacterium leiognathi* with trimethylamine N-oxide, nitrate and fumarate: ecological implications. *Environ. Microbiol.* 2, 399-406.
- Provenzano, D., Schuhmacher, D.A., Barker, J.L., Klose, K.E., 2000. The virulence regulatory protein ToxR mediates enhanced bile resistance in *Vibrio cholerae* and other pathogenic *Vibrio* species. *Infect. Immun.* 68, 1491-1497.
- Pujalte, M.J., Ortiz-Conde, B.A., Steven, S.E., Esteve, C., Garay, E., Colwell, R.R., 1992. Numerical taxonomy and nucleic acid studies of *Vibrio mediterranei*. *Syst. Appl. Microbiol.* 15, 82-91.
- Pujalte, M. J., Sitji-Bobadilla, A., Maci'n, M. C., Belloch, C., Alvarez-Pellitero, P., P'rez-Sanchez, J., Uruburu, F., Garay, E., 2003. Virulence and molecular typing of *Vibrio harveyi* strains isolated from cultured dentex, gilthead, sea bream and European sea bass. *Syst. Appl. Microbiol.* 26, 284-292.
- Raaijmakers, J.M., David, M.W., Thomasshow, L.S., 1997. Frequency of antibiotic-producing *Pseudomonas* spp. in natural environment. *Appl. Environ. Microbiol.* 63, 881-887.
- Rademaker, J.L.W., Hoste, B., Louws, F.J., Kersters, K., Swings, J., Vauterin, L., Vauterin, P., Bruijn, F.J.D., 2000. Comparison of AFLP and rep-PCR genomic fingerprinting with DNA-DNA homology studies: *Xanthomonas* as a model system. *Int. J. Syst. Evol. Microbiol.* 50, 665-677.
- Radstrom, P., Skold, O., Swedberg, G., Flensburg, J., Roy, P.H., Sundstrom, L., 1994. Transposon Tn5090 of plasmid R751, which carries an integron, is related to Tn7, Mu, and the retroelements. *J. Bacteriol.* 176, 3257-3268.
- Raguenes, G., Christen, R., Guezenec, J., Pignet, P., Barbier, G., 1997. *Vibrio diabolicus* sp. nov., a new polysaccharide-secreting organism isolated from a deep-sea hydrothermal vent polychaete annelid, *Alvinella pompejana*. *Int. J. Syst. Bacteriol.* 47, 989-995.

References

- Ramaiah, N., Hill, R.T., Chun, J., Ravel, J., Matte, M.H., Straube, W.L., Colwell, R.R., 2000. Use of a *chiA* probe for detection of chitinase genes in bacteria from the Chesapeake Bay. *FEMS Microb. Ecol.* 34, 63-71.
- Ramaiah, N., Ravel, J., Straube, W.L., Hill, R.T., Colwell, R.R., 2002. Entry of *Vibrio harveyi* and *Vibrio fischeri* into the viable but nonculturable state. *J. Appl. Microbiol.* 93, 108-116.
- Ramesh, A., Venugopalan, V.K., 1989. Role of luminous bacteria in chitin degradation in the intestine of fish. *World. J. Microbiol. Biotech.* 5, 55-59.
- Rao, N.N., Kornberg, A., 1996. Inorganic phosphate supports resistance and survival of stationary-phase *Escherichia coli*. *J. Bacteriol.* 178, 1394-1400.
- Ravelo, C., Magariños, B., López-Romalde, S., Toranzo, A. E., Romalde, J. L., 2003. Molecular fingerprinting of fish-pathogenic *Lactococcus garvieae* strains by random amplified polymorphic DNA analysis. *J. Clin. Microbiol.* 41:751–756.
- Reddy, G.S.M., Aggarwal, R.K., Matsumotto, G.I., Sivaji, S.I. 2000. *Arthrobacter flavus* sp. nov., a psychrophilic bacterium isolated from a pond in McMurdo Dry Vally, Antarctica. *Int. J. Syst. Evol. Microbiol.* 50, 1553-1561.
- Reed, L.J., Muench, H., 1938. A simple method of estimating fifty percent end points. *Am. J. Hyg.* 27, 493-497.
- Reich, K.A., Schoolnik, G.K., 1994. The light organ symbiont *Vibrio fischeri* possesses a homolog of the *Vibrio cholerae* transmembrane transcriptional activator ToxR. *J. Bacteriol.* 176, 3085-3088.
- Reid, G.C., Woods, D.R., Robb, F.T., 1980. Peptone induction and rifampin-insensitive collagenase production by *Vibrio alginolyticus*. *J. Bacteriol.* 142, 447-454.
- Ren, D., Sims, J.J., Wood, T.K., 2002. Inhibition of biofilm formation and swarming of *Bacillus subtilis* by (5Z)-4-bromo-5-(bromomethylene) 3-butyl-2(5H)-furanone. *Lett. Appl. Microbiol.* 34, 293–299.

References

- Rengpipat, S., Phianphak, W., Piyatiratitivorakul, S., Menasveta, P., 1998. Effects of a probiotic bacterium on black tiger shrimp *Penaeus monodon* survival and growth. *Aquaculture*. 167, 301-313.
- Rengpipat, S., Rukpratanporn, S., Piyatiratitivorakul, S., Menasveta, P., 2000. Immunity enhancement in black tiger shrimp (*Penaeus monodon*) by a probiot bacterium (*Bacillus* S11). *Aquaculture*. 191, 271-288.
- Rengpipat, S., Tunyanun, A., Fast, A., Piyatiratitivorakul, S., Menasveta, P., 2003. Enhanced growth and resistance to *Vibrio* challenge in pond reared black tiger shrimp *Penaeus monodon* fed a *Bacillus* probiotic. *Dis. Aqu. Org.* 55, 169-173.
- Rhoden, D.L., Hancock, G.A., Miller, J.M., 1993. Numerical approach to reference identification of *Staphylococcus*, *Stomatococcus*, and *Micrococcus* spp. *J. Clin. Microbiol.* 31, 490-493.
- Rhodes, G., Huys, G., Swings, J., McGann, P., Hiney, M., Smith, P., Pickup, R.W., 2000. Distribution of oxytetracycline resistance plasmids between Aeromonads in hospital and aquaculture environments: implication of Tn1721 in dissemination of the tetracycline resistance determinant *tetA*. *Appl. Environ. Microbiol.* 66, 3883-3890.
- *Rhodes, M., 1959. *J.Gen. Microbiol.* 21, 221-263.
- Rice, S.A., Mcdougald, D., Kjelleberg, S., 2000. *Vibrio vulnificus*: a physiological and genetic approach to the viable but nonculturable response. *J. Infect. Chemother.* 6, 115-120.
- Riemann, L., Azam, F., 2002. Widespread N-acetyl-D-glucosamine uptake among pelagic marine bacteria and its ecological implications. *Appl. Environ. Microbiol.* 68, 5554-5562.
- Ripp, S., Jegier, P., Birmele, M., Johnson, C.M., Daumer, K.A., Garland, J.L., Saylor, G.S., 2006. Linking bacteriophage infection to quorum sensing signalling and bioluminescent bioreporter monitoring for direct detection of bacterial agents. *J. Appl. Microbiol.* 100, 488-499.
- Rivera, I.N.G., Chowdhury, M.A.R., Huq, A., Jacobs, D., Martins, M.T., Colwell, R.R., 1995. Enterobacterial repetitive intergenic consensus sequences and the PCR generate fingerprints of genomic

References

- DNAs from *Vibrio cholerae* O1, O139 and non-O1 strains. *Appl. Environ. Microbiol.* 61, 2898-2904.
- Robertson, P.A.W., Calderon, J., Carrera, L., Stark, J.R., Zherdmant, M., Austin, B., 1998. Experimental *Vibrio harveyi* infections in *Penaeus vannamei* larvae. *Dis. Aqu. Org.* 32, 151-155.
 - Rodney M. Donlan., 2002. Biofilms: Microbial Life on Surfaces. *Emerg. Infect. Dis.* 8, (9): 881–890.
 - Rohlf, F.J., 1998. Numerical Taxonomy and Multivariate Analysis System. 2.0 Version. New York: Exeter Publishing. Exeter. Software
 - Romalde, J. L., Magariños, B., Villar, C., Barja, J. L., Toranzo, A. E., 1999. Genetic analysis of turbot pathogenic *Streptococcus parauberis* strains by ribotyping and random amplified polymorphic DNA. *FEMS Microbiol. Lett.* 459, 297–304.
 - Romalde, J. L., Castro, D., Magariños, B., López-Cortes, L., Borrego, J. J., 2002. Comparison of ribotyping, randomly amplified polymorphic DNA, and pulsed-field gel electrophoresis for molecular typing of *Vibrio tapetis*. *Syst. Appl. Microbiol.* 25, 544–550.
 - Roque, A., Gomez-Gil, B., 2003. Therapeutic effects of enrofloxacin in an experimental infection with a luminescent *Vibrio harveyi* in *Artemia franciscana* Kellog 1906. *Aquaculture.* 220, 37-42.
 - Rosenberg, E., A.Zuckerberg, C.Rubinovitz, D.L.Gutnick, 1979. Emulsifier of *Arthrobacter* RAG-1: isolation and emulsifying properties. *Appl. Environ. Microbiol.* 37, 402–408.
 - Rosenberg, M., 1984. Bacterial adherence to hydrocarbons: a useful technique for studying cell surface hydrophobicity. *FEMS. Microb. Lett.* 22, 289-295.
 - Rosenberg, M., Gutnick, D., Rossenberg, E., 1980. Adherence of bacteria to hydrocarbons: a simple method for measuring cell surface hydrophobicity. *FEMS Microb. Lett.* 9, 29-33.
 - Rosenberg, M., Kjelleberg, S., 1986. Hydrophobic interactions in bacterial adhesion. *Adv. Microb. Ecol.* 9, 353-393.

References

- Rosenberg, E., Ben-Haim, Y., 2002. Microbial diseases of corals and global warming. *Environ. Microbiol.* 4, 318-326.
- Rowe-Magnus, D.A., Davies, J., Mazel, D., 2000. Impact of integrons and transposons on the evolution of resistance and virulence. *Curr. Top. Microbiol. Immunol.* 264, 167-188.
- Rowe-Magnus, D.A., Mazel, D., 2001. Integrons: natural tools for bacterial genome evolution. *Curr. Opin. Microbiol.* 4, 565-569.
- Rowe-Magnus, D.A., Guetout, A.M., Mazel, D., 2002. Bacterial resistance evolution by recruitment of super-integron gene cassettes. *Mol. Microbiol.* 43, 1657-1669.
- Rowe-Magnus, D.A., Guetout, A.M., Biskri, L., Bougie, P., Mazel, D., 2003. Comparative analysis of superintegrons: engineering extensive genetic diversity in Vibrionaceae. *Genome Res.* 13, 428-442.
- Roy, N.K., Ghosh, R.K., Das, J., 1982. Repression of the alkaline phosphatase of *Vibrio cholerae*. *J. Gen. Microbiol.* 128, 348-353.
- Ruangpan, L., Danayadol, Y., Direkbusarakom, S., Siurairatana, S., Flegel, T.W., 1999. Lethal toxicity of *Vibrio harveyi* to cultivated *Penaeus monodon* induced by a bacteriophage. *Dis. Aqua. Org.* 35, 195-201.
- Rui, H., Liu, Q., Ma, Y., Wang, Q., Zhang, Y., 2008. Roles of LuxR in regulating extracellular alkaline serine proteaseA, extracellular polysaccharide and mobility of *Vibrio alginolyticus*. *FEMS. Microbiol. Lett.* 285, 155-162.
- Ruimy, R., Breittmayer, V., Elbaze, P., Lafay, B., Boussemart, O., Gauthier, M., Christen, R., 1994. Phylogenetic analysis and assessment of the genera *Vibrio*, *Photobacterium*, *Aeromonas*, and *Plesiomonas* deduced from small-subunit rRNA sequences. *Int. J. Syst. Bacteriol.* 44, 416-426.
- Ruwandeepika, H.A., Defoirdt, T., Bhowmick, P.P., Shekar, M., Bossier, P., Karunasagar, I., 2010. Presence of typical and atypical virulence genes in vibrio isolates belonging to the Harveyi clade. *J. Appl. Microbiol.* 109, 888-899.
- Ruwandeepika, H.A., Defoirdt, T., Bhowmick, P.P., Karunasagar, I., Karunasagar I, Bossier P., 2011. In vitro and in vivo expression of

References

- virulence genes in *Vibrio* isolates belonging to the Harveyi clade in relation to their virulence towards gnotobiotic brine shrimp (*Artemia franciscana*). *Environ Microbiol.* 13 (2), 506-517.
- Saeed, M.O., 1995. Association of *Vibrio harveyi* with mortalities in cultured marine fish in Kuwait. *Aquaculture.* 136, 21-29.
 - Sahul Hameed, A.S., Rahaman, K.H., Alagan, A., Yoganandhan, K., 2003. Antibiotic resistance in bacteria isolated from hatchery-reared larvae and post-larvae of *Macrobrachium rosenbergii*. *Aquaculture.* 217, 39-48.
 - Saitou, N., Nei, M., 1987. The Neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 4, 406-425.
 - Sambasivam, S., Chandran, R., Khan, S.A., 2003. Role of probiotics on the environment of shrimp pond. *J. Environ. Biol.* 24, 103-106.
 - Sambrook, J., Russell, D.W., 2001. *Molecular Cloning - A Laboratory Manual.* 3rd Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
 - Sangali, S., Brandelli, A., 2000. Feather keratin hydrolysis by a *Vibrio* sp. Strain kr2. *J. Appl. Microbiol.* 89, 735-743.
 - Saulnier, D., Haffner, P., Goarant, C., Levy, P., Ansquer, D., 2000. Experimental infection models for shrimp vibriosis studies: a review. *Aquaculture.* 191, 133-144.
 - Sawabe, T., Sugimura, I., Ohtsuka, M., Nakano, K., Tajima, K., Ezura, Y., Christen, R., 1998. *Vibrio halioticoli* sp. nov., a nonmotile alginolitic marine bacterium isolated from the gut of the abalone *Haliotis discus hannai*. *Int. J. Syst. Bacteriol.* 48, 573-580.
 - Sawabe, T., Setoguchi, N., Inoue, S., Tanaka, R., Ootsubo, M., Yoshimizu, M., Ezura, Y., 2003. Acetic acid production of *Vibrio halioticoli* from alginate, a possible role for establishment of abalone-*V. halioticoli* association. *Aquaculture.* 219, 671-679.
 - Sawabe, T., Hayashi, K., Moriwaki, J., Thompson, F.L., Swings, J., Potin, P., Christen, R., Ezura, Y., 2004a. *Vibrio gallicus* sp. nov., isolated from the gut of the French abalone *Haliotis tuberculata*. *Int. J. Syst. Evol. Microbiol.* 54, 843-846.

References

- Sawabe, T., Hayashi, K., Moriwaki, J., Fukui, Y., Thompson, F.L., Swings, J., Christen, R. 2004b. *Vibrio neonatus* sp.nov. and *Vibrio ezurae* sp.nov. isolated from hr gut of Japaanese abalones. *Syst. Appl. Microbiol.* 27, 527-534.
- Sawabe, T., Kita-Tsukamoto, K., Fabiano L. Thompson, 2007. Inferring the Evolutionary History of Vibrios by Means of Multilocus Sequence Analysis. *J. Bacteriol.* 189, 7932-7936.
- Salyers, A. A., White, D.D., 1994. Bacterial Pathogenesis, a Molecular Approach. 2nd Ed ASM Press, Washington, DC, USA. 47-62.
- Schrank, G. D., Verwey, W. F., 1976. Distribution of cholera organisms in experimental *Vibrio cholerae* infections: proposed mechanism of pathogenesis and antibacterial immunity. *Infect. Immun.* 13, 195-203.
- Schiewe, M.H., Trust, T.J., Crosa, J.H., 1981. *Vibrio ordalii* sp. nov.: a causative agent of vibriosis in fish. *Curr. Microbiol.* 6, 343-348.
- Schmidt, A.S., Bruun, M.S., Dalsgaard, I., Pedersen, K., Larsen, J.L., 2000. Occurrence of antimicrobial resistance in fish-pathogenic and environmental bacteria associated with four Danish rainbow trout farms. *Appl. Environ. Microbiol.* 66, 4908-4915.
- Schmidt, A.S., Bruun, M.S., Dalsgaard, I., Larsen, J.L., 2001. Incidence, distribution, and spread of tetracycline resistance determinants and integron-associated antibiotic resistance genes among motile Aeromonads from a fish farming environment. *Appl. Environ. Microbiol.* 67, 5675-5682.
- Scholz, U, Diaz, G.G., Ricque, D., Suarez, C., Albores, V.F., Latchford, J., 1999. Enhancement of Vibriosis resistance in juvenile *Penaeus vannamei* supplementation of diets with different yeast products. *Aquaculture.* 176, 271-283.
- Schulte, A., Ruamchan, S., Khunkaewla, P., Suginta, W., 2009. The outer membrane protein VhOmp of *Vibrio harveyi*: pore-forming properties in black lipid membranes. *J. Memb. Biol.* 230, 101-111.
- Schulze, A.D., Alabi, A.O., Tattersall-Sheldrake, A.R., Miller, K.M., 2006. Bacterial diversity in a marine hatchery: balance between

References

- pathogenic and potentially probiotic bacterial strains. *Aquaculture*. 256, 50-73.
- Scoaris, D.D.O., Colacite, J.V., Nakamura, C., Ueda-Nakamura, T., Filho, B.A.D.A., Filho, B.P.D., 2008. Virulence and antibiotic susceptibility of *Aeromonas* spp. isolated from drinking water. *Antonie van Leeuwenhoek*. 93, 111-122.
 - Shao, C.P., Hor, L.I., 2001. Regulation of metalloprotease gene expression in *Vibrio vulnificus* by a *Vibrio harveyi* LuxR homologue. *J. Bacteriol*. 183, 1369-1375.
 - *Shewan, J. M., Hodgkiss, W., 1954. *Nature*. 63, 208-209.
 - Shieh, W.Y., Chen, A.L., Chiu, H.H., 2000. *Vibrio aerogenes* sp. nov., a facultatively anaerobic marine bacterium that ferments glucose with gas production. *Int. J. Syst. Evol. Microbiol*. 50, 321-329.
 - Shinoda, S., Matsuoka, H., Tsuchie, T., Miyoshi, S., Yamamoto, S., Taniguchi, H., Mizuguchi, Y., 1991. Purification and characterization of a lecithin-dependent haemolysin from *Escherichia coli* transformed by a *Vibrio parahaemolyticus* gene. *J. Gen. Microbiol*. 137, 2705-2711.
 - Shinoda, S., Kariyama, R., Ogawa, M., Takeda, Y., T.Miwatani, 1976. Flagellar antigens of various species of the genus *Vibrio* and related genera. *Int. J. Syst. Bacteriol*. 26, 97-101.
 - Shyu, Y.C., Chiu, C.C., Lin, F.P., 1999. Cloning, sequencing, expression and characterization of the Manganese superoxide dismutase gene from *Vibrio alginolyticus*. *Biochem. Mol Bio. Int*. 47, (5): 803- 814.
 - Simidu, U., Noguchi, T., Hwang, D., Shida, Y., Hashimoto, K., 1987. Marine bacteria which produce tetrodotoxin. *Appl. Environ. Microbiol*. 53, 1714-1715.
 - Simonet, V. C., Balse, A., Klose, K.E., Delcour, A.H., 2003. The *Vibrio cholerae* porins OmpU and OmpT have distinct channel properties. *J. Biol. Chem*. 278, 17539-17545.
 - Singh, I.S.B., Lakshmanaperumalaswamy, P., Chandramohan, D., 1989. Bacteriology of eggs and larvae of *Penaeus indicus* in hatchery. In: Nair, N.B. (Ed.), Proceedings of Kerala Science Congress, Thiruvananthapuram, Kerala, India, pp. 95 -107.

References

- Skjermo, J., Salvesen, I., Oie, G., Olsen, Y., Vadstein, O., 1997. Microbially matured water: a technique for selection of non-opportunistic bacterial flora in water that may improve performance of marine larvae. *Aquac. Int.* 5, 13-28.
- Skjermo, J., Vadstein, O., 1999. Techniques for microbial control in the intensive rearing of marine larvae. *Aquaculture*. 177, 333-343.
- Sneath, P.H.A., Johnson, R., 1972. The influence of numerical taxonomy similarities of errors in microbiological tests. *J. Gen. Microbiol.* 72, 377-392.
- *Sneath, P.H., Sokal, R., 1973. Numerical Taxonomy. San Francisco: W. H. Freeman.
- Sneath, P.H.A., 1974. Test reproducibility in relation to identification. *Int. J. Syst. Bacteriol.* 24, 508-523.
- Snoussi, M., Noumi, E., Cheriaa, J., Sechi, L.A., Zanetti, S., Bakhrouf, A., 2008a. Adhesive properties of environmental *V. alginolyticus* strains to biotic and abiotic surfaces. *New. Microbiol.* 31, 489-500.
- Snoussi, M., Noumi, E., Usai, D., Sechi, L.A., Zanetti, S., Bakhrouf, A., 2008b. Distribution of some virulence related-properties of *Vibrio alginolyticus* from Mediterranean seawater (Bay of Khenis, Tunisia): investigation of eight *Vibrio cholerae* virulence genes. *World. J. Microbiol. Biotechnol.* 24, 2133-2141.
- Soll, D. R., 2000. The ins and outs of DNA fingerprinting the infectious fungi. *Clin. Microbiol. Rev.* 13, 332-370.
- Somarny, W.M.Z., Mariana, N.S., Neela, V., Rozita, R., Raha, A.R. 2002. Differentiation of pathogenic *Vibrio* species by RAPD. *J. Med. Sci.* 2, 165-169.
- Sommerse, I., Krossoy, B., Biering, E., Frost, P., 2005. Vaccines for fish in aquaculture. *Expert. Rev. Vaccines.* 4, 89-101.
- Son, R.G.R., Sahilah, A.M., Zainuri, A., Raha, A.R., Salmah, I., 1997. Antibiotic resistance and plasmid profile of *A. hydrophila* isolates from cultured fish, Tilapia (*Tilapia mossambica*). *Lett. Appl. Microbiol.* 24, 479-482.

References

- Song, Y.L., Cheng, W., Wang, C.H., 1993. Isolation and characterization of *Vibrio damsela* infectious for cultured shrimp in Taiwan. *J. Invert. Pathol.* 61, 24-31.
- Song, Y.L., W.Cheng, C.H.Shen, Y.C.Ou, H.B.Song, 1990. Occurrence of *Vibrio vulnificus* in cultured shrimp and eel in Taiwan. *NSC Symp. Ser.* 16, 172-179.
- Song, H.H., Kou, G.H. Sung, Y.L., 1994. Vibrosis resistance induced by glucan treatment in tiger prawn (*Penaeus monodon*). *Fish Pathol.* 29, 11-17.
- Sorgeloos, P., Bosswyl, E., Lavina, E., Baeza-Meza, M., Persoone, G., 1977. Decapsulation of Artemia cysts: a simple technique for the improvement of the use of brine shrimp in aquaculture. *Aquaculture.* 12, 311-315.
- Sorum, H., Hvaal, A.B., Heum, M., Daae, F.L., Wiik, R., 1990. Plasmid profiling of *Vibrio salmonicida* for epidemiological studies of cold-water vibrios in Atlantic salmon (*Salmo salar*) and cod (*Gadus morhua*). *Appl. Environ. Microbiol.* 56, 1033-1037.
- Soto Rodríguez, S.A., Roque, A., Partida, A.L., Guerra, M.L., Flores, A.L., Gil, B.G., 2003a. Virulence of luminous Vibrios to *Artemia franciscana* Kellog, 1906, nauplii. *Dis. Aquat. Org.* 53, 231-240.
- Soto Rodríguez, S.A., Simões, N., Jones, D.A., Roque, A., Gómez-Gil, B., 2003b. Assessment of fluorescent labeled bacteria (FLB) for evaluation of in vivo uptake of bacteria (*Vibrio* spp.) by crustacean larvae. *J. Microbiol. Meth.* 52, 101-114.
- Soto-Rodriguez, S., Armenta, M., Gomez Gil, B., 2006. Effects of enrofloxacin and florfenicol on survival and bacterial population in an experimental infection with luminescent *Vibrio campbelli* in shrimp larvae of *Litopenaeus vannamei*. *Aquaculture.* 255, 48-54.
- *Spearman.C, Karber, G., 1996. Viorology methods manual (Eds) Brain W.J. Mahy and Hillar O.Kangro. Academic Press. Harcourt Brace and Company, Publishers. London, San Diego, New York, Boston, Sydney, Tokyo, Toronto. 33-37.
- Sperandio, V., Giro' n, J.A., Silveria, W.D., Kaper, J.B., 1995. The OmpU Outer Membrane Protein, a Potential Adherence Factor of *Vibrio cholerae*. *Infect. Immun.* 63, 4433-4438.

References

- Sperandio, V. L., Mellies, J., Nguyen, W., Shin, S., Kaper, B.J., 1999. Quorum sensing controls expression of the type III secretion gene transcription and protein secretion in enterohemorrhagic and enteropathogenic *Escherichia coli*. *PROC. NATL. ACAD. SCI. USA*. 96, 15196-15201.
- Stackebrandt, E., Frederiksen, W., Garrity, G. M., Grimont, P. A., Kampfer, P., Maiden, M. C., Nesme, X., Rossello-Mora, R., Swings, J., Truper, H. G., Vauterin, L., Ward, A. C., Whitman, W. B., 2002. Report of the ad hoc committee for the re-evaluation of the species definition in bacteriology. *Int. J.Syst. Evol. Microbiol.* 52, 1043–1047.
- Staley, J.T., Irgens, R.L., Brenner, D.J., 1987. *Enhydrobacter aerosaccus* gen. nov., sp. nov., a gas-vacuolated, facultatively anaerobic, heterotrophic rod. *Int. J. Syst. Bacteriol.* 37, 289-291.
- Staples, D.J., Heales, D.S., 1991. Temperature and salinity optima for growth and survival of juvenile banana prawns *Penaeus merguensis*. *J. Exp. Mar. Biol. Ecol.* 154, 251-274.
- Staroscik, A. M., Steven, M. D., David, R. N., 2005. Regulation of the *Vibrio anguillarum* metalloprotease EmpA by post-translational modification. *J. Bacteriol.* 187, 7, 2257-2260.
- Stevens, A. M., Dolan, K. M., Greenberg, E. P., 1994. Synergistic binding of the *Vibrio fischeri* LuxR transcriptional activator domain and RNA polymerase to the lux promoter region. *PROC. NATL. ACAD. SCI. USA* 91, 12619-12623.
- Sudheesh, P.S., Xu, H. S., 2001. Pathogenicity of *V.parahaemolyticus* in tiger prawn *P. monodon* Fabricus: possible role of extracellular protease. *Aquaculture.* 196, 37-46.
- Sudheesh, P. S., Jie, K., Xu, H. S., 2002. Random amplified polymorphic DNA-PCR typing of *Vibrio parahaemolyticus* and *V. alginolyticus* isolated from cultured shrimps. *Aquaculture.* 207, 11-17.
- Sun, W., Cao, J.G., Teng, K., Meighen, E.A., 1994. Biosynthesis of poly-3-hydroxybutyrate in the luminescent bacterium, *Vibrio harveyi*, and regulation by the lux autoinducer, N-(3-hydroxybutanoyl) homoserine lactone. *J. Biol. Chem.* 269, 20785-20790.

References

- Sun, K., Zhang, W.W., Hou, J.H., Sun, L., 2009a. Immunoprotective analysis of vhhP2, a *Vibrio harveyi* vaccine candidate. *Vaccine*. 27, 2733-2740.
- Sun, K., Hu, Y. H., Zhang, X. H., Bai, F. F., Sun, L., 2009b. Identification of vhhP2, a novel genetic marker of *Vibrio harveyi*, and its application in the quick detection of *V. harveyi* from animal specimens and environmental samples. *J. Appl. Microbiol.* 107, 1251–1257.
- Sung, H.H., Hsu, S.F., Chen, C.K., Ting, Y.Y., Chao, W.L., 1999a. Relationships between disease outbreak in culture tiger shrimp (*Penaeus monodon*) and the composition of vibrio communities in pond water and shrimp hepatopancreas during cultivation. *Aquaculture*. 192, 101-110.
- Sung, H.H., Li, H.C., Tsai, F.M., Ting, Y.Y., Chao, W.L., 1999b. Changes in the composition of *Vibrio* communities in pond water during tiger shrimp (*Penaeus monodon*) cultivation and in the hepatopancreas of healthy and disease shrimp. *J. Expt. Marine Ecol.* 236, 261-271.
- Sung, H.H., Hsu, S.F., Chen, C.K., Ting, Y.Y., Chao, W.L., 2001. Relationship between disease out breaks in cultured tiger shrimp (*Penaeus monodon*) and the composition of *Vibrio* communities in pond water and shrimp hepatopancreas during cultivation. *Aquaculture*. 192, 101-110.
- Surette, M.G., B.LBassler, 1999a. Regulation of autoinducer production in *Salmonella typhimurium*. *Mol Microbiol.* 31 585-595.
- Surette, M.G., Miller, M.B., Bassler, B.L., 1999b. Quorum sensing in *Escherichia coli*, *Salmonella typhimurium*, and *Vibrio harveyi*: A new family of genes responsible for autoinducer production. *PROC. NATL. ACAD. SCI.* . 96, 1639-1644.
- Svitil, A.L., Chadhain, S.M.N., Moore, J.A., Kirchman, D.L., 1997. Chitin degradation proteins produced by the marine bacterium *Vibrio harveyi* growing on different forms of chitin. *Appl. Environ. Microbiol.* 63, 408-413.
- Svitil, A., Kirchman, D., 1998. A chitin-binding domain in a marine bacterial chitinase and other microbial chitinases: implications for the ecology and evolution of 1, 4- beta-glycanases. *Microbiology*. 144, 1299-1308.

References

- Swartzman, E., Kapoor, S., Graham, A. F., Meighen, E.A., 1990. A new *Vibrio fischeri* lux gene precedes a bidirectional termination site for the lux operon. *J. Bacteriol.* 172, (12). 6797-6802.
- Swartzman, E., Silverman, M., Meighen, E.A., 1992. The *luxR* gene product of *Vibrio harveyi* is a transcriptional activator of the lux promoter. *J. Bacteriol.* 174, 7490-7493.
- Swartzman, E., Meighen, E.A., 1993. Purification and characterization of a poly(dA-dT) lux-specific DNA-binding protein from *Vibrio harveyi* and identification as *LuxR*. *J. Biol. Chem.* 268, 16706-16716.
- Swift, S., Lynch, M.J., Fish, L., Kirke, D.F., Tomas, J.M., Gordon, S.A., Stewart, B., Williams, P., 1999. Quorum sensing-dependent regulation and blockade of exoprotease production in *Aeromonas hydrophila*. *Infect. Immun.* 67, 5192-5199.
- Tagomori, K., Iida, T., Honda, T., 2002. Comparison of Genome structures of Vibrios, bacteria possessing two chromosomes. *J. Bacteriol.* 4351-43582.
- Takahashi, Y., Itami, T., Nakagawa, A., Nishimura, H., Abe, T., 1985. Therapeutic effects of oxytetracycline trial tablets against vibriosis in cultured Kuruma prawns *Penaeus japonicus* Bate. *Bull. Jpn. Soc. Sci. Fish.* 51, 1639-1643.
- Takahashi, Y., Itami, T., Maeda, M., Kondo, M., 1998. Bacterial and viral diseases of kuruma shrimp (*Penaeus japonicus*) in Japan. *Fish Pathol.* 33, 357-364.
- Tamura, K., Nei, M., Kumar, S., 2004. Prospects for inferring very large phylogenies by using the neighbor-joining method. *PROC. NATL. ACAD. SCI. (USA)* 101, 11030-11035.
- Tamura, K., Dudley, J., Nei, M., Kumar, S., 2007. MEGA4: Molecular Evolutionary genetics Analysis (MEGA) software version 4.0. *Mol. Biol. Evol.* 24, 1596-1599.
- Tang, Y.W., Ellis, N.M., Hopkins, M.K., Dodge, D.E., Persing, D.H., 1998. Comparison of Phenotypic and Genotypic Techniques of Identification of Unusual Aerobic Pathogenic Gram Negative Bacilli. *J. Clin. Microbiol.* 3674-3679.

References

- Taylor, R.K., Miller, V.L., Furlong, D.B., Mekalanos, J.J., 1987. Use of *phoA* gene fusions to identify a pilus colonization factor coordinately regulated with cholera toxin. *Proc. Natl. Acad.Sci. USA* 84, 2833-2837.
- Tendencia, E.A., 2004. The first report of *Vibrio harveyi* infection in the sea horse *Hippocampus kuda* Bleekers 1852 in the Philippines. *Aqua. Res.* 35, 1292-1294.
- Tendencia, E.A., de la Pena, L.D., 2001. Antibiotic resistance of bacteria from shrimp ponds. *Aquaculture.* 195, 193-204.
- Teo, J.W.P., Suwanto, A., Poh, C.L., 2000. Novel β -lactamase genes from two environmental isolates of *Vibrio harveyi*. *Antimicrob. Ag. Chem.* 44, 1309-1314.
- Teo, J.W.P., Zhang, L.H., Poh, C.I., 2003. Cloning and characterization of a metalloprotease from *Vibrio harveyi* strain AP6. *Gene.* 303, 147-156.
- Teuber, M., 1999. Spread of antibiotic resistance with food-borne pathogens. *Cellular and Molecular Life Sciences.* 56, 755-763.
- Teuber, M., 2001. Veterinary use and antibiotic resistance. *Curr. Opin. Microbiol.* 4, 493-499.
- Thompson, J.D., Higgins, D.G., Gibson, T.J., 1994. CLUSTAL-W: Improving the sensitivity of progressive multiple sequence alignment through sequences weighting, position-specific gap penalties and weight matrix choice. *Nuc. Acids. Res.* 22, 4673-4680.
- Thompson, J.D., Gibson, J., Plewniak, T., Jeanmougin, F., Higgins, D.G., 1997. The CLUSTAL-X Windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nuc. Acids. Res.* 25, 4876-4882.
- Thompson, F.L., Hoste, B., Vandemeulebroecke, K., Swings, J., 2001. Genomic diversity amongst *Vibrio* isolates from different sources determined by fluorescent amplified length polymorphism. *Syst. Appl. Microbiol.* 24, 520-538.
- Thompson, F.L., Thompson, C.C., Li, Y., Gomez-Gil, B., Vandenberghe, J., Hoste, B., Swings, J., 2003a. *Vibrio kanaloae* sp. nov., *Vibrio pomeroyi* sp. nov., and *Vibrio chagasii* sp. nov., from

References

- sea water and marine animals. *Int. J. Syst. Evol. Microbiol.* 53, 753-759.
- Thompson, F.L., Thompson, C.C., Swings, J., 2003b. *Vibrio tasmaniensis* sp. nov., isolated from Atlantic salmon (*Salmo salar* L.). *Syst. Appl. Microbiol.* 26, 65-69.
 - Thompson, F. L., Iida, T., Swings, J., 2004a. Biodiversity of vibrios. *Microbiol. Mol. Biol. Rev.* 68, 403-431.
 - Thompson, C.C., Thompson, F.L., Vandemeulebroecke, K., Hoste, B., Dawyndt, P., Swings, J., 2004b. Use of recA as an alternative phylogenetic marker in the family *Vibrionaceae*. *Int. J. Syst. Evol. Microbiol.* 54, 919-924.
 - Thompson, F. L., Gevers, D., Thompson, C. C., Dawyndt, P., Naser, S., Hoste, B., Munn, C. B., Swings, J., 2005. Phylogeny and molecular identification of vibrios on the basis of multilocus sequence analysis. *Appl. Environ. Microbiol.* 71, 5107-5115.
 - Thompson, F. L., Gomez-Gil, B., Vasconcelos, A. T., Sawbae, T., 2007. Multilocus sequence analysis reveals that *V.harveyi* and *V.campbellii* are distinct species. *Appl. Environ. Microbiol.* 73, 4279-4285.
 - Thompson, C.C., Vicente, A.C.P., Souza, R.C., Vasconcelos, A.T.R., Vesth, T., Alves, N., Ussery, D.W., Iida, T., Thompson, F.L., 2009. Genomic taxonomy of vibrios. *BMC Evolutionary Biology.* 9, 1-16.
 - Thompson, J.R., Randa, M.A., Marcelino, L.A., Tomita-Mitchell, A., Lim, E., Polz, M.F., 2004. Diversity and Dynamics of a North Atlantic Coastal *Vibrio* Community. *Appl. Environ. Microbiol.* 70: 7, 4103–4110.
 - Thornley, M.J., 1960. The differentiation of *Pseudomonas* from other gram-negative bacteria on the basis of organic metabolism. *J. Appl. Bacteriol.* 23, 37-52.
 - Thyssen, A., Eygen, S.V., Hauben, L., Goris, J., Swings, J., Ollevier, F., 2000. Application of AFLP for taxonomic and epidemiological studies of *Photobacterium damsela* subsp. *piscicida*. *Int. J. Syst. Evol. Microbiol.* 50, 1013-1019.
 - Tinh, N.T.N., Linh, N.D., Wood, T.K., Dierckens, K., Sorgeloos, P., Bossier, P., 2007. Interference with the quorum sensing systems in a

References

- Vibrio harveyi* strain alters the growth rate of gnotobiotically cultured rotifer *Brachionus plicatilis*. *J. Appl. Microbiol.* 103, 194-203.
- Tolmasky, M.E., Actis, L.A., Toranzo, A.E., Barja, J.L., Crosa, J.H., 1985. Plasmids mediating iron uptake in *Vibrio anguillarum* strains isolated from turbot in Spain. *J. Gen. Microbiol.* 131, 1989-1997.
 - Toranzo, A.E., Barja, J.L., 1990. A review of the taxonomy and seroepizootiology of *Vibrio anguillarum*, with special reference to aquaculture in the northwest of Spain. *Dis. Aquat. Org.* 9, 73-82.
 - Toranzo, A.E., Novoa, B., Romalde, J.L., Nunez, S., Devesa, S., Marino, E., Silva, R., Martinez, E., Figueras, A., Barja, J.L., 1993. Microflora associated with healthy and diseased turbot (*Scophthalmus maximus*) from three farms in northwest Spain. *Aquaculture.* 114, 189-202.
 - Torres, A.G., Zhou, X., Kape, J.B., 2005. Adherence of Diarrheagenic *Escherichia coli* Strains to Epithelial Cells. *Infect. Immun.* 73, (1): 18-29.
 - Touraki, M., Niopas, I., Kastritsis, C., 1999. Bioaccumulation of trimethoprim, sulfamethoxazole and N-acetyl-sulfamethoxazole in *Artemia nauplii* and residual kinetics in seabass larvae after repeated oral dosing of medicated nauplii. *Aquaculture.* 175, 15-30.
 - Tovar, D., Zambonino, J., Cahu, C., Gatesoupe, F.J., Vazquez, J.R., Lesel, R., 2002. Effect of yeast incorporation in compound diet on digestive enzyme activity in sea bass larvae. *Aquaculture.* 204, 113-123.
 - Towbin, H., Staehelin, T., Gordon, J., 1979. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *PROC. NATL. ACAD. SCI. USA.* 6, 4350-4354.
 - Truu, J., Talpsep, E., Heinaru, E., Stottmeister, U., Wand, H., Heinaru, A., 1999. Comparison of API 20NE and Biolog GN identification systems assessed by techniques of multivariate analyses. *J. Microbiol. Methods.* 36 193-201.
 - Urakawa, H., Kita-Tsukamoto, K., Ohwada, K., 1999a. 16SrDNA restriction fragment length polymorphism analysis of

References

- psychrotrophic vibrios from Japanese coastal water. *Can. J. Microbiol.* 45, 1001-1007.
- Urakawa, H., Kita-Tsukamoto, K., Ohwada, K., 1999b. Reassessment of the taxonomic position of *Vibrio iliopiscarius* and proposal for *Photobacterium iliopiscarium comb. nov.* *Can. J. Microbiol.* 45, 67-76.
 - Urwin, R., Maiden, M.C., 2003. Multi-locus sequence typing: a tool for global epidemiology. *Trends Microbiol.* 10, 479-487.
 - van der Meer, J.R., Sentchilo, V., 2003. Genomic islands and the evolution of catabolic pathways in bacteria. *Curr. Opin. Biotechnol.* 14, 248-254.
 - Vandamme, P., Pot, B., Gillis, M., de Vos, P., Kersters, K., Swings, J., 1996. Polyphasic taxonomy, a consensus approach to bacterial systematics. *Microbiol. Rev.* 60, 407-438.
 - Vandenberghe, J., Li, Y., Verdonck, L., Li, J., Sorgeloos, P., Xu, H.S., Swings, J., 1998. Vibrios associated with *Penaeus chinensis* (Crustacea:Decapoda) larvae in Chinese shrimp hatcheries. *Aquaculture.* 169, 121-132.
 - Vandenberghe, J., Verdonck, L., Robles-Arozarena, R., Rivera, G., Bolland, A., Balladares, M., Gomez-Gil, B., Calderon, J., Sorgeloos, P., Swings, J., 1999. Vibrios associated with *Litopenaeus vannamei* larvae, postlarvae, broodstock, and hatchery probionts. *Appl. Environ. Microbiol.* 65, 2592-2597.
 - Vandenberghe, J., Thompson, F.L., Gomez Gil, B., Swings, J., 2003. Phenotypic diversity amongst *Vibrio* isolates from marine aquaculture systems. *Aquaculture.* 219, 9-20.
 - Venkateswaran, K., Dohmoto, N., Harayama, S., 1998. Cloning and nucleotide sequence of the *gyrB* gene of *Vibrio parahaemolyticus* and its application in detection of this pathogen in shrimp. *Appl. Environ. Microbiol.* 64, (2): 681-687.
 - Verschuere, L., Rombaut, G., Sorgeloos, P., Verstraete, W., 2000a. Probiotic bacteria as biological control agents in aquaculture. *Microbiol. Mol Bio. Revs.* 64, 655-671.
 - Verschuere, L., Heang, H., Criel, G., Sorgeloos, P., Verstraete, W., 2000b. Selected bacterial strains protect *Artemia* spp. from the pathogenic effects of *Vibrio proteolyticus* CW8T2. *Appl. Environ. Microbiol.* 66, 1139-1146.

References

- Vici, V., Singh, I.S.B., Bhat, S.G., 2000. Application of bacterins and yeast *Acremonium dyosporii* to protect the larvae of *Macrobrachium rosenbergii* from vibriosis. *Fish and Shellfish Immunology*. 10, 559-563.
- Vidgen, M., Carson, J., Higgins, M., Owens, L., 2006. Changes to the phenotypic profile of *Vibrio harveyi* when infected with the *Vibrio harveyi* myovirus-like (VHML) bacteriophage. *J. Appl. Microbiol.* 100, 481-487.
- Vijayan, K.K., Singh, I.S.B., Jayaprakash, N.S., Alavandi, S.V., Somnath Pai, S., Preetha, R., Rajan, J.J.S., Santiago, T.C., 2006. A brackishwater isolate of *Pseudomonas* PS-102, a potential antagonistic bacterium against pathogenic vibrios in penaeid and non-penaeid rearing systems. *Aquaculture*. 251, 192-200.
- Vine, N.G., Leukes, W.D., Kaiser, H., 2004a. In vitro growth characteristics of five candidate aquaculture probiotics and two fish pathogens grown in fish intestinal mucus. *FEMS. Microbiolo. Lett.* 231, 145-152.
- Vine, N., Leukes, W., Kaiser, H., Daya, S., Baxter, J., Hecht, T., 2004b. Competition for attachment of aquaculture candidate probiotic and pathogenic bacteria on fish intestinal mucus. *J. Fish. Dis.* 27, 319-326.
- Vine, N.G., Leukes, W.D., Kaiser, H., 2006. Probiotics in marine larviculture. *FEMS. Microbiol. Rev.* 30, 404-427.
- Vinod, M.G., Shivu, M.M., Umesh, K.R., Rajeeva, B.C., Krohne, G., Karunasagar, I., Karunasagar, I., 2006. Isolation of *Vibrio harveyi* bacteriophage with a potential for biocontrol of luminous vibriosis in hatchery environments. *Aquaculture*. 255, 117-124.
- Vuddhakul, V., Nakai, T., Matsumoto, C., Oh, T., Nishino, T., Chen, C.H., Nishibuchi, M., Okuda, J., 2000a. Analysis of *gyrB* and *toxR* gene sequences of *Vibrio hollisae* and development of *gyrB*- and *toxR*-targeted PCR methods for isolation of *V.hollisae* from the environment and its identification. *Appl. Environ. Microbiol.* 66, 3506-3514.
- Wagner, P.L., Waldor, M.K., 2002. Bacteriophage Control of Bacterial Virulence. *Infect. Immun.* 70, 3985-3993.

References

- Waldor, M.K., 1998. Bacteriophage biology and bacterial virulence. *Trends in Microbiology*. 6, 295-297.
- Waldor, M.K., D.Raychaudhuri., 2000. Treasure trove for cholerae research. *Nature*. 406, 469-470.
- Waldor, M.K., Friedman, D.I., 2005. Phage regulatory circuits and virulence gene expression. *Curr. Opin. Microbiol.* 8, 459-465.
- *Wallwork, J.A, 1976. The Distribution and Diversity of Soil Fauna Chapter-2. 1st Ed Academic Press, London, 355. S
- Walker, P.J., Winton, J.R., 2010. Emerging viral diseases of fish and shrimp. *Vet.Res.*41: 51, 1-24.
- Wang, J., Jiang, Y., Vincent, M., Sun, Y., Yu, H., Wang, J., Bao, Q., Kong, H., Hu, S., 2005. Complete genome sequence of bacteriophage T5. *Virology*. 332, 45-65.
- Wang, S.Y., Lauritz, J., Jass, J., Milton, D.L., 2003. Role for the major outer-membrane protein from *Vibrio anguillarum* in bile resistance and biofilm formation. *Microbiology*. 149, 1061-1071.
- Wang, Y.B., Xu, Z.R., 2006. Effect of probiotics for common carp (*Cyprinus carpio*) based on growth performance and digestive enzyme activities. *Anim. Feed Sci. Technol.* 127, 283-292.
- Warner, J.M., Oliver, J.D., 1999. Randomly Amplified Polymorphic DNA analysis of clinical and environmental isolates of *Vibrio vulnificus* and other *Vibrio* species. *Appl. Environ. Microbiol.* 65, 1141-1144.
- Watnick, P.I., Lauriano, C.M., Klose, K.E., Croal, L., Kolter, R., 2001. The absence of a flagellum leads to altered colony morphology, biofilm development and virulence in *Vibrio cholerae* O139. *Mol. Microbiol* 39, 223-235.
- Weichart, D., Kjelleberg, S., 1996. Stress resistance and recovery potential of culturable and viable but nonculturable cells of *Vibrio vulnificus*. *Microbiology*. 142, 845-853.
- Welch, T.J., Bartlett, D.H., 1998. Identification of a regulatory protein required for pressure-responsive gene expression in the deep-sea bacterium *Photobacterium* species strain SS9. *Mol. Microbiol.* 27, 977-985.

References

- Welsh, J., McClelland, N.M., 1990. Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acids Res.* 18, 7213-7218.
- Westerdahl, A., Olsson, J.C., Kjelleberg, S., Conway, P.L., 1991. Isolation and characterization of turbot (*Scophthalmus maximus*) associated bacteria with inhibitory effects against *V. anguillarum*. *Appl. Environ. Microbiol.* 57, 2223-2228.
- Westwater, C., Kasman, L.M., Schofield, D.A., Werner, P.A., Dolan, J.W., Schmidt, M.G., Norris, J.S., 2003. Use of genetically engineered phage to deliver antimicrobial agents to bacteria: an alternative therapy for treatment of bacterial infections. *Antimicrobial Agents and Chemotherapy.* 47, 1301-1307.
- Whitehead, N.A., Barnard, A.M., Slater, H., Simpson, N.J., Salmond, G.P.C., 2001. Quorum sensing in Gram-negative bacteria. *FEMS Microbiol. Rev.* 25, 365-404.
- Wiik, R., Andersen, K., Daae, F.L., Hoff, K.A., 1989. Virulence studies based on plasmid profiles of the fish pathogen *Vibrio salmonicida*. *Appl. Environ. Microbiol.* 55, 819-825.
- Wiik, R., Stackebrandt, E., Valle, O., Daae, F.L., Rodseth, O.M., Andersen, K., 1995. Classification of fish-pathogenic vibrios based on comparative 16S rRNA analysis. *Int. J. Syst. Evol. Microbiol.* 45, 421-428.
- Willems, A., Doignon-Bourcier, F., Goris, J., Coopman, R., Lajudie, P.D., Vos, P.D., Gillis, M., 2001. DNA-DNA hybridization study of Bradyrhizobium strains. *Int. J. Syst. Evol. Microbiol.* 51, 623-632.
- Williams, P., 2007. Quorum sensing, communication and cross kingdom signalling in the bacterial world. *Microbiology.* 153, 3923-3938.
- Winans, C.S., Bassler, B.L., 2002. Microbial psychology. *J. Bacteriol.* 184 873-883.
- Winans, C.S., 2004. Reciprocal regulation of bioluminescence and type III protein secretion in *Vibrio harveyi* and *Vibrio parahaemolyticus* in response to diffusible chemical signals. *J. Bacteriol.* 186, 3674-3676.

References

- Wong, H., Liu, C., Pan, T., Lee, C., Shih, D., 1999. Molecular typing of *V.parahaemolyticus* isolates, obtained from patients involved in food poisoning outbreaks in Taiwan, by random amplified polymorphic DNA analysis. *J. Clin. Microbiol.* 37, 1802-1812.
- Wong, H.C., Wang, P., 2004. Induction of viable but nonculturable state in *Vibrio parahaemolyticus* and its susceptibility to environmental stresses. *J. Appl. Microbiol.* 96, 359-366.
- Wong, H.C., Chang, C.N., 2005a. Hydrophobicity, cell adherence, cytotoxicity and enterotoxigenicity of starved *V.parahaemolyticus*. *J. Food. Prot.* 68, 154-156.
- Wong, H.C., Chen, C.H., Chung, Y.J., Liu, S.H., Wang, T.K., Lee, C.L., Chiou, C.S., Nishibuchi, M., Lee, B.K., 2005b. Characterization of new O3:K6 strains and phylogenetically related strains of *Vibrio parahaemolyticus* isolated in Taiwan and other countries. *J. Appl. Microbiol.* 98, 572-580.
- Xavier, K.B., Bassler, B.L., 2003. LuxS quorum sensing: more than just a numbers game. *Curr. Opin. Microbiol.* 6 191-197.
- Yamasaki, S., Shirai, H., Takeda, Y., Nishibuchi, M., 1991. Analysis of the gene of *Vibrio hollisae* encoding the hemolysin similar to the thermostable direct hemolysin of *Vibrio parahaemolyticus*. *FEMS Microbiol. Lett.* 80, 259-264.
- Yumoto, I., Iwata, H, Sawabe, T., Ueno, K., Ichise, N., Matsuyama, H., Okuyama, H., Kawasaki, K., 1999. Characterization of a Facultatively Psychrophilic Bacterium, *Vibrio rumoiensis* sp. nov., That Exhibits High Catalase Activity. *Appl. Environ. Microbiol.* 65, (1): 67-72.
- Yoshikawa, K., Adachi, K., Nishijima, M., Takadera, T., Tamaki, S., Harada, K., Mochida, K., Sano, H., 2000. Beta-cyanoalanine production by marine bacteria on cyanide-free medium and its specific inhibitory activity toward cyanobacteria. *Appl. Environ. Microbiol.* 66, 718-722.
- Zeigler, D.R., 2003. Gene sequences useful for predicting relatedness of whole genomes in bacteria. *Int. J. Syst. Evol. Microbiol.* 53, 1893-1900.
- Zhang, C., Yu, L., Quian, R., 2007. Characterization of OmpK, GAPDH and their fusion OmpK-GAPDH derived from *Vibrio*

References

- harveyi* outer membrane proteins: their immunoprotective ability against vibriosis in large yellow croaker (*Pseudosciaena crocea*). *J. Appl. Microbiol.* 103, 1587-1599.
- Zhang, W., Sun, K., Cheng, S., Sun, I., 2008. Characterization of DegQVh, a serine protease and a protective immunogen from a pathogenic *Vibrio harveyi* strain. *Appl. Environ. Microbiol.* 74, 6254-6262.
 - Zhang, X.H., Austin, B., 2000. Pathogenicity of *Vibrio harveyi* to salmonids. *J. Fish Dis.* 23, 93-102.
 - Zhang, X.H., Austin, B., 2005. Haemolysins in *Vibrio* species. *J. Fish Dis.* 23, 93-102.
 - Zhang, X.H., Meaden, P.G., Austin, B., 2001. Duplication of hemolysin genes in a virulent isolate of *Vibrio harveyi*. *Appl. Environ. Microbiol.* 67, 3161-3167.
 - Zhao, S., White, D.G., Ge, B., Ayers, S., Friedman, S., English, L., Wagner, D., Gaines, S., Meng, J., 2001. Identification and characterization of integron-mediated antibiotic resistance among Shiga toxin-producing *Escherichia coli* isolates. *Appl. Environ. Microbiol.* 67, 1558-1564.
 - Zhong, Y., Zhang, X.H., Chen, J., Chi, Z., Sun, B., Li, Y., B.Austin, 2006. Overexpression, purification, characterization and pathogenicity of *Vibrio harveyi* hemolysin VHH. *Infect. Immun.* 74, 6001-6005.
 - Zhu, J., Miller, M.B., Vance, R.E., Dziejman, M., Bassler, B.L., Mekalanos, J.J., 2002. Quorum-sensing regulators control virulence gene expression in *Vibrio cholerae*. *PROC. NATL. ACAD. SCI.* 99, 3129-3134.
 - Zobell, C.E., 1941. Studies on marine bacteria. The cultural requirements of heterotrophic aerobes. *J. Mar. Res.* 4, 42-75.
 - Zorrilla, I., Arijo, S., Chabrillon, M., Diaz, P., Martinez-Manzanares, E., Balebona, M.C., Morinigo, M.A., 2003. *Vibrio* species isolated from diseased farmed sole, *Solea senegalensis* (Kaup), and evaluation of the potential virulence role of their extracellular products. *J. Fish. Dis.* 26, 103-108.
 - Zuppardo, A.B., DePaola, A., Bowers, J.C., Schully, K.L., Gooch, J.A., Siebeling, R.J., 2001. Heterogeneity of environmental, retail,

References

and clinical isolates of *Vibrio vulnificus* as determined by lipopolysaccharide-specific monoclonal antibodies. *J. Food Prot.* 64, 1172- 1177.

* References not referred to in original.