REFERENCES

Abe, Y. (1979) Antitrypanosomal effect of acridine orange, sodium azide or high temperature on *Leptomonas* sp. *J. Seric. Sci. Japan*., **48:** 106-110.

Abe, Y. and Kawarabata, T. (1988) One the microsporidian isolates derived from the cabbageworm, *Piers rapae* crucifera, *J. Seic. Sci. Japan*., **57:** 147-150.

Allen, H.W. and Brunson, M.H. (1947) Control of Nosema disease of potato tuber worm, a host used in the mass production of *Macrocentrus ancylivorous*, Science., **105:** 394-395.

Alok Sahay, Singh, G.P.; Roy, D.K.; Sahay, D. N. and Suryanarayana, N. (2005) Effect of chemotherapy on Pebrine in Tasar silkworm, *Antherea mylitta* D. The 20th congress of the International Sericulture Commission 15th-18th December, 2005 Banagalore, India. Volume-II, Section 3: Non- Mulberry silkworm, pp: 126-130.

Ananthalakshmi, K. V. V.; Fujiwara, T. and Datta, R. K. (1994) First report on the isolation of three microsporidians (*Nosema* spp.) from the silkworm, *Bombyx mori* L. in India. *Indian J. seric.*, **2:** 146-48.

Andreadis, T.G. (1985a) Experimental transmission of a microsporidian pathogen from mosquitoes to an alternate copepod host. Proc. Natl. Acad. Sci., U.S.A. **82:** 5574-5577.

Andreadis, T.G. (1985b) Life cycle, epizootilogy and horizontal transmission of *Amblyospora* (Microspora: Amblyosporidae) in a univoltine mosquito, *Aedes stimulans*. *J. Invertbr. Pathol.*, **46:** 31-46.

Anonymous (2004) Annual report 2003-04, Central Silk Board. Banagalore. India.

Armstrong, E. (1976) Transmission and infectivity studies on *Nosema kingi* in *Drosophilla willistoni* and other drosophilids., *Z. Parasitenkd*., **50:** 161-165.

Avery, S.W. and Undeen, A.H. (1990) Horizontal transmission of *Parathelohania anophelis* to the copepod, *Microcylops varicans* and the mosquito, *Anopheles quadrimaculatus*. *J. Invertbr. Pathol*., **56:** 98-105.

Baig, M. (1994) Studies on *Nosema bombycis* N. - A pathogen of silkworm *Bombyx mori* L. Ph.D Thesis, University of Mysore, Mysore.

Baig, M.; Ananthalakshmi. K. V. V.; Sasidharan, T.O. and Fujiwara. T. (1997) "*Manual on Microsporidian Disease and its Control*" pp. 1-28.

Baig, M.; Sasidharan, T. O.; Sharma, S. D.; Sen, S. K. and Jolly, M. S. (1989) Efficacy of certain bed disinfectants against nuclear polyhedrosis of silkworm, *Bombyx mori* L. *Indian.J. Seric.*, **33:** 214-221.

Bailey, L. (1963). "*Infectious diseases of honey bees*" Land Books, London.

Baker, M. D.; Vossbrinck, C.; Maddox, J.V. and Undeen, A. H. (1994) Phylogenetic relationships among *Vairimorpha* and *Nosema* species (Microspora) based on ribosomal RNA sequence data. *J. Invertbr. pathol*., **64:** 100-106.

Balavenkatasubbaih, M.; Ananthalakashmi, K.V.V.; Selavakumar, T.; Nataraju, B. and Datta, R.K. (1999) Chlorine dioxide, a new disinfectant in sericulture, *Indian.J. Seric.,* **38:** 125-130.

Balavenkatasubbaih, M.; Hyasaka, S. and Nataraju, B. (2003) Efficacy of "Kao Haiter" as a disinfectant against the pathogens of the silkworm, *Bombyx mori* L. *Advance in Tropical Sericulture*., pp 326-329

Balavenkatasubbaih, M.; Nataraju, B.; Sharma, S.D.; Selavakumar, T.; Chandrasekharan K. and Sudhakara, Rao. (2006) Serichlor, A disinfectant in Indian Sericulture, *Int. J. Indust. Entomol.*, **12:** 7-14.

Baribeau, M.F and Burkhardt, C. C. (1970) Effect of heat and ultraviolet rays on *Nosema apis* spores in relation to honey bee infection. *J. Kansa Entomol. Soc*., **43:** 455-458.

Bauer, L, S. and Pankratz, H. S. (1993) *Nosema scripta* N. Sp. (Microsporidia : Nosematidae), a Microsporidian parasite of the cottonwood leaf beetle *Chrysomela scripta* (Coleoptera : Chrysomelidae), *J. Euk. Microbio.,* **40:** 135-141.

Becnel, J.J. and Andredis, T.G. (1999) Microsporidia in Insects; in the *microsporidia and microsporiodiosis*. Wittner, M and L. M. Weiss (eds.) pp. 447-501. ASM press, Washington D.C.

Becnel, J.J. and Fukada, T. (1992) Ultra structure of *Culicosporella lunata* (Microsporidia: Culicosporellidae famn) in mosquito *Culex pilosus* (Diptera: Culicudae) with new information on the developmental cycle. *Eur. J. Protistol*., **26:** 319-329.

Brooks, W. M.; Cranford, J. D. and Pearce, L. W. (1978) Benomyle: Effectiveness against the microsporidian *Nosema heliothis* in the corn earworm, *Helithis zea*. *J. Invertebr. Pathol*., **31:** 239-245.

Brooks, W. M. (1988) Entomogenous protozoa. *In* C. M. Ignoff (ed.) CRC Handbook of natural pesticides. Microbial Insecticides. Part A Entomogenous Protozoa and Fungi. CRC Press. Boca raton Florida.

Burges, H. D.; Canning, E.U. and Hulls, I.K. (1974) Ultra structure of *Nosema oryzaephilli* and the taxonomic value of the polar filament, *J.Invert. Path.*, **23:** 135-139.

Canning, E. U and Vavra, J. (2000) Phylum microsporidia. I The Illustrated Guide to the Protozoa 2nd ed. (J.J. Lee, G.F. Leedale and P. Brad, eds.) Vol. 1, pp. 39-126. Allen press, Lawrence, K.S.

Canning, E.U. (1990). Phylum Microspora. *In* " Handbook of Protoctista." (L. Margulis, J. O. Corliss, M. Melkonian, and D. J. Chapman. Eds.), pp. 53-72. Jones and Bartlett, Boston.

Canning, E. U. and Curry, A. (2004) Further observations on the Ultrastructure of *Cystosporogenes operophtera brumata* L. (Lepidoptera, Geometridae) *J.Invert. Path.*, **87:** 1-7.

Cantwell, G. E. and Shimanuki, H. (1969) Heat treatment as means of eliminating Nosema and increasing production, *Am. Bee. J*., 109: 52-54.

Cantwell, G. E. (1974) Insect diseases. Vol. II, Marcel Dekkar Inc. New York, pp. 501-547.

Chinnaswamy, K. P. and Devaiah, M. C. (1984) Susceptibility of different races of silkworm *Bombyx mori* L. to aspergillosis caused by *Aspergillus tamarii* Katta. *Sericolgia.,* **24:** 513-517.

Chitra, C. N.; Karanth, C. K. and Vasantharjan, V. N. (1975) Diseases of mulberry silkworm, *Bombyx mori* L. *J. scientific and industries res*., **3:** 386-410.

Choi, J. Y.; Kim, J. G.; Choi, Y.C.; Goo, T. W.; Chang, J. H.; Je, Y. H. and Kim, K. H. (2002) *Nosema* sp. isolated from cabbage white butterfly (Pieris rapae) collected in Korea, *The J. of Microb.,* **40:** 199-204.

Choudary, S. N. (1967) The silkworm and its culture. Mysore Printing and Publishing House, Mysore, India, pp. 76.

Colbourn, N. I.; Hollister, W. S.; Curry, A. and Canning, E. U. (1994) Activity of albendazole against *Ecepphalitozoon cuniculi* in vitro. *Eur. J. Protistol*., **30:** 211-220.

Dandin, S. B. (2006) Production and supply of quality Bivoltine eggs to meet the set target of raw silk production, "*Workshop on Pebrine*" Disease management in south India, 6th March-2006, pp. 12-15.

Dash, A. K and Nayak, B. K. (2003) Seasonal variation of the parasitic effect of *Nosema* sp. on reproductive success of Indain tasar silk insect *Antheraea myltta* Drury (Saturnidae) *Bull. Ind. Acad. Seri*., **7:** 17-20.

Datta, R. K. (1992) *Guidelines for bivoltine rearing*, Central Silk Board, Banglore, India.

David, Oi, H.; Briano, J.A.; Vallea, S. M. and Williams, D.F. (2005) Transmission of Vairimorpha invictae (Microsporidia : Burenellidae) infections between red imported fire ant (Hymenoptera :Formicidae) Colonies. *J.Invert. Path.,* **88:** 108-115.

Devaiah, M. C. (1975) Studies on the pebrine disease of the silkworm, Ph.D., thesis, University of Agricultural Sciences, GKVK., Bangalore, India.

Devaiah, M. C.; Thontadaraya, T. S. and Krishnaswami, S. (1975) Susceptibility of different races of silkworm *Bombyx mori* L. to pebrine disease. *Current Res*., **4:** 99.

Fang, D.; Liao, S.; Zheng, M.; Zhu.D.; Chen.H.; Liu.J.; Qil.; Luk.; Nong. Z.; Bai. B.; Mao. K. and Chen. J. (1991) Study on the two microsporidians (MG1, MG2) from the silkworm, *Bombyx mori*, *Guangdong Agricultural Sci*., **2:** 35-38.

Finney, D. J. (1971) Estimation of the median effective dose, In: *Probit analysis*.Third Edition. S. Chand & Company Ltd., New Delhi, pp. 20-49.

Frankenhuyzen, K. V.; Ebling. P.; McCron, B.; Ladd, T.; Gauthier, D. and Vossbrinck, C. (2004) Occurrence of *Cystosporogenes* sp (Protozoa, Microsporidia) in multi species insect production facility and its elimination from a colony of the eastern spruce budworm, *Choristoneura fumiferana* (Clem.) (Lepidoptera : Tortricidae). *J.Invert. Path.,* **87:** 16-28.

Fujiwara, T. (1993) Comprehensive report on the silkworm disease control. A report on "Bivoltine Sericulture Technology Development in India" submitted to Central Silk Board, Bangalore, India, pp. 110.

Fujiwara, T. (1980) Three microsporidians *(Nosema spp.)* from the silkworm *Bombyx mori*. *J.Seric. Sci. Japn.,* **49:** 229-236 (in Japanese with English summary).

Fujiwara, T. (1985) Microsporidian fron silkmoths in egg production sericulture. *J. Seric. Sci. Japn*., 54: 108-111 (in Japanese with English summary).

Fujiwara, T. (1984a) A Pleistophora like microsporidian isolated from silkworm, *Bombyx mori* L. *J. Seric. Sci. Jpn* ., **53:** 398-402.

Fujiwara, T. (1984b) Thelanhania spp. (Microsporidia : Thelanhaniidae) isolated from silkworm, *Bombyx mori* L. *J. Seric. Sci. Jpn* ., **53:** 459-460.

Fujiwara, T.; Ogimoto, T.; Sudo, K. and Ueda, K. (1966) Immunoflurescence study of *Nosema bombycis* Naegeli, a causative agent of pebrine. *Acta Sericol*., **58:** 14-19.

Girijidevi, A. (2006) Studies on the development of botanical based suppression / control of *Nosema bombycis* infection in the silkworm. *Bombyx mori* L. M. Sc. Dissertation NGM college (Autonomous), Pollachi (Tamil Nadu), Affiliated to Bharathiar University.

Goertz, D.; Pilarska, D.; Kereselidze, M.; Solter, L. F.; and Linde, A. (2004) Studies on the impact of two *Nosema* isolates from Bulgaria on the gypsy moth (*Lymantria dispar* L), *J. Invert. Path.*, **87:** 105-113.

Govindan, R; Narayanaswamy, T. K.; Devaiah, M. C. (1998) Principles of Silkworm Pathology Ser. Scientific Publishers, Bangalore, India, pp. 420.

Griyaghey, U.P.; Jolly, M.S. and Kumar, P. (1976) Studies on the thermic control of microsporodiosis of the tropical Tasar silkworm, *Antherea mylitta* D. *Ind. J. Seric*., **14:** 27-30.

Han, M. S. and Watanabe, H. (1988) Transovarial transmission of two microsoridia in the silkworm, *Bombyx mori*, and disease occurrence in the progeny population. *J. Invertebr. Pathol*., **51:** 41-45.

Hanumappa, H. G. (1986) Sericulture for rural development, Himalaya Publishing House, Bombay, India, p. 152.

Haque, M. A. and Haberkom, A. (1993) Possible mode of action of toltrazuril; studies on the *Eimeria* species and mammalian and *Ascar*is suum enzyme, *Parasitol. Res*., **76:** 8-12.

Hatakeyama, Y.; Bansal, A. K.; Iwano, H.; Kawakami, Y.; and Ishihar, R. (2000) Characterzation of ssu-rRNA sequence of a new microsporidium *Nosema* sp. (Nosematidae: Microsporidia), isolated from *Antheraea mylitta* Drury (Lepidoptera :Saturniidae) in India. *Indian J. seric.*, **39:** 131-134.

Hayasaka, S. (1991) Inhibitory effects of antimicrobial chemicals on the sporogenesis of *Nosema bombycis* infecting larvae of silkworm, *Bombyx mori* L. *Acta. Seric. Entomol.*, **4:** 53-54.

Hayasaka, S.; Nozaki, M.; Yasunaga-Aoki, C.; Teramoto, N.; Iiyama, K.; Hatakeyama, Y.; Hashiguchi, M.; and Kawarabata, T. (2002) A new microsporidian occurring in larvae of *Antheraea pernyi* (Lepidoptera:

Saturniidae) released on an oak tree, *Quercus acutissima* and a possible insect-vector, *Lepidogma kiiensis* (Lepidoptera: Pyralidae) inhibiting the same plant, *Sericologia*., **42:** 107-111.

Hazard,E I. and Weiser, J. (1968) Spores of *Thelohania* in adult female *Anopheles*: Development and transovarial transmission and redescriptions of *T.legeri* Hesse and *T. obesa* Kudo. *J. Protozool*., **15:** 817-823.

Hsiao, T. H. and Hsiao, C. (1973) Benomyl: anovel drug for controlling a microsporidian disease of alfalfa weevil. *J. Inverterbr. Pathol.*, **22:** 303-30.

Howard, (1925) Cf. Liu Shi Xian and Zhong., W.B. (1988).The research channels in the prevention and control of silkworm diseases. *Sericolgia*. **29:** 287-295.

Hylis, M.; Pilarska, D. K.; Obornik, M.; Vavra, J.; Solter, L. F.; Weiser, Linde, A. and McManus, M.L (2006) *Nosema chysorrhoea* N. sp. (Microsporidia), isolated from brown tail moth (*Euproctis chysorrhoea* L) (Lepidoptera, Lymantriidae) in Bulgaria: Characterisation and phylogenetic relationships. *J. Invertebr. Pathol*., **91:** 105-114.

Inoue, H. (1977) Thermal therapy of virus diseases of the silkworm *Bombyx mori*. *J. Seric. Sci. Japn*. **46:** 306-312.

Ishihara, R. (1968) Some observations on the fine structure of sporoplasm discharged from spores of microsporidian, *Nosema bombycis*. *J. Invertbr, patholo*., **12:** 245-258.

Ishihara, R. and Fujiwara, T. (1965) The spread of pebrine in a colony of silkwom, *Bombyx mori,* L. *J. Invertebr. Patol.,* **7:** 126-131.

Issi, I.V. (1986) Microsporidia phylum of parasitic protozoa, *Protozoology*., (Leningrad) **10:** 6-136.

Iwano, H and Isihara, R. (1981) Inhibition effect of several chemicals against the hatch of *Nosema bombycis* spores. *J. Seric. Sci. jpan*. **50:** 276-281.

Iwano, H and Isihara, R. (1991) Isolation of *Nosema bombycis* from the moths of the lawn grass cutworm. Spodoptera depravata Butler., *J. Seric. Sci. jpan*. 6**0:** 279-287.

Iwashita, Y.; Suzuki, T.; Nakazato, S. and Kobayashi. H. (1990) Multiplication and tranovarial transmission of three microsporidians infecting the silkworms, *Bombyx mori*. *Tech. Bull. Seric. Sci. Inst*., **38:** 35- 43. (in Japanese).

Iwashita, Y. and Zhou, C. D. (1988) Inactivation by treatment of a polyhedrosis virus of the silkworm, *Bombyx mori* L. with calcium hydroxide solution. *J.Seric.Sci.japan*., **57:** 511-518.

Jaiswal, K and Deka, G.M. (2005) study of protein in the leaf of medicinal plant extract for controlling different diseases in *Bombyx mori*. The 20th congress of international sericulture Commision Vol. III Sec. II pp 38-42.

Jankiraman, A. T. (1961) Disease affecting the Indian silkworm races, *J. Silkworm.*, **13:** 91-101.

Jesper, A. B and Ward, M. A. De. (1993) Natural products in plant protection. Netherland, *J. Pl. Path*., **99:** 109-117.

Johny, S. and Muralirangan, M. C. (2005) A microsporidian (Protozoa: Microspora) Infecting *Spodoptera litura* (Fabricius) (Lepidotera: Noctutuidae) in India: its morphology, host range and efficacy. *Entomon*., **30:** 147-155.

Kagawa, T. (1980) The efficacy of formalin as a disinfectant of *Nosema bombycis* spores. *J.Seric. Sci. Japan*., **49:** 218-222.

Kalaivani and Anathalakshmi, R. (2003) Effects of aqueous neem extracts on pebrine in *Spodoptera*. *Natinal Confrence on Tropics.*, pp. 95

Kaya, H. K. (1977) Survival of spres of *Vairimorpha* (*Nosema necatrix*) (Microsporidia: Nosematidae) exposed to sunlight, ultraviolet radiation and high temperature. *J. Invertebr. Pathol.,* **30:** 192-198.

Kawarabata, T. (2003) Biology of Microsporidians infecting the silkworm, *Bombyx mori*, in Japan, Journal of *Insect Biotecnology and Sericology.,* **72:** 1-32.

Kawarabata, T. and Hayasaka, S. (1987) An enzyme-linked immunosorbent assay to detect alkali-soluble spore surface antigens of strains of *Nosema bombycis* (Microspora - Nosematidae). *J. Invertebrt. Pathol*., **50:** 118-123.

Kawokami, Y.; Inoue, T.; Ito, K.; Kitamizu, K.; Hanawa, C.; Sunairi, M.; Ando, T.; Iwano, H. and Ishihaara, R.. (1994) Comparison of Chromosomal DNA from four microsporidia pathogenic to the silkworm, *Bombyx mori. Appl. Entomol. Zool.*, **29:** 120-123.

Keeling, P. J. and Fast, N. M. (2002) Microsporidian: biology and evolution of highly reduced intracellular parasites. *Annu. Rew. Microbial.*, **56:** 93-116.

Keeling, P. J.; Fast, N. M.; Law , J. S.; Williams, B. A. P. and Slamovits, C. H. (2005) Comparative genomics of microsporidia. *Folia Parasitologica.*, **53:** 8-14.

Kellen, W. R.; Chapman, J. E.; Clark, T.B.; and Lindegren.J. E. (1965) Host-parasite relationships of some of the *Thelohania* from mosquitoes (Nosematidae: Microsporidia). *J. Invertebr. Pathol*., **51:** 161-166.

Kellen, W. W. and Lindegren, J.E. (1971) Mode of transmission of *Nosema Plodiae* Kellen and Lindegren, a pathogen of *Plodia interpunctella* (Hubner) *J. stored Prod. Res.*, **7:** 31-34.

Keohan, E. M. and Weiss, L.M. (1999) The structure, function and composition of microsporidian polar tube, In *The Microsporidia and Microsporopdiosis*, p. 196-224. ASM Press, Washington, D. C.

Kfir, R. and Walter, H. S. (1997) Control of microspoprdian disease in laboratory cultures of stem borer *Chilo partellus* (Lepidoptera: Pyralidae) *Afr. Entomol*., **5:** 155-172.

Kishore, S.; Baig. M.; Nataraju, B.; Balavenkatasubbaiah, M.; Sivaprasad, V.; Iyengar, M. N. S. and Datta, R.K. (1994) Cross infectivity microsporidians isolated from wild Lepidopterous insects to silkworm, *Bombyx mori* L. *Indian J. Seric*., **33:** 126-130.

Kobayashi, H. and Yamazaki, D. (1987) Discrimination of microsporidian spores with fluorescent antibody techniques in the insects, *Gunma J. of Agric. Res*., Series B, **4:** 25-28.

Kramer, M. (1976) the extra-corporeal ecology of microsporidia. In "Comparative Pathobiology" (L. A. Bulla, Jr. and T. C. Cheng, eds.), Vol. 1, pp. 127-135. Plenum Pres, New York.

Kuhn, D. (1988) Textile technology: Spinning and reeling. In *Science and Civilization in China*. Part XI, edited by J. Needham, Cambridge: Cambridge University Press.

Kumar, S.; Sigh, A and Sharma, A. (1999) Asian Region Inventory of Medical and aromatic plant and polyherbal formulation. Department of Biotechnology, New Delhi, India, pp. 91.

Laemmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature.*, **227:** 680-685.

Lewis, L, C. and Lynchi, R.E. (1970) Treatment of Ostrinia nubilalis larvae with Fumidil B to control infections caused by *Perezia pyraustae*. *J. Inverterbr. Pathol.*, **15:** 43-48.

Li Aiqun, (1962) The proceedings of Confrence of Sericulture community, China, p. 198.

Lim, J.S.; Lee, Y.K.; Cho, S.Y. and Han, M. S. (1982) Characteristics of a New Microsporidian S80 Isolated from the silkworm, *Bombyx mori* in *Korea*. *Res. Proj. Rep., Korean sericult. Assoco., Seoul.,* (in Korean with English summary).

Liu Shixian, (1972) Cf. Liu Shi Xian and Zhong., W.B. (1988).The research channels in the prevention and control of silkworm diseases. *Sericolgia*. **29:** 287-295.

Liu, (1987). Cf. Liu Shi Xian and Zhong., W.B. (1988).The research channels in the prevention and control of silkworm diseases. *Sericolgia*. **29:** 287-295.

Liu Shi Xian. and Zhong, W.B. (1980) Cf. Liu Shi Xian and Zhong., W.B. (1988).The research channels in the prevention and control of silkworm diseases. *Sericolgia*. **29:** 287-295.

Liu-Yaung, (1960). Cf. Liu Shi Xian and Zhong., W.B. (1988).The research channels in the prevention and control of silkworm diseases. *Sericolgia*. **29:** 287-295.

Liu, S. X. (1984) Identification on the resistance of silkworm *Bombyx mori* races to six types of silkworm diseases. *Sericologia.,* **24:** 377-382.

Liu, S. X. (1980) The effect of Chemotherapy on pebrine disease of *Bombyx mori*. *Sericologia.,* **27:** 495-410.

Lom, J. and Vavar, J. (1963a) The mode of sporoplasm extrusion in microsporidian spores. *Acta. Porotozool.,* **1:** 81-89

Lu Heming.; Cf. Liu Shi, Xian. and Zhong, W. B. (1981) The research cahhels in the prevention and control of silkworm diseases. *Sericologia*., **29:** 287-295.

Lu Yuling, (1960) Cf. Liu Shi Xian and Zhong., W.B. (1988).The research channels in the prevention and control of silkworm diseases. *Sericolgia*. **29:** 287-295.

Lu-haung (1984) Cf. Liu Shi Xian and Zhong., W.B. (1988).The research channels in the prevention and control of silkworm diseases. *Sericolgia*. **29:** 287-295.

Maddox, J.V. (1973) The persistence of the microsporidia in the environment. *Mis. Pub. Entomol. Soc. Amer*., **9:** 99-104.

Maddox, J.V. (1977) Stability of entomopathogenic protozoa. Misc. Publ. *Entomol. Soc. Amer*., **10**: 3-8.

Masera. (1938) Azione biologica di metallic sulle uova di *Bombyx mori* L. infetti di pebrina (nota preliminera). *Mem. R. Accad. Sci. Arti. Padova*., **54:** 7.

Matos, E. and Azevedo, C. (2004) Ultrastructure of Microsporidium brevirostris sp. n., parasite of the Teleostean Brachyhypopomus brevirostris (Hypopomidae) from Amazon River. *Acta protozoologica*., **43:** 261-267.

Miyajima, S. (1979) Effect of some disinfectants on the viruses of silkworm, *Bombyx mori* L. *Res. Bull. Achi. Agric. Res. Centre.*, **11:** 165-168.

Mohan, Madana, N.; Krishnan, N.; Mitra, P.; Das, N.K.; Saratchandra, B. and Haldar, D,P. (2005) Seasonal impact of microsporidian infection on the reproductive potential of silkworm, *Bombyx mori* L. (lepidoptera: Bombycidae) *Int. J. Indust. Entomol.*, **11:** 107-111.

Mozziconacci, A. (1921) Le ber a Soiedu Murier, Libraivie Hachette, Paris.

Nageswara Rao.; Muthulaskshmi.; M. Kanginakudra.; S. and Nagraju, J. (2004) Phylogenetic relationships of three microsporidian isolates from silkworm, *Bombyx mori*, *J. Invertebrt. Pathol*., **3:** 87-95.

Nageswara Rao.; Surrendra Nath, B. and Sarachandra, B. (2005) Characterisation and phylogenetic relationships amoung microsporidian infecting silkworm, Bombyx mori L, using inter simple sequence repeat (ISSR) and small subunit rRNA (SSu-rRNA) sequence analysis, *Genome*., **48:** 355-366.

Nagraju, J. G.; Klimenko, V. and Couble, P. (2001) The silkworm, *Bombyx mori*: A model genetics system, *Encycloedia of Genetics*. Edited by Eric, C, Reeve and Isobel Black, Fitzroy Dearborn publishers London: Chiago.

Nataraju, B. and Dandin, S.B. (2006) Recent trends in Pebrine disease management in Silkworm, "*Workshop on Pebrine*" Disease management in south India, 6th March-2006, pp: 21-27.

Nataraju, B.; Sathyaprasad.; Manjunath, D. and Aswani Kumar, C. (2005) A text book on Silkworm crop protection (Central Silk Board) pp. 1-320.

Nataraju, B.; Balavenkatasubbaih, M.; Sharma, S.D.; Selavakumar, T.; Thiagarajan, V. and Dandin, S. B. (2002) A practical technology for diagonosis and management of diseases in silkworm rearing., *Int. J. Entomol.* **4:** 169-173.

Nataraju, B. (1995) Studies on Diagnosis and prevention of Nuclear polyhedrosis in silkworm *Bombyx mori* L. Ph.D Thesis, University of Mysore, Mysore.

Nathan, S. and Kalaivani. (2005) Efficacy of Azidirachtin on pebrine in *Spotoptera*. *Biological Contro*l **17:** 95 95.

Olsen, P.E.; Rice, W.A. and Liu, T.P. (1986) In Vitro germination of *Nosema apis* spores under conditions favorable for generation and maintenance of sporoplasm. *J. invertebr. Patohlol*., **47:** 65-73.

Oshima, K. (1964) Effect of potassium ion on filament evagination os spores of *Nosema bombycis* as studied by neutralization method. *Annot. Zool. Japn*., **37:** 102.

Oumouna, M.; Matbouli, M. El. Hoffmann, R.W. and Bouix, G. (2000) Electron microscopic study of a new microsporean *Microsporidium epithelialis* sp. n. infecting *Tubifex* sp. (Oligochaeta) *Filia parasitologica*., **47:** 257-265.

Ovcharenko, M. and Wita, I. (2005) The ultrastructure of *Nosema artemiae* (Codreanu, 1957) (Microsporidia : Nosematidae) *Acta protozoologica*., **44:** 33-41.

Overstreet, R. M. (1975) Buquinolate as a preventive drug to control microsporidiosis in blue crab. *J. Invertebrt. Pathol*., **26:** 213-216.

Pasteur, L. (1870) Etude sur la maladie des vers a soie. Tome 1, p.322, Tome ll, p.327, Gauthier-Villars, Paris.

Peter, A.; Sadatulla, F. and Devaiah, M. C. (1999) The viral, bacterial and protozoan diseases of the silkworm *Bombyx mori* L.; In Advances in mulberry sericulture, Devaiah, M. C Narayanasawamy, K.C. and Maribashetty, V, G. (eds.) pp. 378-457. CVG publications, banagalore, India.

Patil, C. S. (1991) Studies on the evaluation of calcium hydroxide against cytoplasmic polyhedrosis of the silkworm, *Bombyx mori* L *Entomon*., **16 :** 147-150.

Patil, C.S. and Geethabai. (1989) Studies on the susceptibility of silkworm races to pebrine spore *Nosema bombycis* N. *J.Appl. Entomol*., **108:** 421-425.

Petri, M. (1969) Studies on the *Nosema cniculi* found in the transplantable aseites tumors with a survey of microsporodiosis in mammals. *Acta. Pathol. Microbiol. Scand. Suppl*., **204:** 1-9.

Petri, M. and Shiodt, T. (1966) On the ultra structure of *Nosema cuniculi* in the cells of the Yoshida rat *Ascites sarcoma*, *Acta. Pathol. Microbiol. Scand*., **66:** 437-446

Ramegowda, T. and Geethabai, M. (2005) Transovarian transmission of Pebrine spores, *Nosema bombycis* Naegeli in the leaf-roller pest of mulberry, *Diaphania pulverulentalis*, *Sericologia*., **45:** 137-141.

Raun, E. S. (1961) Elimination of microsporidiosis in laboratory reared European corn borers by the use of heat. *J. Insect. Pathol.,* **3:** 446-448.

Rice, R, N. (2001) A report for Rural Industries Research and Development Corporation. *Nosema* disease in honey bees (Genetic variation and control), RIRDC Publication. pp. 1-36.

Sahaf, K. A. (2002) Incidence of protozoan disease (pebrine) of silkworm, *Bombyx mori* L. in Jammu and Kashmir. *J. Ent. Res.,* **26:** 305-307.

Samson, M. V.; Santha P.C.; Singh R. N. and Sasidharan T.O. (1999a) A new microsporidian infecting *Bombyx mori* L. *Indian Silk.,* **37:** 10-12.

Samson, M. V.; Santha P.C.; Singh R. N. and Sasidharan T.O. (1999b) Microsporidian spore isolated from Pieris sp. *Indian Silk.,* **38:** 5-8.

Sasidharan, T. O.; Singh, R. N. and Santha, P. C. (2003) Studies on the microsporidiosis of silkworm, *Bombyx mori*, L. *Annual Report* - SSTL Banagalore, pp. 1-3.

Sato, R. and Watanabe, H. (1980) Purification mature Microsporodian spores by isodensity equilibrium centrifugation, *J. Sericult. Sci.Japan.,* **49:** 512-516.

Sato, R., Kobayashi, M.; Watanabe, H. and Fujiwara, T. (1981) Serological discrimination of several kinds of microsporidian spore isolated from the silkworm, *Bombyx mori*, by an indirect fluorescent antibody technique. *J. Seric. Sci. Jpn.,* **50:** 180-184.

Savitri (2006) Studies on the development of thermotherapy against inctious flacherie in silkworm *Bombyx mori* L. M. Sc. Dissertation: NGM college (Autonomous), Pollachi (Tamil Nadu), Affiliated to Bharathiar university.

Schimaha, G. and Benini, J. (1998) Treatment of fish parasites. 11. Effects of different benzimidazole derivatives (albendazole, mebendizole, febendazole) on *Glugea anomala*, 1887 (Microsporidia) Ultrastructure aspects and efficacy studies, *Parasitol. Res.,* **84:** 41-49.

Scholtyseck, E. and Danneel, R. (1962) Uber die Feinstruktur der spore von *Nosema apis*, *Dtsch. Entomol. Z.,* **9:** 471-476.

Selvakumar, T.; Nataraju, B.; Balavenkatasubbaiah, M.; Shivaprasad, V.; Baig, M.; Virendra Kumar.; Sharama, S. D.; Thaigarajan, V. and Datta, R.K. (2002) A report on the prevalence of silkworm diseases and estimated crop loss.

Proceedings of national conference on strategies for sericulture research and development 16-18 Nov. 2000, PP. 354-357 edited by S.B. Dandin and V.P. Gupta published by CSR and TI (CSB) Mysore.

Selvakumar, T.; Nataraju, B.; Chandrasekharan, K.; Sharma, S.D.; Balavenkatasubbaiah, M.; Sudahakara, Rao.; Thiagarajan, V. and Dandin, S.B. (2005) Isolation of a new microsporidian sp. (NIK-5hm) forming spores within the haemocytes of silkworm, *B. mori* L. *Int. J. Indust. Entomol.*, **11:** 63-66.

Sharma, S. D.; Balavenkatasubbaiah, M.; and Baig, M. (1989) A report on the presence of various pathogenic microbes in a wild population of Bihar hairy caterpillar, *Diacrasia oblique*. *Curent Sci.*, **58:** 762-763.

Sharma, S. D.; Chandrasekharan, K.; Nataraju, B.; Balavenkatasubbaiah, M.; Selvakumar, T.; Thiagarajan, V. and Dandin, S.B. (2003) The cross infectivity between a pathogens of silkworm, *Bombyx mori* L. and mulberry leaf roller, *Diaphania pulverulentalis* (Hampson). *Sericologia*., **43:** 203-209.

Sheeba, R.; Devaiah, M.C.; Chinaswamy, K.P. and Govindan, R.K. (1999) Thermotherapy of pebrinized cocoons and its effect on larval progeny; in *Proc. Natl. Semi. Trop. Seric.* Govindan, R.K., K.P. Chinnaswamy., N.K. Krishnaprasad and D.N.R. Reddy (eds.), Vol. II., pp.266-272, UAS and Swiss Agency for Development and Cooperation, Bangalore, India.

Shi, L. and Jin, W. (1997) Study on the differential diagnosis of *Nosema bombycis* of the silkworm, *Bombyx mori*, by monoclonal antibody-sensitized latex., *Sericologia*., **1:** 1-6.

Sidorov, N. G.; Koptev, V.S. and Mugalimov, I. A. (1975) Resistance of honeybees to *Nosema* disease and a genetic method of control. *Veterinariya Mosco.*, **7:** 63-65.

Singh, G.P.; Alok Sahay Roy, D,K.; Sahay D, N. and Suryanarayana, N. (2005a) Screening of medicinal plants against Virosis in Tasar silkworm, *Antherea mylitta* D. The 20th congress of the International Sericulture Commission 15th - 18th December 2005 Banagalore, India. Volume-II, Section 3: Non-Mulberry silkworm, pp:131-135.

Singh, G.P.; Alok Sahay. Roy. D, K.; Sahay D.N. (2005b) Efficacy of disinfectants against Cytoplasmic polyhedrosis virus and Microsporidia of Tasar silkworm, *Antherea mylitta* D. *Int. J.Indust. Entomol*., **10:** 69-72.

Singh, G.P.; Sharma, S.D.; Selvakumar, T.; Nataraju, B. and Datta, R.K.(2002) Screening of fungicides and natural plant products and their efficacy on the

control of Aspergillosis in silkworm *Bombyx mori* L.; *Int. J. Indust. Entomol.*, 4**:** 5-11.

Singh, T. and Saratchandra, B. (2003) Microsporidian disease of the silkworm *Bombyx mori* L. (Lepidoptera : Bombycidae) *Int. J. Indust. Entomol.*, **6:** 1-9.

Sokolova, Y. K and Lange, C. E. (2002) An ultrastructural study of Nosema locustae Canning (Microsporidia) from three species of *Acrididae* (Orthoptera). *Acta protozoologica.*, **41:** 229-237.

Sprague, V. (1977) Classification and Phylogeny of microsporidia. *In* "Comparative Pathobiology. Systematics of the Microsporidia" (L.A. Bulla, Jr. And T.C. Cheng, eds.), Vol.2, pp1-30. Plenum Press, New York.

Sprague, V. (1982). Microspora. *In* " Synopsis and Classification of Living organisms*"* (S.P Parker,ed.) Vol. 1.pp589-594, McGraw-Hill, New York.

Sprague, V.; Becnel, J.J. and Hazard, E.I. (1992) Taxomony of phylum Microspora. *Critical Review in Microbiology.*, **18:** 285-395.

Steinhaus, E. A. (1949) Protozoan infections. In: *Priciples of the insect pathology*, McGraw-Hills Book Company, INC., New York. Pp. 592-602.

Steinhaus, E. A. (1956) The effect of disease on insect poulation, *Hilgardia*, **23:** 193-216.

Sujatha, K.; Purusotham, A. R. and Sammaiah, Ch. (2005) Turmeric - nature's precious gift as bed disinfectant against silkworm diseases and its effect on growth and cocoon characters. The 20th congress of the International Sericulture Commission 15th-18th December, 2005 Bangalore, India. Volume-II, Section 3 : Non-Mulberry silkworm, pp: 371-374.

Sweeney, A.W.; Hazard, E.I. and Graham, M. F. (1985) Intermediate host for an Amblyospora sp. (Microspora) infecting mosquito, *Culex annulirostris*. *J. Inverttebr. Pathol.*, **46:** 98-102.

Tanada, Y. and Kaya, H.K. (1993) Protozoan infection: Apicomplexa. Micospora, In "*Insect Pathology*" pp 414-458. Academic press, Inc. San. Diego.

Tanaka, S.; Shimizu, T.; Kobayashi, M., and Ishihara, R. (1972) A new microsporidian pathogenic to the silkworm. *Bombyx mori. J. Seric. Sci. Japn.*, **41:** 89-95 (in Japanese with English summary).

Tatsuke, K. (1971) Pebrine disease of Silkworm a technical report, Overseas Technical Cooperation agency, Tokyo, Japan, pp. 1-2.

Thomas, Bartram (1998) Encyclopedia of herbal medicine (Vol. I) Marlove and company, New York.

Toguebaye, B.S. and Bouix, G. (1989) *Nosema galerucella* sp. n. Microsporidian (Protozoa, Microspora) parasite of *Galerucella luteola* Muller (Chrysomelidae, Coleoptera); Development cycle and ultrastructure. *Europ. J. Protistol*., **24:** 346-353.

Toguebaye, B.S. and Marchand, R.E. (1986) Etude d'une infection microsporidienne due a *Nosema birgli* n. sp. (Microsporida, Nosematidae) chez, *Mesoplatys cincta* Olivier, 1790 (Coleoptera, Chrysomelidae). *Z. Parasitenkd*., **72:** 723-737.

Toguebaye, B.S. and Marchand, R.E. (1989) Observiation en microscopie electronique a transmission des stades de developpement de *Nosema nisotrae* sp. n (Microsporida, Nosematidae) parasite de *Nisotra* sp. (Coleoptera, Chrysomelidae). Development structure and Ultrastructure. *Europ. J. Protistol*., **24:** 346-353.

Toguebaye, B.S. and and Marchand, R.E. (1984) Etude histopathologique et cytopathologique d'une microsporidiose naturelle chz la coccinelle des cucuabitacees d' Afrique Henosepilachna elaterii [Col.: coccinellidae] *Entomophaga*., **29:** 421-429.

Tsai, S. J.; Lo. C.H.; Soichi, Y. and Wang, C. H. (2003) The Characterisation of microsporidian isolates (Nosematidae: Nosema) from five important lepidopteran pests in Taiwan. *J. Invertebrt. Pathol*., **83:** 51-59.

Undeen, A. H. (1975) Growth of *Nosema algerae* in Pig Kidney cell culture. *J. Protozool*. **22:** 107-110.

Undeen, A. H.; Johson, M.A. and Becnell, J.J. (1993) The effect of temperature on the survival of *Edhazardia aedis* (Microspora: Amblyosoridae), a pathogen of *Aedes aegypti*. *J. Invertebrt. Pathol*., **67:** 80-91.

Undeen, A.H. (1990) A proposed mechanism for the germination of microsoridian (Protozoa : Microspora) spores . *J. Theoret. Biol*., **142:** 223- 235.

Undeen, A.H. and Epsky, N. D. (1990) *Invitro* germination and *invivo* germination of *Nosema locustae* (Microsporidia : Nosematidae) spores. J. *Invertebr. Pathol*. **56:** 371. 379.

Vavra, J. (1976) Sytructure of the microsporidia. In "Comparative Pathobiology. Biology of the Microsporidia." (L.A.Bulla, Jr. and T.C Cheng. Eds,) Vol. 1. pp. 1-85. Plenum Press. New York.

Venkata Reddy.; Dandin, S.B.; Baig, M.; Sengupta K.; Giridhar, K. and Singal, B. K. (1990) Efficacy of Asiphor as a disinfectant against incidance of silkworm, *Bombyx mori* L. *Indian J. Seric*., **29:** 147-148.

Virender Kumar.; Nataraju, B.; Thiagarjan, V and Datta, R. K. (2002) Application of systemic fungicides for the control of white muscardine in silkworm, *Bombyx mori* L., *Int. J. Indust. Entomol.*, **5:** 171-174.

Vossbrinck, C.; Baker, M.D.; Didier, E.S.; Debrunner-Vossbrinck, B.A. and Shadduck, J.A. (1993) Ribosomal DNA sequences of *Encephalitozoon hellem* and *Encephalitozoon cuniculi*: Sepcies identification and phylogenetic constriction. *J. Eukaryot. Microbiol.*, **40:** 354-362.

Vossbrinck, C.R.; Maddox, J.V.; Friendman, S.; Debrunner-Vossbrinck, B.A. and Woese. C, R. (1987) Ribosomal RNA sequence suggests microsporidia are extremely ancient eukaryotes. *Nature*., 326: 411-414.

Wang, C.Y.; Solter, L.F.; T'sui, W.H. and Wang, C. H. (2005) An Endoreticulatus species from Ocinara lida (Lepidoptera: Bombycidae) In Taiwan. *J. Inverttebr. Pathol.*, **89:** 123-135.

Weidner, F. (1972) Ultrastuructural study of microsporidian invasion into cells. *Z. parasitenk*., **40:** 227-242.

Weiser, J. (1961) die Mikrosporidien als Parasiten der Insekten, *Monogr. Agew. Ent*., **17:** 1-149.

Wieser, J. (1969) Immunity of Insects to protozoa, In "*Immunity to parasitic animals"* (G. J. Jackson, R. Herman and I. Singer, eds.) Appleton century crafts, New York. P. 129-147.

Wilson, G.G. (1979) Reduced spore production of *Nosema fumiferance* (Microsporidia) in spruce budworm (Choristoneura fumiferana) reared at the elevated temperature. *Can. J. Zool*. **57:** 1167-1168.

Wilson, G. G. and Sohi, S. S. (1977) Effect of temperature on healthy microsporidia infected continuous cultures of *Malacosoma disstria* hemocytes. *Can. J. Zool.* 55: 713-717.

Wilson, G.G. (1974) The effect of temperature and ultraviolet radiation on the infection of *Choristoneura fumiferana* and *Malacosoma pluviale* by a microsporidian parasite, *Nosema* (Perezia) *fumiferana* (Thom) *Canad. J. Zool*., **52:** 59-63.

Yaman, M. and Radek, R. (2003) *Nosema chaetocnemae* sp. n. (Microsporo : Nosematidae), Microsporidian parasite of *Chaetocnema tibialis* (Coleoptera : Chrysonelidae). *Acta protozoologica.*, **42:** 231-237.

Yaman, M.; Radek, R.; Aslan, I. and Erturk. (2005) Characteristic features of Nosema phyllotretae Weiser 1961, a microsporidian parasite of Phyllotreta atra (Coleoptera: Chrysomelidae) in Turkey. Zoological studies 44: 368-372.