

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE NO.
1.1	Adult female worm of <i>Wuchereria bancrofti</i>	11
1.2	Microfilaria of <i>Brugia malayi</i>	11
1.3	Microfilarial sheath as observed by Giemsa stain (A) and phase contrast microscopy (B)	11
1.4	Comparative analysis of the life cycles of three human parasites reveal a similarity in developmental patterns. The life cycle of filarial parasite alternates between two hosts - <i>Brugia malayi</i> infective larvae are transmitted to the human host by arthropod vector, <i>C. quinquefasciatus</i>	12
1.5	Cross-section of a lymph node infested with Bancroftian parasites	17
1.6	Granulomatous lesion surrounding filarial parasitic debris (white ring)	17
1.7	Dense eosinophilic reaction around <i>W. bancrofti microfilaria</i> (Giemsa nuclear stain)	17
1.8	Illustration of the morphological features of necrosis and apoptosis	20
1.9	Pathways of apoptotic induction are closely associated with survival signals	23
1.10	Two models are proposed to explain <i>c-myc's</i> role in induction of apoptosis - the "conflict" model and the "dual signal model"	32
1.11A	Bcl-2 Family of Proteins	35
1.11B	Bcl-2 is an inner mitochondrial membrane channel regulating proton flux	35

FIGURE NO.	TITLE	PAGE NO.
1.12A	The Ras-Raf-1-MAPkinase - cascade transmit proliferative signals to the nucleus leading to activation of Myc. Activated PKC may exploit this cascade to trigger Myc-mediated apoptosis, while downregulating <i>bcl2</i>	39
1.12B	Synergistic cooperation between Ras and Bcl-2: Activation of Raf-1 by Ras enables its translocation to the mitochondrial membrane, bringing about phosphorylation of Bcl-2	42
3.1A	A confluent monolayer of HEp-2 cells at 6 days in culture (untreated)	88
3.1B	HEp2 cells exhibit apoptotic morphology on 6 days of treatment with filarial parasitic sheath proteins	88
3.2	Dose response analysis of HEp2 cells on exposure to filarial parasitic sheath proteins	90
3.3	Time course analysis of HEp2 cells on exposure to filarial parasitic sheath proteins	90
3.4A	PSSFV <i>neobcl2</i>	92
3.4B	Western Blot to demonstrate overexpression of Bcl-2 in transfected clones (lanes 2-5) in comparison to untransfected cells (lane 1)	92
3.5A	HEp-2 <i>bcl2</i> cells in culture for 6 days (untreated) show similar morphology and growth characteristics as wildtype cells	94
3.5B	HEp-2 <i>bcl2</i> cells are resistant to apoptosis induced on 6 days of treatment with filarial parasitic sheath proteins	94
3.6	Dose response analysis of HEp2 <i>bcl2</i> cells reveals abortive proliferation at 125µg (62.5 µg/ml), which was used as the dose for all future experiments	95
3.7	Time course analysis of HEp2 <i>bcl2</i> cells on exposure to filarial parasitic sheath proteins, show abortive proliferation at 5 and 6 days of treatment with filarial proteins	95

FIGURE NO.	TITLE	PAGE NO.
3.8	Immunostaining of actin filaments in HEp2 cells, control (A), and treated (B) show rearrangement of actin in apoptotic cells. HEp2 <i>bcl2</i> cells, both control (C) and treated (D), show no rearrangement	97
3.9	Immunoblot for actin indicates intact actin protein in HEp2 controls (Lane 1), treated cells (Lane 2), and in HEp2 <i>bcl2</i> cells, both control (lane 3) and treated (Lane 4). Lane M represents molecular weights in KDa	99
3.10	Immunostaining for tubulin provides evidence for cycling HEp2 cells in treated (B) population. HEp2 <i>bcl2</i> cells display abortive proliferation on treatment (D(i)) and (D(ii)). Untreated cells show random distribution of tubulin (A) and (C)	98
3.11	Immunoblot for tubulin indicates intact tubulin protein in HEp2 controls (Lane 1), treated cells (Lane 2), and in HEp2 <i>bcl2</i> cells, both control (lane 3) and treated (lane 4). Lane M represents molecular marker in KDa	99
3.12	Immunoblot for $\alpha 5\beta 1$ (Fibronectin Receptor) (8% Gel) indicates cleavage of receptor in HEp2 treated cells (Lane 2), absent in controls (Lane 1), and absent in HEp2 <i>bcl2</i> cells, both control (Lane 3) and treated (Lane 4). Lane M represents molecular weights in KDa	101
3.13	DNA laddering observed HEp2 cells on 6 days exposure (Lane 4) absent in untreated controls and 24 hours treated cells (Lane 2 and 3). HEp2 <i>bcl2</i> cells do not show laddering in control (Lane 6) or 24 hours or 6 days treatment (Lanes 7 and 8 respectively). Actinomycin D was used as a positive control shows laddering in HEp2 in HEp2 cells (Lane 5) but not in HEp2 <i>bcl2</i> cells (Lane 9) Lane M represents 1 Kb ladder.	101
3.14	Apoptosis in HEp2 cells observed on 6 days of exposure to filarial parasitic sheath proteins, absent in HEp2 <i>bcl2</i> cells (A-C) - Phase contrast micrographs of HEp2:untreated controls (A) represent cells maintained for 6 days in culture show no change in morphology whereas cells exposed to filarial parasitic sheath protein for 6 days (B) and Actinomycin D treated cells (C) show apoptotic morphology ; (D-F) - Visualisation of DNA strand breaks using TUNEL staining: Untreated control HEp2 cells (D) do not stain positive for the assay whereas DNA strand breaks observed on 6 days exposure (E) and in Actinomycin D treated cells (F) (G-I) - Propidium iodide staining shows condensed nuclei in HEp2 cells on 6 days treatment (H) and Actinomycin D treated cells (I), untreated cells show normal nuclear morphology (G) No such changes observed in HEp2 <i>bcl2</i> cells (J-L, M-O, P-R).	103

FIGURE NO.	TITLE	PAGE NO.
3.15	Flow cytometric analysis of HEp2 and HEp2 <i>bcl2</i> cells on exposure to filarial proteins for 24 hours, 2 days and 6 days HEp2 cells treated with filarial parasitic sheath proteins show sub-G1 peaks whereas HEp2 <i>bcl2</i> cells do not show any such peak. Untreated controls of both HEp2 and HEp2 <i>bcl2</i> cells taken at 6 days of culture show no apoptotic characteristics	105
3.16	(A) and (B) Comparison of [³ H] Thymidine incorporation with Northern Blot for <i>c-myc</i> shows elevated levels of <i>c-myc</i> at initial stimulation (24 hours) and at apoptotic window (6 days) in HEp2 cells. (B) HEp2 <i>bcl2</i> cells show a significant increase in proliferation and <i>c-myc</i> expression at 6 days on treatment with filarial proteins, supporting abortive proliferation	107
3.17	Northern blot for c-H-ras indicates elevation of expression in HEp2 <i>bcl2</i> cells on treatment with filarial proteins, both at early and late time points, indicating a cooperative effort by <i>bcl2</i> and Ras in the prevention of apoptosis	107
3.18	(A) PKC activation observed in HEP2 cells on 6 days treatment with filarial parasitic sheath proteins (B) PKC is inhibited at similar time points in HEP2 <i>bcl2</i> cells	110
3.19	Tyrosine kinase activity is downregulated in apoptosing HEp2 cells at 6 days (A) whereas in HEp2 <i>bcl2</i> cells (B) it increases to almost double that of untreated controls	110
3.20	Cytokine gene expression studies in HEp2 cells by Northern blot indicates elevation in levels of IL-8 and, to a lesser extent, IL-6, on treatment with filarial proteins. Similar expression levels are detected in <i>c-fos</i> , indicating a role for Fos as a transcriptional activator for cytokine gene expression	112
3.21	Cytokine ELISA for IL-6(A) and IL-8(B) indicate release of the cytokines into cell culture supernatant on treatment with filarial protein. LPS was used as a positive inducer of cytokine secretion in both cell lines	113
4.1	Hypothetical model to explain apoptotic signalling in HEp2 cells : PKC drives apoptotic cascade via ras effectors and NF-κB, which converge on Myc. Growth factor - mediated mitogenic signals "conflict" with apoptotic signals due a downregulation of Bcl-2 or loss of anchorage (?)	125

FIGURE NO.	TITLE	PAGE NO.
4.2	Hypothelial model to explain survival in HEp2 <i>bcl2</i> cells : Activated <i>ras</i> synergises with Bcl-2 energised by tyrosine phosphorylation (by integrin associated <i>c-abl</i> ?). PKC inactivation annulus apoptotic cascade	129
4.3	Protein kinase C plays an important role in governing apoptotic (brown shading) and proliferative (green shading) cascading via Ras family members	131
4.4	Tyrosine kinases suppress apoptosis via activation of Raf-1 activation followed by Bcl-2 upregulation Growth factor associated RTKs and integrin associated <i>c-abl</i> transmit survival signals to the nucleus	135