

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE NO.
1.1	Continuous absorption chromatography column	5
1.2	Gradient sublimation for the separation of impurities from organic material	6
1.3	Phase diagram showing impurity segregation behavior at low impurity concentration	7
1.4	Directional solidification process	8
1.5	Bridgman growth setup	10
1.6	Modified furnace	10
1.7	Ampoule designs for single crystal growth	11
1.8	Apparatus used for Czochralski growth of organic materials	13
1.9	Schematic diagram of apparatus used for growth from solution at low temperatures and undercooled melt	14
1.10	Ampoule designs used by different authors (Sherwood 1960)	22
1.11	Ampoule design used by Marciniak	23
1.12	Containers listed by Lipsett employed by various authors	24
1.13	Ampoule design used by Lipsett	24
1.14	Ampoule design for single crystal growth given by Brian Pamplin (1980)	25
1.15	Ampoule design for single crystal growth (Brice 1986)	26
1.16	Temperature profile for organic material crystal growth (Lipsett 1957)	27
1.17	Schematic diagram of Vapour Growth technique (Sloan and Vapour Transport technique)	30

FIGURE NO.	TITLE	PAGE NO.
1.18	Energy levels of an organic molecule with π -electron structure (Birks 1964)	36
1.19	Scintillation fluorescence and quenching	37
1.20	Energy resolution setup	44
1.21	Time resolution setup	44
1.22	Block diagram of PSD circuit (Sperr 1974)	46
1.23	Schematic showing the thermalisation of positrons in solids	52
2.1	Schematic diagram of Vertical Bridgman setup	61
2.2	Photograph of growth setup	62
2.3a	Temperature profile of 30 cm length furnace for the growth of stilbene crystal	63
2.3b	Temperature profile of 60 cm length furnace for the growth of stilbene crystal	63
2.4a	Schematic of furnace heating element used for winding	65
2.4b	Special translation assembly used for zone refining	65
2.4c	Zone refined material with tube	66
2.5	Fluorescence spectra of <i>trans</i> -stilbene (Commercial)	68
2.6	Fluorescence spectra zone purified <i>trans</i> -stilbene	69
2.7	X-ray spectrum of 10 times zone passed stilbene	70
2.8	X-ray spectrum of 30 times zone passed stilbene	70
2.9	Loss of single crystallinity due to constant growth rate 2 mm/hr	71
2.10	Crystal grown with various ampoule design	73
2.10	Crystals grown with growth varying from 2 to 0.5 mm /hr	73
2.12a	Strongest diffraction peak	75
2.12b	Powder X-ray diffraction peak showing low intensity peaks	75
2.13	FTIR spectrum of grown ingot	76

FIGURE NO.	TITLE	PAGE NO.
2.14	UV – Visible transmission spectrum of 8 mm thick stilbene crystal	76
2.15	Time resolution spectrum of stilbene	77
2.16	Temperature profile of 30 cm length furnace for the growth of anthracene crystal	79
2.17	Fluorescence spectra of anthracene	81
2.18	Fluorescence spectra of commercial anthracene	84
2.19	Fluorescence spectra of DMSO Purified anthracene	84
2.20	Fluorescence spectra of DMF purified anthracene	85
2.21	Fluorescence spectra of sublimation purified anthracene	85
2.22	Fluorescence spectra of Zone purified anthracene	86
2.23	X-ray spectrum of DMSO purified anthracene	87
2.24	X-ray spectrum of DMSO, DMF, Hexane purified anthracene	87
2.25	X-ray spectrum of zone refined anthracene	88
2.26	X-ray spectrum of sublimation purified anthracene	88
2.27	Grown anthracene crystals with modified ampoule design	90
2.28	Grown stilbene crystal with modified ampoule design	90
2.29	Transmission spectrum of double walled ampoule (with modification) grown anthracene crystal	93
2.30	Transmission spectrum of double walled ampoule with modification) grown stilbene crystal	94
2.31	NMR Spectra of anthracene	95
2.32	NMR Spectra of stilbene	95
2.33	Decay time measurement setup	97
2.34	Grown stilbene crystals	98
2.35	Grown anthracene crystals	99
3.1a	Illustrative figure of ampoule with selective self-seeding technique	104

FIGURE NO.	TITLE	PAGE NO.
3.1b	Photograph of SSVBT grown crystals	105
3.2	Photograph of the multicrystals obtained from the ampoule with 3mm gap with the inner capillary edge and outer tube edge	106
3.3	Necked-capillary ampoule grown crystals used for the optical observation	107
3.4	X-ray rocking curve for the selective self seeded <i>trans</i> -stilbene crystal	108
3.5	X-ray rocking curve for the conventional vertical Bridgman grown <i>trans</i> -stilbene crystal	108
3.6	Photograph of the DRSSVBT-grown anthracene crystal	110
3.7	IR spectrum of anthracene	111
3.8	X-ray rocking curve for the conventional method grown crystal	113
3.9	X-ray rocking curve for the DRSSVBT-grown crystal	114
3.10	Timing resolution for ^{22}Na 511 KeV gamma ray	114
3.11a,b	Single wall ampoule design and grown crystal	117
3.12.a,b	Double wall ampoule design and grown crystal	118
3.13a,b	Conical bottom and cylindrical top design and grown crystal	118
3.14a,b	Full conical selective self-seeded ampoule design and grown crystal	119
3.15	X-ray rocking curve of stilbene crystal grown by using single wall ampoule	120
3.16	X-ray rocking curve of stilbene crystal grown by using double wall ampoule without selective self-seeding	121
3.17	X-ray rocking curve of stilbene crystal grown with the gradual decrease of growth rate	121
3.18a,b	Optical micrograph of selective self seeded ampoule grown crystal, surface etched using CCl_4	122

FIGURE NO.	TITLE	PAGE NO.
3.19	X-ray rocking curve of crystal in conical region of the ampoule	123
3.20	X-ray rocking curve of the crystal in the cylindrical region of the ampoule	124
3.21	X-ray rocking curve of the crystal ingot grown from low cone angle ampoules	124
3.22	X-ray rocking curve of the crystal grown from high cone angle ampoules	125
3.23	X-ray rocking curve of the crystal near the wall region	125
3.24	Optical micrograph of crystals grown without selective self seeding	126
3.25	X-ray rocking curve of the crystals grown with high and low twist angles of the capillary	127
3.26a,b	Optical micrograph of the crystals grown with high and low twist angles of the capillary	128
3.27	X-ray rocking curve of fast grown crystals without selective self seeding	129
3.28	X-ray rocking curve of fast grown crystals using double walled ampoule without selective self seeding	130
3.29	Optical micrograph of etched crystal indicating low angle grain boundary	130
3.30	X-ray rocking curve of slow seeded crystal grown with conical ampoule	132
3.31	X-ray rocking curve of crystals grown with slow seeding using full conical ampoule (Top region)	132
4.1	Raman Phonon modes spectrum of irradiated and unirradiated stilbene crystals	138
4.2	Raman molecular mode spectrum of irradiated and unirradiated stilbene crystal	139
4.3	Raman spectrum of stilbene powder	148
4.4	Photoluminescence setup	151

FIGURE NO.	TITLE	PAGE NO.
4.5	Photoluminescence spectra of stilbene (on the bubble spot)	151
4.6	Photoluminescence spectra of stilbene (Edge of the bubble)	152
4.7	Photoluminescence spectra of stilbene (1 mm away from bubble spot)	152
4.8	Photoluminescence spectra of stilbene (Bubble free crystal)	153
4.9	Photoluminescence spectra of stilbene (multi bubble spot)	153
4.10	Photoluminescence spectrum of DRSSVBT grown anthracene crystal	157
4.11	Photoluminescence spectrum of conventional grown anthracene crystal	157
4.12	Photoluminescence spectrum of vapour grown anthracene crystal	158
4.13	Photoluminescence spectrum of solution grown anthracene crystal	158
4.14	Raman spectrum of solution grown anthracene crystal	159
4.15	Raman spectrum of vapour grown anthracene crystal	159
4.16	Raman spectrum of conventional method grown anthracene crystal	160
4.17	Raman spectrum of DRSSVBT grown anthracene crystal	160
4.18	Microphotograph of etched (001) surface with indentation	163
4.19	Hardness Vs. load plots for (001) surface	163
4.19	Microphotograph of etched (110) surface with indentation	163
4.21	Hardness Vs. load for (110) surface	164

FIGURE NO.	TITLE	PAGE NO.
4.22	Microphotograph of etched (100) surface with indentation	164
4.23	Hardness Vs. load for (100) surface	164
4.24	Microhardness plot of anthracene crystals grown by conventional method and DRSSVBT	167
5.1	Rise time shift with proton energy in stilbene scintillator (Theoretical curve) (Papadopoulos 1997)	179
5.2	Scintillation response curve of stilbene for protons (Papadopoulos 1999)	180
5.3	Crystal elements used in the experiment	182
5.4	14 MV UD Tandem Pelletron (Nuclear Science Center, New Delhi)	185
5.5	Block diagram of PSD	186
5.6	Time relationship between the pulses at various points of the PSD circuit a) Normal operation b) During adjustment	187
5.7	Experimental setup used in the experiment	188
5.8	Timing calibration using 25 MeV proton + ^{60}Co gamma source	189
5.9	Timing calibration using proton beam + ^{60}Co gamma source	190
5.10	Energy calibration using 25 MeV proton beam	190
5.11	Block diagram of SSBD for energy calibration spectrum of Protons	191
5.12	Energy calibration using SSBD in steps of 5 MeV up to 25 MeV	192
5.13	Response curve of SSBD to Protons	192
5.14	Two-dimensional energy Vs TAC spectrum of 5 MeV proton + ^{60}Co gamma(-1600 V PMT bias)	194
5.15	Two-dimensional energy Vs TAC spectrum of 10 MeV proton + ^{60}Co gamma(-1600 V PMT bias)	194

FIGURE NO.	TITLE	PAGE NO.
5.16	Two-dimensional energy Vs TAC spectrum of 15 MeV proton + ^{60}Co gamma(-1600 V PMT bias)	195
5.17	Two-dimensional energy Vs TAC spectrum of 20 MeV proton + ^{60}Co gamma(-1600 V PMT bias)	195
5.18	Two-dimensional energy Vs TAC spectrum of 25 MeV proton + ^{60}Co gamma(-1600 V PMT bias)	196
5.19	Intensity Vs energy spectrum (-1600 V PMT bias)	196
5.20	TAC spectrum showing P- γ discrimination (-1600 V PMT bias)	197
5.21	Two-dimensional energy Vs TAC spectrum of 5 MeV proton + ^{60}Co gamma (-1800 V PMT bias)	197
5.22	Two-dimensional energy Vs TAC spectrum of 10 MeV proton + ^{60}Co gamma(-1800 V PMT bias)	198
5.23	Two-dimensional energy Vs TAC spectrum of 15 MeV proton + ^{60}Co gamma(-1800 V PMT bias)	198
5.24	Two-dimensional energy Vs TAC spectrum of 20 MeV proton + ^{60}Co gamma(-1800 V PMT bias)	199
5.25	Two-dimensional energy Vs TAC spectrum of 25 MeV proton + ^{60}Co gamma(-1800 V PMT bias)	199
5.26	Intensity vs Energy spectrum (-1800 V PMT bias)	200
5.27	TAC spectrum showing P- γ discrimination (-1800 V PMT bias voltage)	200
5.28	Energy response of stilbene (-1600 V PMT bias)	203
5.29	Energy response of stilbene (-1800 V PMT bias)	204
5.30	Shift in rise time for protons with energy (-1600 V PMT bias)	204
5.31	Shift in rise time for protons with energy (-1800 V PMT)	205
5.32	Two-dimensional energy Vs TAC spectrum of C(P,n γ) reaction (-1600V PMT bias)	208

FIGURE NO.	TITLE	PAGE NO.
5.33	TAC spectrum showing C(P, γ) reaction (-1600 PMT bias voltage)	208
5.34	Energy response of stilbene for C(P, γ) reaction (-1600 V PMT bias)	209
5.34	Energy resolution of stilbene for ^{60}Co and ^{137}Cs source	213
5.36	Plot of T_{max} Vs channels	214
5.37	Time resolution spectrum of BGO-stilbene using ^{22}Na gamma source	214
5.38	Timing resolution spectrum for Na^{22} 511 keV gamma ray using conventional method grown crystal	219
5.39	Timing resolution spectrum for ^{22}Na 511 keV gamma ray using selective self seeded method grown crystal	219
5.40	n- γ pulse shape discrimination spectra of ^{252}Cf source	220
5.41	α - γ pulse shape discrimination spectrum of ^{241}Am source	220
5.42	Time resolution setup	224
5.43	Time resolution spectrum of the single walled ampoule grown crystal	224
5.44	Optical photomicrograph showing hollow pore with striations of single walled ampoule grown crystal	226
5.45	Time resolution spectrum of the double walled ampoule grown crystal	225
5.46	SEM micrograph of double walled ampoule grown crystal	226
5.47	Time resolution spectrum of SSVBT grown crystal	225
5.48	SEM micrograph of the SSVBT grown crystal	226
6.1	Photographic view of the 2D-ACAR system at University of Madras, Chennai	236

FIGURE NO.	TITLE	PAGE NO.
6.2	Coincident 511 keV energy spectra sampled simultaneously from the energy signals using <i>trans</i> -stilbene single crystal	236
6.3	Photograph of the low temperature setup used for present study at University of Madras, Chennai	238
6.4a	Perspective surface plot of 2D-ACAR spectrum $N(p_x, p_y)$ for <i>trans</i> -stilbene projected along [010] direction such that p_x is along [001] and p_y along [100] direction at RT	241
6.4b	Contour plot of 2D-ACAR spectrum $N(p_x, p_y)$ for <i>trans</i> -stilbene projected along [010] direction such that p_x is along [001] and p_y along [100] direction at RT. The contour spacing is 1/16 of the maximum	241
6.5	Cross-section of 2D-ACAR spectrum $N(p_x, p_y)$ –RT	242
6.6a	Perspective surface plot of 2D-ACAR spectrum $N(p_x, p_y)$ for <i>trans</i> -stilbene (measured at 210 K) projected along [010] such that p_x is along [001] and p_y along [100]	243
6.6b	Contour plot of 2D-ACAR spectrum $N(p_x, p_y)$ of <i>trans</i> -stilbene (measured at 210 K) projected to [010] such that p_x is along [001] and p_y along [100]	243
6.7a	Perspective surface plot of 2D-ACAR spectrum $N(p_x, p_y)$ for <i>trans</i> -stilbene (measured at 110 K) projected along [010] such that p_x is along [001] and p_y along [100]	244
6.7b	Contour plot of 2D-ACAR spectrum $N(p_x, p_y)$ of <i>trans</i> -stilbene (measured at 110 K) projected to [010] such that p_x is along [001] and p_y along [100]	244
6.8	Cross-section of 2D-ACAR spectrum $N(p_x, p_y)$ (RT, 210 K and 110 K)	246