

LIST OF FIGURES

Figure No.	Description
3.1	Map of intermediate vector pGEM-T Easy
3.2	Map of expression vector pQE-30
4.1	PCR amplification of <i>asd</i> gene using primers F1 and R1
4.2	Map of the cloning vector pGEM-T Easy with <i>M. tb asd</i> insert
4.3	PCR amplification of <i>asd</i> gene using primers F2 and R2
4.4	Diagrammatic representation of cloning of <i>asd</i> gene in intermediate vector pGEM-T Easy
4.5	Double digestion of pGEM-T Easy and pQE-30 vectors
4.6	Plasmid map of the construct pSST1
4.7	Diagrammatic representation of subcloning of <i>asd</i> gene into pQE-30, induction of its expression and DNA sequencing
4.8 (a)	Results of the forward sequencing of the cloned <i>asd</i> gene
4.8 (b)	Results of the reverse sequencing of the cloned <i>asd</i> gene
4.9	BLAST-2 of sequencing result against sequence of <i>asd</i> gene from <i>M. tb</i> genome sequence present in NCBI database
4.10	Submission of <i>M. tb asd</i> gene sequence to GenBank database
4.11	SDS-PAGE of induction of expression of recombinant <i>M. tb</i> ASD in <i>E. coli</i>
4.12 (a)	SDS-PAGE of the induced His-tagged ASD used for transfer to nitrocellulose membrane
4.12 (b)	Western blot of His-tagged <i>M. tb</i> ASD using primary anti-His antibodies
4.13	SDS-PAGE of purification of recombinant <i>M. tb</i> ASD using one step Ni-NTA affinity chromatography
4.14	Double reciprocal (Lineweaver-Burk) plots of the three substrates ASA, Pi and NADP
4.15	ClustalX multiple alignment of amino acid sequences of ASD of <i>M. tb</i> and seven other bacteria
4.16	Theoretical pI and molecular weight of native <i>M. tb</i> ASD calculated by ExPASy pI/Mw tool
5.1	DAP biosynthetic pathway present in bacteria