

## REFERENCE

- Alexandrov, E.A., 1973. The Precambrian banded iron-formation of the Soviet Union. *Econ.Geol.* 68, pp.1035-1062.
- Anantha Iyer, G.V. and Vasudev, V.N., 1979. Geochemistry of Archaean metavolcanic rocks of Kolar and Hutti Gold Fields, Karnataka, India. *Jour.Geol.Soc. India*, V.20, pp.419-432.
- Appel, P.W.U., 1974. On an unmetamorphosed iron-formation in the early Precambrian of Southwest Greenland. *Miner.Deposita*, V.9, pp.75-82.
- Appel, P.W.U., 1983, Rare earth elements in the early Archaean iron-formation, West Greenland. *Precamb-Research* V 20, pp.243-258.
- Appel, P.W.U., 1987. Geochemistry of the early Archaean Isua iron-formation, West Greenland. In: Appel, P.W.U. and Gene L.Laberge (Edts). *Theopraster Publ. S.A. Greece*. pp.31-68.
- Appel, P.W.U. and Mahabaleswar, B., 1988. Secular trends in rare earth element patterns of Precambrian iron-formation from India and Greenland. *Jour.Geol.Soc.India*, Vol.32, No.3, pp.214-226.
- Balaji Rao, B., 1913. Report on prospecting work to the east and south of Chikmagalur. *Rec.Mysore Geol.Dept.V.13*, pp.139-149.
- Bayley, R.W. and James, H.L., 1973. Precambrian iron-formation of the United State. *Econ.Geol.*, V,68. pp 934-959
- Beckinsale, R.D., Reeves Smith, G., Gale, N.H., Holt, R.W. and Thompson, B., 1982. Rb-Sr and Pb-Pb whole rock isochron ages and REE data for Archaean gneisses and granites, Karnataka State, South India. Abstract, Indo-US Workshop on Precambrian of South India, Hyderabad.
- Beckinsale, R.D., Drury, S.A. and Holt, R.W., 1980. 3360 M.y. old Gneisses from the Indian Craton. *Nature*. V. 283, PP. 469-470.
- Berge, J.W 1977. Geology and origin of the hematite ores of the Nimba Range, Liberia. *Eco. Geo*, V. 72, pp.582-607.
- Beukes, N.J., 1973. Precambrian iron-formations of South Africa. *Econ.Geol.*68: 960-1004.

- Beukes, N.J., 1980. Suggestions towards a classification and nomenclature for iron-formation. *Trans.Geol.Soc. S.Africa*, V.83, pp.285-290.
- Bhaskara Rao, Y.J. and Naqvi, S.M., 1978. Geochemistry of metavolcanics from the Bababudan schist belt: a late Archaean/early proterozoic volcano sedimentary pile from India. In B.F.Windley and S.M.Naqvi (Ed.), *Archaean Geochemistry*, Elsevier, Amsterdam, pp.325-341.
- Bhaskara Rao, Y.J., and Drury, S.A. 1982. Incompatible trace element geochemistry of Archaean metavolcanic rocks from the Bababudan volcanic-sedimentary belt, Karnataka. *Jour.Geol.Soc. India*, V.23, pp.1-12.
- Bhaskara Rao, Y.J., Beck, W., Murthy, V.R., Charan, S.N. and Naqvi, S.M. 1983. Geology, geochemistry and age of metamorphism of Archaean grey gneisses around Channarayaptna, Hassan, Dist, Karnataka, South India. In *Precambrian of South India*. S.M.Naqvi and J.J.W.Rogers (Eds). *Geol. Soc. India. Mem. 4*. pp.309-328.
- Borchert, H., 1965. Formation of marine sedimentary iron ores. In J.P.Riley & G.Skirrow (Eds.), *Chemical oceanography*, Vol.2. Academic Press, New York, pp.159-204.
- Breitkopf, J.H., 1988. Iron formation related to mafic volcanism and ensialic rifting in the southern margin zone of the Damara orogen. Namibia. *Precamb. Res.* 38.111-130.
- Bridgwater, D., Watson, J. and Windley, B.F., 1973. The Archaean craton of the North Atlantic region. *Phil.Trans.R.Soc.Lond.* V.A273, pp.493-512.
- Bruce Foote, 1888. 'The Dharwar System', the chief auriferous rocks in South India. *Rec.Geol.Surv.India*, V.21 (2), pp.1-40.
- Bruce Foote, 1889. Dharwar System, the chief auriferous series in South India. *Rec.Geol.Surv.India.*, V.22 (1), pp.17-39.
- Bruce Foote, 1900. Geological notes an traversed through the Mysore State. *Mem.Mys. Geol.Dept.*, V.1, pp.103.
- Chadwick, B., Ramakrishnan, M. and Viswanatha, M.N., 1981b. Structural and metamorphic relationship between Sargur and Dharwar supracrustal rocks and Peninsular gneiss in Central Karnataka. *JOUR.Geol.Soc. India.*, V.22, pp.557-569.
- Chadwick, B., Ramakrishnan, M. and Viswanath, M.N., 1985.

- Bababudan Schist Belt. Geol.Soc.Ind., V.26, pp.769-821.
- Chadwick, B., Garrioch, N.H.G, Ramakrishnan, M., and Viswanath, M.N., 1986. Mineral composition, texture and deformation in late Archaean banded iron-formation rich in magnesio riebeckite and aegirine, Bababudan, Karnataka, Southern India. Geol.Soc.Ind. Vol.28, pp.189-200.
- Chakrapani Naidu, M.G. and Rama Rao, G.V.S., 1968. Mineralogy of the banded ferruginous quartzite, Bababudan Hills, Mysore State. India Mineralogist, Vol.9, pp 59.67.
- Chakraborty, K.L. and Majumder, T., 1986. Geological aspects of the banded iron-formation of Bihar and Orissa. Geol.Soc.India, Vol.28, pp 109-133.
- Davy, R., 1983. Chemical composition of BIF. In : A.F.Trendall and R.C.Morris (Eds) Iron-formation: Facts and Problems. Elsevier, Amsterdam, pp.325-343.
- Deer, W.A., Howie, R.A. and Zussman, J., 1969. An introduction to rock forming minerals. Longmans, London.
- Devaraju, T.C. Anantha Murthy, K.S. and Khandali, S.D., 1986. Iron-formation of Chiknayakanhalli, Greenstone belt, Karnataka. Geol.Soc.India , Vol.28, pp.201-217.
- Dimroth, E., 1975. Paleo-environment of iron-rich sedimentary rocks. Geol.Rundschau, V.64, pp.751-767.
- Dimroth, E. and Chauvel, J.J., 1973. Petrography of the Sokomon iron-formation in the part of the Central Labrador Trough, Quebec, Canada. Geol.Soc.of America. Bull. V.84, pp.111-134.
- Dorr, J.V.N., 1973. Iron-formation and associated manganese in Brazil. In: Genesis of Precambrian iron and manganese deposits. Paris, UNESCO. Earth.Sci., v.9, pp. 106-113.
- Drury, S.A., Holt, R.W., Van Clasteren, P.C. and Beckinsale, R.D., 1983. Sm-Nd and Rb-Sr ages for Archaean rocks in W. Karnataka. Jour.Geol.Soc.India, Vol.24, pp 454-459.
- Dunn, J.A., 1935. The origin of iron ores in Singhbhum, India. Eco.Geol.V.30, pp.643-654.
- Dymek, R.F., and Klein, C., 1988. Chemistry, petrology and origin of banded iron formation lithologies from the 3800 M.A. Isua supracrustal belt, West Greenland. Precamb. Res. 39, pp 247-302
- Edwards, A.B., 1965. Textures of ore minerals and their

significance (2nd edition). Australian institute of mining and metallurgy.

- Eichler, J., 1976. Origin of the Precambrian banded iron-formation. Handbook of Stratiform and stratabound ore deposits. Elsevier, Amsterdam, V.7, p.201.
- Ernst, W.G., 1968. Amphiboles. Crystal chemistry, phase relations and occurrence. Springer-Verlag, Inc., New York, p125.
- Eskola., 1920. The mineral facies of rocks. Nork Geol. tidsskr., v.6, pp 143- 194.
- ESkola., 1939. die entstehung der Gesteine(Barth, Correns, Eskola). Springer- verlag, Berlin.
- Fermor, L.L., 1909. Manganese ore deposits of India. Mem.Geol.Surv. India.
- Fermor, L L., 1936. An attempt at the correlation of ancient schistose formations of Peninsular India. Mem.Geol.Surv. India., V.70 (1) and 70 (2) pp.1-128.
- Floran, R.J. and Papike, J.J., 1975. Petrology of Gunflint iron-formation, Ontario-Minnesota: Pt.I. The lowgrade rocks. Bull.Geol.Soc.Amer., Vol.86, pp. 1169-1190.
- Floran,R.J. and Papike, J.J., 1978. Mineralogy and petrology of the Gunflint iron-formation, Minnesota-Ontario: Correlation of compositional and asemblage variations at low to moderate grade. Jour.Petrol. Vol.19, pp. 215-288.
- French, B.M., 1968. Progressive contact metamorphism of the Biwabik Iron-formation, Mesabi Range, Minnesota. Minn.Geol.Surv. Bull., Vol.45, pp.103.
- French, B.M., 1975. Mineral assemblages in diagenetic and low grade metamorphic iron-formation. Econ.Geol. Vol.68, pp.1063-1074.
- Friend, C.R.L., 1984. The origin of Closepet granites and the implications for the crustal evolution of Southern Karnataka. JOur.Geol.Soc.India. Vol.25, pp.73-84.
- Frietsch, R., 1970. Trace elements in magnetite and hematite, Sverges. Geol.Unders, Arshok, Vol.64, pp.1-136.
- Fryer, B.J., 1977a. Rare earth evidence in iron-formation for changing Precambrian oxidation states. Geochim.Cosmochim.Acta, Vol.41, pp.361-367.
- Fryer,B.J., 1977b. Trace element geochemistry of the Sokoman

- iron-formation. *Can.Jour. Earth Sci.*, Vol.14, pp.1598-1600.
- Fryer, B.J., 1983. Rare earth elements in iron-formation. In: A.F. Trendall and R.C. Morris (Eds). *Iron-formation: Facts and Problems*. Elsevier, pp.345-358.
- Gavisetty, K., 1975. Asbestos deposits of Manikyadara area, Bababudan hills, Chikmagalur district, Karnataka State. *Geological studies, Dept. of Mines and Geology.*, No.96, pp. 1-9.
- Gole, M.J. 1980a. Mineralogy and petrology of very low metamorphic grade Archaean banded iron-formations, Weld Range, Western Australia. *Am.Mineral.*, Vol.65. pp.8-25.
- Gole, M.J., 1981. Archaean banded iron-formation, Yilgarn Block, W.Australia. *Econ.Geol.*, Vol.76, pp.1954-1975.
- Gole, M.J. and Klein, C., 1981. Banded iron-formation through much of Precambrian time. *Jour. Geol.*, Vol.89. pp.169-183.
- Gole, M.J. and Klein, C., 1981b. Archaean banded iron-formation, Yilgarn block, Western Australia. *Econ.Geol.*, Vol.76. pp.1954-1974.
- Goldschmidt, V.M., 1958. *Geochemistry*. Oxford Uni.Press. Amer.House, London, pp.1-730.
- Goodwin, A.M., 1962. Stratigraphy and origin of iron formation, Michipicoten area, Algoma district. Ontario, Canada. *Geol.Soc.Am. Bull.* 73, pp. 561- 586.
- Goodwin, A.M., 1973. Archaean iron-formation and tectonic basin of the Canadian shield. *Econ.Geol.*, Vol.68, pp.915-933.
- Govett, G.J.S., 1966. Origin of banded iron-formation. *Geol.Soc.of America. Bull.* Vol.77. pp.1181-1212.
- Graf, J.L., 1977. Rare earth elements as hydrothermal tracers during the formation of sulfide deposits in volcanic rocks. *Econ.Geol.*, Vol.72, pp.527-548.
- Graf, J.L., 1978. Rare earth elements, iron-formation and sea water. *Geochim.Cosmochim.Acta.*, Vol.42, pp.1845-1850.
- Grigoryev, V.M., 1971. Quoted by Melnik: In: *Precambrian banded iron formations*. Elsevier. Publ.
- Gross, G.A., 1965,. The geology of the iron deposits in Canada: 1. General Geology and evolution of iron deposits. *Geol.Surv. Canada, Econ.Geol.Ser.Rep.*, No.22(1), pp.181.

- Gross, G.A., 1973. The depositional environment of the principal types of Precambrian iron formation. In: Genesis of Precambrian iron and manganese deposits. UNESCO, Paris. Earth Sciences, v.9, pp.15-21.
- Gross, G.A., 1980. A classification of iron-formation based on depositional environment. Canadian Mineral., Vol.18, pp.215-222.
- Gross, G.A., 1986. The metallogenetic significance of iron-formation and related stratafer rocks. Jour.Geol.Soc. of India., Vol.28, pp.92-108.
- Gundersen, J.N., and Schwartz, G.M., 1962. The geology of the metamorphosed Biwabik iron formation, Eastern Mesabi district, Minnesota, Univ. Minn, Minn. Geol. Surv. Bull. 43
- Haase, C.S., 1979b., Multisystems analysis of phase relations in the system  $\text{CaO}-(\text{Fe}, \text{Mg})\text{O}-\text{Al}_2\text{O}_3-\text{Fe}_2\text{O}_3-\text{SiO}_2-\text{CO}_2-\text{H}_2\text{O}$ : a petrographic grid for metamorphosed iron-formation. Geol.Soc.Am.Abstr. Programmes, Vol.11, pp.436.
- Haase, C.S., 1982a. Metamorphic petrographic petrology of the Negaunee iron-formation, Marquette district, northern Michigan: Mineralogy, metamorphic reactions and phase equilibria. Econ.Geol. Vol.77, pp.60-81.
- Han, T.M., 1971. Diagenetic-metamorphic replacement features in the Negaunee iron-formation of the Marquette iron range, Lake Superior district. Soc.Min.Geol.Jph. Spec. Iss. Vol.3, pp.430-438.
- Harrison, C.G.D., and Peterson, M.N.A., 1965. A magnetic mineral from the Indian ocean, Am. Mineral. 50: pp704-712.
- Hegemann, F., and Albrecht, F., 1954. Zur geochemie oxydischer eisenerze, chemie d. Erde, Vol.17, pp.81-103.
- Hietanen, A., 1938. On the petrology of Finnish Quartzites. Bull.Comm. Geol. Finland. Vol.122, pp.118.
- Holland, T.H., 1902. The mica deposits of India. Mem.Geol.Surv. India. Vol.34, pp.65-89.
- Hussain, S.M., Naqvi, S.M. and Gnaneshvara Rao, T., 1982. Geochemistry of mafic-ultramafic rocks from southern part of the Holenarsipur schist belt, Karnataka, India. Jour.Geol.Soc. India, Vol.23, pp.19-31.
- Iyengar, B.R.C., 1971. Cycles of Dharwar sedimentation. Seminar on Precambrian Geology of Peninsular shield,

Calcutta, pp.5-6 (Abst.).

- Jacobsen, S.B and Pimentel-Klose, M.R., 1988. A Nd isotopic study of the Hamersley and Michipicoten banded iron formations: the source of REE and Fe in Archen oceans, Earth Plant. Sci. Lett. 88, pp 29-44.
- Jafri, S.H. and Saxena, R., 1981. Geochemistry of anorthositic dykes from Nuggihalli schist belt of Dharwar craton, Karnataka, India. Jour.Geol.Soc. India, Vol.22, pp.85-91.
- James, H.L., 1954. Sedimentary facies iron-formation. Econ.Geol. Vol.49. pp.253-293.
- James, H.L., 1955. Zones of regional metamorphism in the Precambrians of northern Michigan. Geol.Soc. Am. Bull., 66. pp.1455-1488.
- James, H.L., 1966. Chemistry of iron rich sedimentary rocks. U.S.Geol.Surv., Prof. Pap., 440-W. pp.1-61.
- Janardhan, A.S., Newton, R.C. and Hansen, E.C., 1982. The transformation of amphibolite facies gneiss to charnockite in Southern Karnataka and Northern Tamil Nadu, India. Contrib. Min.Petrol. Vol.79. pp.139-149.
- Janardhan, A.S. and Gopalakrishna, D., 1983. Pressure-temperature estimates of the basic granulites and conditions of metamorphism in Sargur terrain, south Karnataka and adjoining areas. Jour.Geol.Soc.India, Vol.24. pp.219-228.
- Janardhan, A.S., Shadaksharaswamy, N. and Capdevila, R., 1986. Banded iron formation and associated manganese horizons of Sargur Supracrustals, Southern Karnataka. Jour.Geol.Soc.India., Vol.28. pp.179-188.
- Jayaram, B., 1922. Report on the work done during the field season 1917-1918. Rec.Mysore Geol.Dept.No, 18.pp.43-87.
- Jayaram, B., 1923. Progress report on the work done during the field season 1919-20. Rec.Mysore Geol. Dept., 20(2) pp.1-49.
- Jayaram, S., Venkatasubramanian, V.S. and Radhakrishna, B.P., 1983. Geochronology and trace element distribution in some tonalitic and granitic gneisses of the Dharwar craton. In. Naqvi, S.M. and Rogers, J.J.W. (Eds). Precambrian of South India: Geol.Soc.India. Mem. 4. pp.377-388.
- Kimberley, M.M., 1977. Geochemical distinctions among environmental types of iron-formations. Chem.Geol.

Vol.25, pp.185-212.

- Kimberley, M.M., 1978. Paleoenvironmental classification of iron-formations. *Econ.Geol.* Vol.73. pp.215-229.
- Klein, C., 1973. Changes in mineral assemblages with metamorphism of some banded Precambrian iron-formations. *Econ.Geol.*, Vol.68. pp.1075-1088.
- Klein, C., 1974. Greenalite, stilpnomelane, minnesotaite, crocidolite and carbonates in a very low grade metamorphic Precambrian iron-formations. *Can.Min.*, Vol.12, pp.475-498.
- Klein, C., 1978. Regional metamorphism of proterozoic iron-formation, Labrador Trough, Canada. *Am. Min.*, Vol.63. pp.898-912.
- Klein, C., 1983. Diagenesis and metamorphism of Precambrian banded iron-formations. In: A.F.Trendall and R.C.Morris (Eds). *Iron-formation: Facts and problems.* Elsevier. pp.417-470.
- Klein, C. and Bricker, O.P., 1977. Some aspects of the sedimentary and diagenetic environments of Proterozoic banded iron-formation. *Econ.Geol.*, Vol.72. pp.1457-1470.
- Klein, C., and Fink, R.P., 1976. Petrology of the Sokoman iron-formation in the Howells River area, at the western edge of the Labrador Trough. *Econ.Geol.*, Vol.71. pp.453-487.
- Klein, C., and Gole, M.J., 1981. Mineralogy and petrology of parts of the Marra Mamba iron-formation, Hamersley Basin, Western Australia. *Am.Min.*, Vol.66. pp.507-525.
- Klein, C., and Gole, M.J., 1983. Mineralogy and petrology of the Joffre member of the Brockman iron-formation, Hamersley Basin, Western Australia. *Am.Min.*
- Krauseopf, K.B., 1967. *Introduction to geochemistry.* McGraw Hill, New York. pp.721.
- Laajoki, K., 1975. Rare-earth elements in Precambrian iron-formation in Vayrylankyla, South puolanka area, Finland. *Bull.Geol.Soc.Finland.* Vol.47. pp.93-107.
- Laajoki, K. and Lavikainen, S., 1977. Rare-earth elements in the Archaean iron-formation and associated schists in Ukkolanvaara, 9, Lomantsi, S.E. Finland. *Bull.Geol.Soc. Finland.* Vol.47. pp.105-123.
- Laajoki, K. and Saikkonen, R., 1977. On the geology and geochemistry of the Precambrian iron-formation of Vayrytonkyla, South Puolanka area, Finland.



Bull.Geol.Sur. Finland, Vol.47. pp.93-107.

- Laaajoki, K. and Devaraju., T.C. 1989. Fe- chlorite, Grunerite, Stilpnomelane, Ankerite Siderite Occurrence in the Iron-Formation of Chiknayakanahalli Schist Belt, Karnataka. Geol. Sco. India., V.33, pp 175-182.
- Laberge, G.L., 1964. Development of magnetite in iron formations of the Lake Superior Region. Econ .Geol., v.59, pp. 1313-1342.
- Laberge, G.L., Robbins. E.I., and Han, T.M., 1987. A model for the biological precipitation of Precambrian iron formation - B: Morphologic evidence and modern analogs. In: P.W.U.Appel & Gene L.Laberge. Theophrastar Publ. S.A. Greece. pp.97-140.
- Landergren, S., 1948. On the geochemistry of Swedish iron ores and associated rocks. Sveriges. Geol.under.Ser.C 42(5). pp.1-182.
- Lepp, H. 1987. Chemistry and origin of Precambrian iron formations. In: Precambrian Iron- Formation: Appel, P.W.U., and LaBerge. G.L., Theophrastus. Athens. Greece.
- Lepp, H. and Goldich. S.S., 1964. Origin of Precambrian iron-formation. Econ .Geol. v.59, pp 1025-1060.
- Leshner, C.M., 1978. Mineralogy and petrology of the Sokoman iron-formation near Ardua Lake, Quebec. Can.Jour.Earth Sci. Vol.15(4).pp.480-500.
- McClaren, J.M., 1906. On some auriferous tracts in Southern India. Rec.Geol.Surv. India., Vol.34(2). pp.45-136.
- Mahabaleswar, B., and Naganna, C., 1981. Geothermometry of Karnataka Charnockites. Bull.Min., Vol.104. pp.848-855.
- Mahabaleswar, B., Basavanna, M., and Maaskant, P. 1987. Iron-formation of Archaean high-grade region, Karnataka, India. In: P.W.U.Appel and G.L.Laberge (Eds). Precambrian iron-formation. Theophrastar Publ., S.A. Greece. pp.487-512.
- Majumder. T., 1985. Environment and petrogenesis of the Precambrian banded iron-formation of the Iron Ore Group, Eastern India. Geol.Sur.Finland Bull, Vol.331. pp.201-213.
- Majumder, T., Pal.J.C., and Das, S.B., 1980. Trace element study of banded iron-formation and associated rocks of Orissa. Geol.Sur.India Spl.Publ. Series No.1, pp.587-597.
- Majumder, T., Chakraborty, K.L. and Bhattacharyya, A., 1982. Geochemistry of banded iron-formation of Orissa, India.

Geol.Sur.India. Spl.Publ. Series No.1. pp.587-597.

- Majumder, T., Whitley, J.E. and Chakraborty, K.L., 1984. Rare earth elements in the Indian banded iron-formation. *Chem.Geol.* Vol.45. pp.203-211.
- McLennan, S.M., Fryer, B.J. and Youg.G.M., 1979. Rare earth elements in Huronian (lower proterozoic) sedimentary rocks: Composition and evolution of the post Kemoran upper crust. *Geochim.Cosmochim.Acta*, Vol.43. pp.375-388.
- McLennan, S.M., Taylor, S.R., McGregor, V.R. 1984. Geochemistry of Archean metasedimentary rocks from west Greenland. *Geo chim. Cosmochim. Acta.*,v.48, pp 1-13.
- Mel'Nik, Y.P., 1982. Precambrian banded iron-formations. Elsevier. pp.1-31.
- Mishra, R.N., 1985. Precambrian BIFs of Karnataka: Abs. Geol. Soc. India. Group Discussion. Dharwad.
- Miyano, T., 1978b. Phase relations in the system Fe-Mg-Si-O-H and environments during low grade metamorphism of some Precambrian iron-formation. *Jour.Geol.Soc. Jpn.*, Vol.84. pp.679-690.
- Miyano, T., 1987. Diagenetic to low-grade metamorphic conditions of Precambrian iron-formation. In: Precambrian Iron-formation. P.W.U.Appel and Gene,L.Laberge (Eds). Theophrastar Publ., S.A.Greece. pp.155-186.
- Miyano, T., and Miyano,S., 1982. Ferri-annite from the Dales Gorge Member iron-formations, Wittenoom area, Western Australia. *Am.Min.* Vol.67. pp.1179-1195.
- Miyashiro., 1973. Metamorphism and metamorphic belts. John Wiley and sons. New York, pp 1- 492.
- Moorbath, S., 1976. Age and isotope constraints for the evolution of Archaean crust. In: B.F.Windley (Ed). The early history of the earth. Wiley London. pp.351-360.
- Morris, R.C., 1980. A textural and mineralogical study of relationships of iron ores to banded iron-formation in the Hamersley iron province of Western Australia. *Econ.Geol.* Vol.75. pp.184-209.
- Mukhopadhyay, D. Baral, M.C. and Neogi, R.K., 1980. Mineralogy of the banded iron formation in the southwestern Bababudan Hills, Karnataka, India. *N.Jb.Mineral. Abh.*, v.139(3), pp303- 327.

- Mukhopadhyay, D., 1986. Structural pattern in the Dharwar Craton. *Jour.Geo.*, Vol.94. pp.167-186.
- Mukhopadhyay, D., and Ghosh, D., 1983. Superposed deformation in the Dharwar rocks of the Southern part of the Chitradurga schist belt, near Dodguni, Karnataka. In: S.M.Naqvi and J.J.W.Rogers (Eds). *Precambrian of South India: Geol.Soc.India Mem 4*, pp.275-292.
- Naganna,C. and Shivakumr,B.S.,1977.Banded Iron Formation and the associated rocks of Kudremukh Region ,Karnataka; *Proc. Sym. Geology. etc. of ferrous and ferro-alloy minerals ., Bangalore.* pp 15-21.
- Naha. K. and Chatterjee, A.K., 1982. Axial plane folding in Bababudan hill ranges of Karnataka. *Indian Jour. Earth Sci.*, Vol.9, pp.37-43.
- Naha. K., Srinivasan,R and Naqvi,S.M., 1986. Structural unity in the early Precambrian Dharwar tectonic province, Peninsular India. *Quart.Jour.Geol. Min. Met. Soc. India.* Vol.58 (4). pp.215-243.
- Naqvi. S.M., 1981. The oldest supracrustal of Dharwar craton, India. *Jour.Geol.Soc. India.*, Vol.22. pp.458-469.
- Naqvi, S.M., 1983. Early Precambrian clastic metasediments of Dharwar Greenstone belts: Implications to sima-sial transformation processes. In: S.M.Naqvi and J.J.W.Rogers (Eds). *Precambrian of South India. Geol. Soc.India. Mem.4*, pp.309-328.
- Naqvi. S.M., Divakar Rao, V. and Hari Narain, 1978. The primitive crust: evidence from Indian shield. *Precam. Res.*, Vol.6. pp.323-345.
- Naqvi,S.M. and Hussain, S.M., 1979. Geochemistry of meta-anorthosites from a greenstone belt in Karnataka, India. *Can.Jour.Earth Sci.*, Vol.16. pp.1254-1264.
- Narayan Kutty, T.R., Anantha Iyer, G.V. and Ramakrishnan, M.,1974. Coexisting aegirine and magnesio-riebeckite from Bababudan hills, Mysore State. *Current Sci.* Vol.43. pp.1-3.
- Nautiyal, S.P., 1966. Precambrians of Mysore plateau. Presidential address to the Geology and Geography section. *Indian Sci.Cong. 53rd Session*, pp.1-14.
- Nicols, G.D. 1967. Trace elements in sediments; as assesment of their possible utility as depth indicators. *Marine Geol.*,v.5 , pp 539- 556.

- Parak, T., 1975. Kiruna iron ores are not "Intrusive-magmatic ores of the Kiruna type". *Econ.Geol.* 70. pp.1242-1258.
- Pichamuthu, C.S. 1935c. The banded ferruginous quartzites of the Bababudan hills, Mysore State. *Current Sci.*, Vol.3. pp.606-608.
- Pichamuthu, C.S., 1935f. The iron-formation and associated rocks of the Eastern Bababudans, Kadur district, Mysore. Half yly. *Jour. Mysore Univ.*, Vol.8, pp.1-48.
- Pichamuthu, C.S., 1951. Some observation on structure and classification of Dharwars of Mysore State. *Curr.Sci.* Vol.20. pp.117-119.
- Pichamuthu, C.S., 1952. Bababudanite. *Mysore.Geol. Dept. Rec.*, Vol.47, pp.1-26.
- Pichamuthu, C.S., 1953. Metamorphic facies in Mysore State. *Mys.Geol.Assn.inf. Circ.*, 47.
- Pichamuthu, C.S. 1974. On the banded iron formation of Precambrian age in India. *Jour.Geol.Soc. Ind.*, Vol.15, pp.1-30.
- Pichamuthu, C.S., and Srinivasan, R., 1983. A billion year history of the Dharwar craton. In: S.M.Naqvi and J.J.W.Rogers (Eds). *Precambrian of South India. Geol.Soc.Ind. Mem.*4. pp.121-142.
- Pichamuthu, C.S. and Srinivasan, R., 1984. The Dharwar craton - a Golden Jubilee publication: New Delhi. *India. Natl.Sci.Aca.* pp.34.
- Piper, D.Z., 1974. Rare-earth elements in ferro-manganese nodules and other marine phases. *Geochim.Cosmochim. Acta.* Vol.38. pp.1007-1022.
- Raase. P., Raith, M., Ackermant, D. and Lal, R.K., 1986. Progressive metamorphism of mafic rocks from greenschist to granulite facies in the Dharwar craton of South India. *Jour.Geol.*, Vol.94. pp.261-282.
- Radhakrishna, B.P., 1964. Some aspects of the stratigraphy and structure of Dharwar of Mysore State, South India, *Proc.Symp. on Stratigraphy, age and correlation of the Archaean provinces of India.* pp.33-34.
- Radhakrishna, B.P., 1967. Reconsideration of some problems in the Archaean complex of Mysore. *Jour.Geol. Soc. Ind.*Vol.18, pp.102-110.
- Radhakrishna, B.P., 1982. Archaean granite-greenstone terrain of the South Indian sheild. *Mem.Geol. Soc. Ind.*

Vol.4, pp.1-46.

- Radhakrishna, B.P. and Vasudev, V.N., 1977. The early Precambrian of the South Indian shield. Jour.Geol. Soc. Ind., Vol.18. pp.525-541.
- Radhakrishna, B.P., and Naqvi, S.M., 1986. Precambrian continental crust of India and its evolution. Jour. Geol. Vol.94. pp.145-166.
- Radhakrishna, B.P., Devaraju, T.C. and Mahabaleswar, B., 1986. B.I.F. of India: Jour.Geol.Soc.India, V. 28, pp 71-91.
- Raith, M., Raase, D., Ackermant, D., and Lal, R.K., 1983. Metamorphic condition in charnockite-khodalite zone of South India, Geothermobarometry on garnet-pyroxene-plagioclase rocks. In: Naqvi, S.M., and Rogers, J.J.W (Eds). Precambrian of South India. Geol.Soc.India. Mem. 4. pp.346-449.
- Rajagopalan, P.T., Jayaram, S. and Venkatasubramanian, V.S., 1980. Rb-Sr isochron ages of gneisses in the Western region of the Dharwar craton. Jour.Geol.Soc. India. Vol.21. pp.54-56.
- Rajamani, V., Shivkumar, K., Hanson, G.N. and Granath, J.W., 1981. Petrogenesis of amphibolites in the Kolar schist belt, India - a preliminary report. Jour.Geol.Soc. India, Vol.22 pp.470-487.
- Ramachandra Rao, M.B., 1938. A note on the Bababudanite-rhodusite relationship. Rec.Mysore Geol.Dept., Vol.37. pp.116-126.
- Ramakrishnan, M., Moorbath, S. Taylor, P.N., Ananth Iyer, G.V. and Viswanatha, M.N., 1984. Rb-Sr and Pb-Pb whole rock isochron ages of basement gneisses in Karnataka craton. Jour.Geol.Soc. India, Vol.25. pp.20-34.
- Ramakrishnan, M., and Viswanatha, M.N., 1987. Angular unconformity, structural unity argument and Sargur-Dharwar relations in Bababudan basin. Jour.Geol.Soc. India. Vol.29. pp.471-482.
- Rama Rao, B., 1936a. Note on some conglomerates of Mysore. Rec. Mysore Geol.Dept. Vol.34. pp.33-45.
- Rama Rao, B., 1940. Archaean complex of Mysore. Mysore Geol. Dept. Bull. 17. pp.95.
- Rama Rao, B., 1962. A hand book of the geology of Mysore State, Southern India. Bangalore Printing & Publishing Co., Bangalore. pp.264.
- Ramakrishnan, M. and Viswanatha, M.N., 1978. Charnockite-

gneiss complex relationship in Southern Karnataka. Jour.Geol.Soc. India., Vol.19. pp.411-419.

Roger, J.J.W., 1986. The Dharwar craton and the assembly of Peninsular India. Jour.Geol. Vol.94(2). pp.129-143.

Rollinson, H.R., Windley, B.F. and Ramakrishnan, M., 1981. Contrasting high intermediate pressure of metamorphism in the Archaean Sargur schist of South India. Contrib. Min. Petrol. Vol.76. pp.420-429.

Sakamoto, T. 1950. The origin of the Precambrian banded iron ores: Am, Jour. Sci. V.248, PP449-474.

Sambashiva Iyer, V.S., 1899. Report on geological survey work in the Chitradurga and Tumkur districts during the field seasons 1896-1899. Rec.Mysore Geol.Dept. Vol.2, pp.82-106.

Sampath Iyengar, P., 1908. Report on the geology of the part of Hassan and Kadur districts. Rec. Mysore Geol. Dept., Vol.9. pp.59-84

Sampath Iyengar, P., 1909. Geology of the western portion of Hassan district. Rec. Mysore Geol. Dept., Vol.10. pp.31-62.

Sampath Iyengar, P., 1915. The schistose rocks of the Bababudans with special reference to the auriferous series in their neighbourhood. Rec. Mysore Geol. Dept. pp.109-134.

Semenko, M.P., 1973. Genesis of Precambrian Iron-Manganese Deposits: Proc. Kiev. Symp, 1970. UNESCO, Paris.

Shatskiy, N.S., 1954. Quoted by Melnik: In. Precambrian banded iron formations. Elsevier. Publ.

Shivakumar, B.S., 1976. Study of the iron ores of Sandur schist belt Bellary Dist. Karnataka State, In: studies in Precambrian. C.Naganna (Ed) Dept. Geology, Bagalore University. Bangalore.

Simonsen, B.M., 1985. Sedimentological constraints on the origins of Precambrian iron formations. Geol, Soc. Am. Bull., v.96, pp244-252.

Slater, H.K., 1899. Preliminary report on the geological survey work in the Shimoga district. Rec. Mysore Geol. Dept. Vol.2. pp.118-130.

Slater, H.K., 1904. Report on survey work in portions of the Shimoga and Kadur districts. Rec. Mysore Geol. Dept. Vol.5. pp.35-56.

- Slater, H.K., 1908. Report on the geological survey of portion of Kadur district. Rec. Mysore Geol. Dept. Vol.9. pp.35-58.
- Smeeth, W.F., 1899. General report for the year 1898-1899. Rec. Mysore Geol. Dept. Vol.2. pp.1-51.
- Smeeth, W.F., 1901. General report on the work of the Department for the year 1900-1901. Rec. Mysore Geol. Dept. Vol.3. pp.1-43.
- Smeeth, W.F., 1908. General report of the work of the Department for the year 1907-1908. Rec. Mysore Geol. Dept. Vol.9. pp.1-25.
- Smeeth, W.F., 1915. General report of the year 1914. Rec. Mysore Geol. Dept., Vol.14. pp.21-46.
- Smeeth, W.F., 1916. Outline of the geological history of Mysore. Mysore Geol. Dept. Bull.6, pp.22.
- Srikantappa, C., Raith, M. and Ackermant, D., 1985. High-grade regional metamorphism of ultramafic and mafic rocks from the Archaean Sargur terrain, Karnataka, South India. Precamb.Res., Vol.24.
- Srinivasan, R. and Sreenivas, B.L., 1972. Dharwar stratigraphy. Jour. Geol.Soc. India. Vol.13. pp.72-83.
- Srinivasan, R. and Sreenivas, B.L., 1975. A reappraisal of some aspects of the geology of Dharwar of Mysore State, India. Geol.Sur. India Misc.Publ.23(1). pp.49-69.
- Stroh, P.T., Monrad, J.R., Fullagar, P.D., Naqvi, S.M., Hussain, S.M. and Rogers, J.J.W., 1982. 3,000 m.y. old Halekote trondhjemite, a record of stabilization of the Dharwar Precambrian of South India. Geol.Sur. India. Mem.4. pp.365-376.
- Swami Nath. J., Ramakrishnan, M. and Viswanatha, M.N., 1976. Dharwar stratigraphic model and Karnataka craton evolution. Rec. Geol. Surv. India. Vol.107. pp.149-175.
- Swami Nath, J. and Ramakrishnan, M., 1981. Early Precambrian supracrustals of South Karnataka. Geol. Surv. India. Mem.112. pp.350.
- Taylor, S.R. and McLennan, S.M., 1981. The composition and evolution of the continental crust, rare earth evidence from sedimentary rocks. Phil.Trans.R.Soc. London, Ser.A, 301: pp.381-399.
- Taylor, P.N., Moorbath, S., Chadwick, B., Ramakrishnan and Viswanatha, M.N., 1984. Petrography, chemistry and

T 5679

isotopic ages of Peninsular gneiss, Dharwar acid volcanic rocks and the Chitradurga granite with special reference to the late Archaean evolution of the Karnataka craton, Southern India. Preamb.REs., Vol.23. pp.349-375.

- Trendall, A.F., 1973. Iron formation of the Hamersley Group of manganese deposits, UNESCO, Paris, Earth Science, No.9, pp. 227-268.
- Trendall, A.F., 1983. Introduction: In: Iron-formation: Facts and Problems. Elsevier, pp.1-12.
- Trendall, A.F., and Blockley, J.G., 1970. The iron-formation of the Precambrian Hamersley Group, W. Australia. Geological Survey of Western Australia Bulletin. v. 119, pp 1-366.
- Van Hise, C.R. and Leith, C.K. 1911. The Geology of the Lake Superior region, Mon. USGS - LII.52. pp. 641.
- Veblen, D.R. and Ribbe, P.H. 1982. Reviews in mineralogy. V. 9B, Mineralogical Society of America.
- Venkatasubramanian, V.S., 1974. Geochronology of the Dharwar craton: a review. Jour.Geol.Soc. India. Vol.15. pp.463-468.
- Viswanath, S. 1974. Basaltic komatiite occurrences in the Kolar Gold Field of India. Geol. Mag. v. 111, pp 353-354.
- Viswanath, M.N. and Ramakrishnan, M., 1981. Bababudan belt. In: J.Swami Nath and M.Ramakrishnan (Eds). Early Precambrian Supracrustals of Southern Karnataka. Mem. Geol. Surv. India. Vol.112. pp.91-114.
- Viswanathaiah, M.N., and Venkatachalapathy, V., 1980. Microbiota from the Bababudan iron-formation, Karnataka. Geol.Soc.India. Vol.21. pp.16-20.
- Walter, M.R. and Hofmann, 1983. In; A.F. Trendall and R.C. Morris( Eds.). Iron-formation: Facts and Problems, Elsevier, pp. 373-396.
- Wetherell, E.W., 1904. Report on the country lying between 76 35' and 76 59' longitude and between 13 9'E and 14 10'N latitude in parts of Chitaldurg and Tumkur districts. Rec. Mysore Geol. Dept. Vol.5(2). pp.1-34.
- Windley, B.F. and Bridgwater, D., 1971. The evolution of Archaean low and high grade terrains. Geol. Soc. Austr. Spel. Publ. Vol.3, pp.33-46.
- Zelenov 1963 . Quoted by Mel'Nik, Y.P., 1982. Precambrian banded iron formation , Elsevier, pp 1- 310.