

LIST OF FIGURES

S. No.	Title	After Page
1	Determination of geographical coordinates of the selected maize accessions using Google Earth	33
2	Locations of collection of selected maize landrace accessions from various agro-ecological zones of India (as defined by AICRP on Maize)	33
3	Representative ears of maize accessions analyzed in this study	49
4	Representative ears of trial entries (Bajaura <i>Kharif</i> 2006)	51
5	Representative ears of trial entries (Delhi <i>Kharif</i> 2007)	53
6	Representative ears of trial entries (Hyderabad <i>Rabi</i> 2006-07)	55
7	Dendrogram depicting the clustering of the selected Indian maize landrace accessions using Euclidean distance and Ward's method of minimum variance, based on nine morpho-agronomic traits recorded at Bajaura	59
8	Principal component analysis of phenotypic data recorded on the selected maize landrace accessions	61
9	DNA sequencer images of SSR alleles of the loci (labelled with FAM dye)	62
10	DNA sequencer images of SSR alleles of the loci (labelled with HEX dye)	62
11	DNA sequencer images of SSR alleles of the loci (labelled with TET dye)	62
12	Graphical representation of pair-wise genetic similarities based on Roger's distance (1972)	63
13	Rogers' genetic distance-based unrooted tree depicting the genetic relationships among the selected maize landrace accessions	64
14	Plot of ΔK values, the second order rate of change of $\ln P(D)$, in relation to the number of inferred clusters (K)	65
15	Model-based population structure analysis of the Indian maize landraces implemented using STRUCTURE	65
16	POWERCORE-based identification of a 'minicore' based on phenotypic analysis of Bajaura data and molecular data	66
17	'Sikkim Primitives' showing prolificacy at Bajaura (<i>Kharif</i> 2008)	68
18	Representative ears of maize landraces/inbreds (Bajaura <i>Kharif</i> 2008) evaluated for prolificacy	68
19	Representative ears of maize landraces (Chinyalisaur <i>Kharif</i> 2008) evaluated for prolificacy	68
20	<i>tb1</i> allele mining using primer pair <i>tb1-3</i> : (a) A 550 bp amplicon resolved in three replicates in 1% agarose gel after purification. (b) Alignment of the sequences, revealing SNPs and InDels, particularly in 'Sikkim Primitives'	69
21	<i>tb1</i> allele mining using primer pair <i>tb1-5</i> : (a) A 532 bp amplicon resolved in three replicates in 1% agarose gel after purification. (b) Alignment of the sequences of the amplicons, revealing SNPs and InDels in the Sikkim Primitive IML602	71

S. No.	Title	After Page
22	Plot of LD decay in terms of r^2 (square of the correlation coefficient between two markers) against distances in bp for the 1.1 kb region of 5' NTR, upstream of <i>tb1</i> gene	73
23	Linkage disequilibrium (LD) matrix for polymorphic sites within 5' NTR region of <i>tb1</i> gene	73
24	Six generations of the cross DMRQPM-28-03 x CM139, evaluated for prolificacy, at Bajaura in <i>Kharif</i> 2008	76
25	Six generations of the cross DMRQPM-28-03 x CM140, that showed maximum heritability for prolificacy, evaluated at Delhi in <i>Kharif</i> 2008	76
26	Six generations of the cross DMRQPM-28-03 x CM139 evaluated for prolificacy, at Meerut in <i>Kharif</i> 2008	76