

References



References

- ❖ Abbas AK, Lichtman AH. Autoimmunity. In: Cellular and Molecular Immunology, 5th ed. Philadelphia: Elsevier Science Publications, 2003; p. 217-239.
- ❖ Abraham DJ, Varga J. Scleroderma: from cell and molecular mechanisms to disease models. Trends Immunol. 2005; 26(11):587–595.
- ❖ Agarwal K, Jones DE, Daly AK, James OF, Vaidya B, Pearce S, Bassendine MF. CTLA-4 gene polymorphism confers susceptibility to primary biliary cirrhosis. J Hepatol. 2000; 32(4):538-541.
- ❖ Ahmad YA, Bruce IN. Genetic epidemiology: systemic lupus erythematosus. Arthritis Res Ther. 2001; 3(6):331–336.
- ❖ Ahmed AR, Rosen GB. Viruses in pemphigus. Int J Dermatol. 1989; 28(4): 209-217.
- ❖ Ajjan RA, Watson PF, Weetman AP. Cytokines and thyroid function. Adv Neuroimmunol. 1996; 6(4):359–386.
- ❖ Akbar AN, Salmon M, Savill J, Janossy G. A possible role for bcl-2 in regulating T cell memory - a balancing act between cell-death and survival. Immunol Today. 1993; 14(11):526-532.
- ❖ Akbar AN, Terry L, Timms A, Beverley PC, Janossy G. Loss of CD45R and gain of UCHL1 reactivity is a feature of primed T cells. J Immunol. 1988; 140(7):2171–2178.
- ❖ Alcocer-Varela J, Alarcon-Segovia D. Decreased production of and response to IL-2 by cultured lymphocytes from patients with Systemic Lupus Erythematosus. J Clin Invest. 1982; 69(6):1388–1392.
- ❖ Allan SE, Crome SQ, Crellin NK, Passerini L, Steiner TS, Bacchetta R, Roncarolo MG, Levings MK. Activation-induced FoxP3 in human T-effector cells does not suppress proliferation or cytokine production. Int Immunol. 2007; 19(4):345–354.
- ❖ Al-Lazikani B, Lesk AM, Chothia C. Canonical structures for the hypervariable regions of T cell $\alpha\beta$ receptors. J Mol Biol. 2000; 295:979-995.
- ❖ Allison JP, McIntyre BW, Bloch D. Tumor-specific antigen of murine T lymphoma defined with monoclonal antibody. J Immunol. 1982; 129(5):2293-2300.
- ❖ Al-Mutairi S, Al-Awadhi A, Raghupathy R, Al-Khawari H, Sada P, Al-Herz A, Rawoot P. Lupus patients with pulmonary involvement have a pro-inflammatory cytokines profile. Rheumatol Int. 2007; 27(7):621–630.

-
- ❖ Alococer-Varela J, Alarcon-Segovia D. Decreased production of and response to interleukin 2 by cultured lymphocytes from patients with systemic lupus erythematosus. *J Clin Invest.* 1982; 69(6):1388-1392.
 - ❖ Alvarado-Sa´nchez B, Herna´ndez-Castro B, Portales-Pe´rez D, Baranda L, Layseca-Espinosa E, Abud-Mendoza C, Cubillas-Tejedab AC, Gonza´lez-Amaro R. Regulatory T cells in patients with systemic lupus erythematosus. *J Autoimmun.* 2006; 27(2):110–118.
 - ❖ Amagai M, Karpati S, Prussick R, Klaus-Kovtun V, Stanley JR. Autoantibodies against the amino-terminal cadherin-like binding domain of pemphigus vulgaris antigen are pathogenic. *J Clin Invest.* 1992; 90(3):919–926.
 - ❖ Amagai M, Klaus-Kovtun V, Stanley JR. Autoantibodies against a novel epithelial cadherin in pemphigus vulgaris, a disease of cell adhesion. *Cell.* 1991; 67(5):869–877.
 - ❖ American College of Rheumatology. Guidelines for referral and management of systemic lupus erythematosus in adults. *Arthritis Rheum* 1999; 42(9):1785-96.
 - ❖ Amzel LM, Poljak RJ, Sank F, Varga JM, Richards FFR. The three dimensional structure of a combining region-ligand complex of immunoglobulin NEW at 3.5A resolution. *Proc Natl Acad Sci USA.* 1974; 71(4):1427-1430.
 - ❖ Anaya JM, Correa PA, Herrera M, Eskdale J, Gallagher G. Interleukin 10 (IL-10) influences autoimmune response in primary Sjögren’s syndrome and is linked to IL-10 gene polymorphism. *J Rheumatol.* 2002; 29(9):1874–1876.
 - ❖ Andersen HR, Vinggaard AM, Rasmussen TH, Gjermansen IM, Bonefeld-Jorgensen EC. Effects of currently used pesticides in assays for estrogenicity, androgenicity, and aromatase activity in vitro. *Toxicol Appl Pharmacol.* 2002; 179(1):1–12.
 - ❖ Anderson MS, Venanzi ES, Klein L, Chen Z, Berzins SP, Turley SJ, Boehmer HV, Bronson R, Dierich A, Benoist C, Mathis D. Projection of an immunological self shadow within the thymus by the aire protein. *Science.* 2002; 298(5597):1395–1401.
 - ❖ Andriankes AA, Sharp JT, Person DA, Duffy J. Cell mediated immunity in rheumatoid arthritis. *Ann Rheum Dis.* 1977; 36(1):13-20.

- ❖ Anhalt GJ, Labib RS, Voorhees JJ, Beals TF, Diaz LA. Induction of Pemphigus in neonatal mice by passive transfer of IgG from patients with the disease. *N Engl J Med.* 1982; 306(20):1189–1196.
- ❖ Ansel KM, Lee DU, Rao A. An epigenetic view of helper T cell differentiation. *Nat Immunol.* 2003; 4(7):616-623.
- ❖ Arend WP, Gabay C. Cytokines in the rheumatic diseases. *Rheum Dis Clin North Am.* 2004; 30(1):41–67.
- ❖ Artlett CM, Smith JB, Jimenez SA. Identification of fetal DNA and cells in skin lesions from women with systemic sclerosis. *N Engl J Med.* 1998; 338(17):1186-1191.
- ❖ Awad MR, El-Gamel A, Hasleton P, Turner DM, Sinnott PJ, Hutchinson IV. Genotypic variation in the transforming growth factor beta1 gene: association with transforming growth factor production, fibrotic lung disease, and graft fibrosis after lung transplantation. *Transplantation.* 1998; 66(8):1014–1020.
- ❖ Awata T, Kurihara S, Iitaka M, Takei S, Inoue I, Ishii C, Negishi K, Izumida T, Yoshida Y, Hagura R, Kuzuya N, Kanazawa Y, Katayama S. Association of CTLA-4 gene A-G polymorphism (IDDM12 locus) with acute-onset and insulin-depleted IDDM as well as autoimmune thyroid disease (Graves' disease and Hashimoto's thyroiditis) in the Japanese population. *Diabetes.* 1998; 47(1):128-129.
- ❖ Backer LC. Pet dogs as sentinels for human exposure to environmental pollution. In: Butterworth FM, Corkum LD, Guzmán-Rincon J, editors. *Biomonitoring and biomarkers as indicators of environmental change.* New York: Plenum Press, 1995; p. 193–206.
- ❖ Baecher-Allan C, Brown JA, Freeman GJ, Hafler DA. CD4+CD25 high regulatory cells in human peripheral blood. *J Immunol.* 2001; 167(3):1245–1253.
- ❖ Baldwin TA, Hogquist KA, Jameson SC. The fourth way? Harnessing aggressive tendencies in the thymus. *J Immunol.* 2004; 173(11):6515–6520.
- ❖ Ballestar E, Esteller M, Richardson BC. The epigenetic face of systemic lupus erythematosus. *J Immunol.* 2006; 176(12):7143-7147.
- ❖ Ballou SP, Khan MA, Kushner I. Clinical features of systemic lupus erythematosus. Differences related to race and age of onset. *Arthritis Rheum.* 1982; 25(1):55-60.

-
- ❖ Banerjee BD, Koner BC, Ray A. Immunotoxicity of pesticides: Perspectives and trends. *Indian J Exp Biol.* 1996; 34(8):723–733.
 - ❖ Baraut J, Michel L, Verrecchia F, Farge D. Relationship between cytokine profiles and clinical outcomes in patients with systemic sclerosis. *Autoimmun Rev.* 2010; 10(2):65-73.
 - ❖ Barcellini W, Rizzardi GP, Borghi MO, Nicoletti F, Fain C, Del Papa N, Meroni PL. In vitro type-1 and type-2 cytokine production in systemic lupus erythematosus: lack of relationship with clinical disease activity. *Lupus.* 1996; 5(2):139–145.
 - ❖ Barnett AJ, Miller MH, Littlejohn GO. A survival study of patients with scleroderma diagnosed over 30 years (1953–1983): the value of a simple cutaneous classification in the early stages of the disease. *J Rheumatol.* 1988; 15(2):276–283.
 - ❖ Barnett AJ, Tait BD, Barnett MA, Toh BH. T lymphocyte subset abnormalities and HLA antigens in scleroderma (Systemic sclerosis). *Clin Exp Immunol.* 1989; 76(1):24-29.
 - ❖ Barnett JB, Rodgers HE. Pesticides. In: Dean JH, Luster MI, Munson AE, Kimber I, editors. *Immunotoxicology and Immunopharmacology*, 2nd ed. New York: Raven Press, 1994; p. 191–212.
 - ❖ Barnhill RL. *Textbook of Dermatopathology*, 1st ed. New York: Mc Graw-Hill, 1998.
 - ❖ Baroja ML, Vijayakrishnan L, Bettelli E, Darlington PJ, Chau TA, Ling V, Collins M, Carreno BM, Madrenas J, Kuchroo VK. Inhibition of CTLA-4 functions by the regulatory subunit of serine/threonine phosphatase 2A. *J Immunol.* 2002; 168(10):5070–5078.
 - ❖ Barzilai O, Sherer Y, Ram M, Izhaky D, Anaya JM, Shoenfeld Y. Epstein-Barr virus and cytomegalovirus in autoimmune diseases: are they truly notorious? A preliminary report. *Ann N Y Acad Sci.* 2007; 1108:567-577.
 - ❖ Becker KG. Comparative genetics of type 1 diabetes and autoimmune disease: common loci, common pathways? *Diabetes.* 1999; 48(7):1353-1358.
 - ❖ Belkaid Y, Rouse BT. Natural regulatory T cells in infectious disease. *Nat Immunol.* 2005; 6:353–360.
 - ❖ Bellaney G, Rycroft R. Pemphigus vulgaris following a hyperimmune response to typhoid booster. *Clin Exp Dermatol.* 1996; 21(6):434- 436.

-
- ❖ Bendelac A, Rivera MN, Park SH, Roark JH. Mouse CD1-specific NK1 T cells: development, specificity and function. *Annu Rev Immunol.* 1997; 15:535–562.
 - ❖ Benedek TG. History of the Rheumatic Diseases. In: Klippel JH, editor. *Primer on the Rheumatic Diseases*, 11th ed. Atlanta: Arthritis Foundation, 1997.
 - ❖ Bentley GA, Mariuzza RA. The structure of the T cell antigen receptor. *Annu Rev Immunol.* 1996; 14:563-590.
 - ❖ Bentwich Z, Kalinkovich A, Weisman Z, Borkow G, Beyers N, Beyers AD. Can eradication of helminthic infections change the face of AIDS and Tuberculosis? *Immunol Today.* 1999; 20(11):485-487.
 - ❖ Bentwich Z, Kalinkovich A, Weisman Z. Immune activation is a dominant factor in the pathogenesis of AIDS in Africa. *Immunol Today.* 1995; 16(4):187-191.
 - ❖ Beretta L, Bertolotti F, Cappiello F, Barili M, Masciocchi M, Toussoun K, Caronni M, Scorza R. Interleukin-1 gene complex polymorphisms in systemic sclerosis patients with severe restrictive lung physiology. *Hum Immunol.* 2007a; 68(7):603–609.
 - ❖ Beretta L, Cappiello F, Barili M, Scorza R. Proximal interleukin-10 gene polymorphisms in Italian patients with systemic sclerosis. *Tissue Antigens.* 2007b; 69(4):305–312.
 - ❖ Bette M, Schafer MK, van Rooijen N, Weihe E, Fleischer B. Distribution and kinetics of superantigen-induced cytokine gene expression in mouse spleen. *J Exp Med.* 1993; 178(5):1531–1539.
 - ❖ Bettelli E, Korn T, Kuchroo VK. Th17: the third member of the effector T cell trilogy. *Curr Opin Immunol.* 2007; 19(6):652–657.
 - ❖ Beverly B, Kang SM, Leonardo MJ, Schwartz RH. Reversal of in vitro T cell clonal anergy by IL-2 stimulation. *Int Immunol.* 1992; 4(6):661-671.
 - ❖ Bhol KC, Desai A, Kumari S, Colon JE, Ahmed AR. Pemphigus vulgaris: the role of IL-1 and IL-1 receptor antagonist in pathogenesis and effects of intravenous immunoglobulin on their production. *Clin Immunol.* 2001; 100(2):172–180.
 - ❖ Bhol KC, Rojas AI, Khan IU, Ahmed AR. Presence of interleukin 10 in the serum and blister fluid of patients with pemphigus vulgaris and pemphigoid. *Cytokine.* 2000; 12(7):1076–1083.

-
- ❖ Bickle KM, Roark TR, Sylvia HSU. Autoimmune Bullous Dermatoses: A Review. *American Family Physician*. 2002; 65(9):1861-1870.
 - ❖ Bidwell J, Keen L, Gallagher G, Kimberly R, Huizinga T, McDermott MF, Oksenberg J, McNicholl J, Pociot F, Hardt C, D'Alfonso S. Cytokine gene polymorphism in human disease: on-line databases. *Genes Immun*. 1999; 1(1):3-19.
 - ❖ Bierer BE, Burakoff SJ. T-lymphocyte activation: the biology and function of CD2 and CD4. *Immunol Rev*. 1989; 111:267-294.
 - ❖ Billiau A, Heremans H, Vermeire K, Matthys P. Immunomodulatory properties of interferon-gamma. An update. *Ann NY Acad Sci*. 1998; 856:22-32.
 - ❖ Bjorses P, Aaltonen J, Horelli-Kuitunen N, Yaspo ML, Peltonen L. Gene defect behind APECED: a new clue to autoimmunity. *Hum Mol Genet*. 1998; 7(10):1547-1553.
 - ❖ Blakemore AI, Tarlow JK, Cork MJ, Gordon C, Emery P, Duff GW. Interleukin-1 receptor antagonist gene polymorphism as a disease severity factor in systemic lupus erythematosus. *Arthritis Rheum*. 1994; 37(9):1380-1385.
 - ❖ Blakemore AIF, Cox A, Gonzalez AM, Maskil JK, Hughes ME, Wilson RM, Ward JD, Duff GW. Interleukin-1 receptor antagonist allele (IL1RN*2) associated with nephropathy in diabetes mellitus. *Hum Genet*. 1996; 97(3):369-374.
 - ❖ Blakemore AIF, Watson PF, Weetman AP. Association of Graves' disease with an allele of the interleukin-1 receptor antagonist gene. *J Clin Endocrinol Metab*. 1995; 80(1):111-115.
 - ❖ Blasini AM, Stekman IL, Leon-Ponte M, Caldera D, Rodriguez MA. Increased proportion of responders to a murine anti-CD3 monoclonal antibody of the IgG1 class in patients with systemic lupus erythematosus (SLE). *Clin Exp Immunol*. 1993; 94(3):423-428.
 - ❖ Boehm T, Rabbitts TH. The human T cell receptor genes are targets for chromosomal abnormalities in T cell tumors. *FASEB J*. 1989; 3:2344-2359.
 - ❖ Boin F, Wigley FM, Schneck JP, Oelke M, Rosen A. Evaluation of topoisomerase-1-specific CD8+ T-cell response in systemic sclerosis. *Ann N Y Acad Sci*. 2005; 1062:137-145.

-
- ❖ Boles JW, Pitt ML, LeClaire RD, Gibbs PH, Torres E, Dyas B, Ulrich RG, Bavari S. Generation of protective immunity by inactivated recombinant staphylococcal enterotoxin B vaccine in nonhuman primates and identification of correlates of immunity. *Clin Immunol*. 2003; 108(1):51–59.
 - ❖ Bolster MB, Ludwicka A, Sutherland SE, Strange C, Silver RM. Cytokine concentrations in bronchoalveolar lavage fluid of patients with systemic sclerosis. *Arthritis Rheum*. 1997; 40(4):743-751.
 - ❖ Bonelli M, Savitskaya A, Steiner CW, Rath E, Smolen JS, Scheinecker C. Phenotypic and functional analysis of CD4+ CD25- Foxp3+ T cells in patients with systemic lupus erythematosus. *J Immunol*. 2009; 182(3):1689–1695.
 - ❖ Bonelli M, Savitskaya A, Steiner CW, Rath E, Smolen JS, Scheinecker C. Phenotypic and functional analysis of CD4+ CD25– Foxp3+ T cells in patients with systemic lupus erythematosus. *J Immunol*. 2009; 182(3):1689-1695.
 - ❖ Bonelli M, Smolen JS, Scheinecker C. Treg and lupus. *Ann Rheum Dis*. 2010; 69:i65–i66.
 - ❖ Bonelli M, von Dalwigk K, Savitskaya A, Smolen JS, Scheinecker C. Foxp3 expression in CD4+ T cells of patients with systemic lupus erythematosus: a comparative phenotypic analysis. *Ann Rheum*. 2008; 67:664–671.
 - ❖ Borkow G, Leng Q, Weisman Z, Stein M, Galai N, Kalinkovich A, Bentwich Z. Chronic immune activation associated with intestinal helminthic infection results in impaired signal transduction and anergy. *J Clin Investig*. 2000; 106(8):1053-1060.
 - ❖ Boyum A. Isolation of mononuclear cells and granulocytes from human blood. Isolation of monuclear cells by one centrifugation, and of granulocytes by combining centrifugation and sedimentation at 1 g. *Scand J Clin Lab Invest Suppl*. 1968; 97:77–89.
 - ❖ Bremell T, Tarkowski A. Preferential induction of septic arthritis and mortality by superantigen-producing staphylococci. *Infect Immun*. 1995; 63(10):4185–4187.
 - ❖ Brenner S, Bialy Golan A, Anhalt G. Recognition of pemphigus antigens in drug-induced pemphigus vulgaris and pemphigus foliaceus. *J Am Acad Dermatol*. 1997; 36(6 pt 1):919-923.

-
- ❖ Brenner S, Mashiah J, Tamir E, Goldberg I, Wohl Y. Pemphigus: An acronym for a disease with multiple causes. *SKINmed*. 2003; 2:163–167.
 - ❖ Brenner S, Ruocco V. Pemphigus registry: genetic and inducing factors to be explored. *Int J Dermatol*. 1994; 33(4):287-289.
 - ❖ Brenner S, Wolf R, Ruocco V. Contact pemphigus: a subgroup of induced pemphigus. *Int J Dermatol*. 1994; 33:843–845.
 - ❖ Brinkman BM, Zuijdgheest D, Kaijzel EL, Breedveld FC, Verweij CL. Relevance of the tumor necrosis factor alpha (TNF alpha)-308 promoter polymorphism in TNF alpha gene regulation. *J Inflamm*. 1996; 46(1):32–41.
 - ❖ Brocke S, Gaur A, Piercy C, Gautam A, Gijbels K, Fathman CG, Steinman L. Induction of relapsing paralysis in experimental autoimmune encephalomyelitis by bacterial superantigen. *Nature*. 1993; 365(6447):642–644.
 - ❖ Brocke S, Hausmann S, Steinman LW, Wucherpfennig KW. Microbial peptides and superantigens in the pathogenesis of autoimmune diseases of the central nervous system. *Semin Immunol*. 1998; 10(1):57–67.
 - ❖ Bronson R, Brenner S, Frusic Zlotkin M, Michelle B, Tamir A, Milner Y. Circulating pemphigus IgG in families of patients with pemphigus. *J Am Acad Dermatol*. 1997; 36(1):44-52.
 - ❖ Broughton A, Thrasher JD. Chronic health effects and immunological alterations associated with exposure to pesticides. *Comm Toxicol*. 1990; 4:59-71.
 - ❖ Buchman AL. Side effects of corticosteroid therapy. *J Clin Gastroenterol*. 2001; 33(4):289-294.
 - ❖ Buckner JH. Mechanisms of impaired regulation by CD4 (+) CD25 (+) FOXP3 (+) regulatory T cells in human autoimmune diseases. *Nat Rev Immunol*. 2010; 10:849–859.
 - ❖ Buonpane RA, Churchill HR, Moza B, Sundberg EJ, Peterson ML, Schlievert PM, Kranz DM. Neutralization of staphylococcal enterotoxin B by soluble, high-affinity receptor antagonists. *Nat Med*. 2007; 13(6):725–729.
 - ❖ Burns C, Burns P, Holsapple M. “Of Mice and Men” (John Steinbeck) — How do we determine the potential for immunotoxicity in humans? *Ann Epidemiol*. 2000; 10(7): 471– 472.

-
- ❖ Bush B, Snow J, Koblitz R. Polychlorobiphenyl (PCB) congeners, p, p'-DDE and hexachlorobenzene in maternal and fetal cord blood from mothers in upstate New York. *Arch Environ Contam Toxicol*. 1984; 13(5):517–527.
 - ❖ Campbell DJ, Koch MA. Phenotypical and functional specialization of FOXP3+ regulatory T cells. *Nat Rev Immunol*. 2011; 11(2):119–130.
 - ❖ Campbell DJ, Ziegler SF. FoxP3 modifies the phenotypic and functional properties of regulatory T-cells. *Nat Rev Immunol*. 2007; 7(4):305–310.
 - ❖ Cantagrel A, Navaux F, Loubet-Lescoulié P, Nourhashemi F, Enault G, Abbal M, Constantin A, Laroche M, Mazières B. Interleukin-1beta, interleukin-1 receptor antagonist, interleukin-4, and interleukin-10 gene polymorphisms. Relationship to occurrence and severity of rheumatoid arthritis. *Arthritis Rheum*. 1999; 42(6):1093–1100.
 - ❖ Caproni M, Giomi B, Cardinali C, Salvatore E, Pestelli E, D'Agata A, Bianchi B, Toto P, Feliciani C, Fabbri P. Further support for a role for Th2-like cytokines in blister formation of pemphigus. *Clin Immunol*. 2001; 98(2):264–271.
 - ❖ Carlsson R, Sjogren HO. Kinetics of IL-2 and IFN gamma production, expression of IL-2 receptors, and cell proliferation in human mononuclear cells exposed to SEA. *Cell Immunol*. 1985; 96(1):175–183.
 - ❖ Carvalho LP, Bacellar O, Neves N, de Jesus AR, Carvalho EM. Downregulation of IFN-gamma production in patients with recurrent vaginal candidiasis. *J Allergy Clin Immunol*. 2002; 109(1):102–105.
 - ❖ Casciola-Rosen LA, Anhalt G, Rosen A. Autoantigens targeted in systemic lupus erythematosus are clustered in two populations of surface structures on apoptotic keratinocytes. *J Exp Med*. 1994; 179(4):1317-1330.
 - ❖ Cervera R, Khamashia MA, Font J, Sebastiani GD, Gil A, Lavilla P, Doménech I, Aydintug AO, Jedryka-Góral A, de Ramón E, et al. SLE: Clinical and immunologic patterns of disease expression in a cohort of 1000 patients. *Medicine*. 1993; 72(2):113-124.
 - ❖ Chan A, Mak TW. Genomic organization of the T cell receptor. *Cancer Detect Prev*. 1989; 14(2):261-267.

-
- ❖ Chan SH, Waltzinger C, Baron A, Benoist C, Mathis D. Role of coreceptors in positive selection and lineage commitment. *EMBO J.* 1994; 13(19):4482-4489.
 - ❖ Chandrasekaran AN, Krishnamurthy V, Porkodi R, Ramakrishna S, Rajendran CP, Radhakrishna B, Radha Madhavan. Spectrum of clinical and immunological features of systemic rheumatic disorders in a referral hospital in south India: progressive systemic sclerosis. *J Ind Rheum Assoc* 1994; 2:54-60.
 - ❖ Chen X, Murakami T, Oppenheim JJ, Howard OM. Differential response of murine CD4+CD25+ and CD4+CD25- T cells to dexamethasone-induced cell death. *Eur J Immunol.* 2004; 34(3):859–869.
 - ❖ Chen X, Oppenheim JJ. Resolving the identity myth: key markers of functional CD4 (+) FoxP3 (+) regulatory T cells. *Int Immunopharmacol.* 2011; 11(10):1489–1496.
 - ❖ Chong WP, Ip WK, Wong WH-S, Lau CS, Chan TM, Lau YL. Association of interleukin-10 promoter polymorphisms with systemic lupus erythematosus. *Genes Immun.* 2004; 5(6):484–492.
 - ❖ Clements PJ, Hurwitz EL, Wong WK, Seibold JR, Mayes M, White B, Wigley F, Weisman M, Barr W, Moreland L, Medsger TA Jr, Steen VD, Martin RW, Collier D, Weinstein A, Lally E, Varga J, Weiner SR, Andrews B, Abeles M, Furst DE. Skin thickness score as a predictor and correlate of outcome in systemic sclerosis: high-dose versus low-dose penicillamine trial. *Arthritis Rheum.* 2000; 43(11):2445–2454.
 - ❖ Clements PJ, Lachenbruch PA, Ng SC, Simmons M, Sterz M, Furst DE. Skin score. A semiquantitative measure of cutaneous involvement that improves prediction of prognosis in systemic sclerosis. *Arthritis Rheum.* 1990; 33(8):1256–1263.
 - ❖ Coaccioli S, Di-Cato L, Marioli D, Patucchi E, Pizzuti C, Ponteggia M, Puxeddu A. Impaired cutaneous cell-mediated immunity in newly diagnosed rheumatoid arthritis. *Pan Minerva Med.* 2000; 42(4):263-266.
 - ❖ Coffman JD, Zhu J, Roach JM, Bavari S, Ulrich RG, Giardina SL. Production and purification of a recombinant Staphylococcal enterotoxin B vaccine candidate expressed in *Escherichia coli*. *Protein Expr Purif.* 2002; 24(2):302–312.
 - ❖ Cohen N, Kehri H, Berglund B, O’Leary A, Ross G, Seltzer J, Weisel C. Psychoneuroimmunology. *Environ Health Perspect.* 1997; 105(Suppl 2):527-529.

-
- ❖ Cole BC, Knudtson KL, Oliphant A, Sawitzke AD, Pole A, Manohar M, Benson LS, Ahmed E, Atkin CL. The sequence of the *Mycoplasma arthritis* superantigen, MAM: identification of functional domains and comparison with microbial superantigens and plant lectin mitogens. *J Exp Med*. 1996; 183(3):1105–1110.
 - ❖ Colosio C, Birindelli S, Corsini E, Galli CL, Maroni M. Low level exposure to chemicals and immune system. *Toxicol Appl Pharmacol*. 2005; 207(Suppl 2):320–328.
 - ❖ Committee SfSCotARADaTC. Preliminary criteria for the classification of systemic sclerosis (scleroderma). *Arthritis Rheum*. 1980; 23(5):581-590.
 - ❖ Conrad B, Weissmahr RN, Boni J, Arcari R, Schupbach J, Mach B. A human endogenous retroviral superantigen as candidate autoimmune gene in type I diabetes. *Cell*. 1997; 90(2):303–313.
 - ❖ Cooper GS, Dooley MA, Treadwell EL, St Clair EW, Parks CG, Gilkeson GS. Hormonal, environmental, and infectious risk factors for developing systemic lupus erythematosus. *Arthritis Rheum*. 1998; 41(10):1714-1724.
 - ❖ Cooper GS, Miller FW, Pandey JP. The role of genetic factors in autoimmune disease: Implications for environmental research. *Environ Health Perspect*. 1999; 107(Suppl 5):693–700.
 - ❖ Cooper GS, Parks CG, Treadwell EL, St Clair EW, Gilkeson GS, Dooley MA. Occupational risk factors for the development of systemic lupus erythematosus. *J Rheumatol*. 2004; 31(10):1928–1933.
 - ❖ Costenbader KH, Feskanich D, Stampfer MJ, Karlson EW. Reproductive and menopausal factors and risk of systemic lupus erythematosus in women. *Arthritis Rheum*. 2007; 56(4):1251-1262.
 - ❖ Costenbader KH, Karlson EW. Epstein-Barr virus and rheumatoid arthritis: is there a link? *Arthritis Res Ther*. 2006; 8(1):204.
 - ❖ Couper JJ, Kallincos N, Pollard A, Honeyman M, Prager P, Harrison LC, Rischmueller M. Toxic shock syndrome associated with newly diagnosed type I diabetes. *J Paediatr Child Health*. 2000; 36(3):279-282.
 - ❖ Crabtree GR. Contingent genetic regulatory events in T lymphocyte activation. *Science*. 1989; 243(4889):355–361.

-
- ❖ Crilly A, Hamilton J, Clark CJ, Jardine A, Madhok R. Analysis of the 5' flanking region of the interleukin 10 gene in patients with systemic sclerosis. *Rheumatology (Oxford)*. 2003; 42(11):1295–1298.
 - ❖ Crilly A, Hamilton J, Clark CJ, Jardine A, Madhok R. Analysis of transforming growth factor β 1 gene polymorphisms in patients with systemic sclerosis. *Ann Rheum Dis*. 2002; 61(8):678–681.
 - ❖ Criscuoli V, Rizzuto MR, Cottone M. Cytomegalovirus and inflammatory bowel disease: is there a link? *World J Gastroenterol*. 2006; 12(30):4813–4818.
 - ❖ Crispin JC, Martinez A, Alcocer-Varela J. Quantification of regulatory T cells in patients with systemic lupus erythematosus. *J Autoimmun*. 2003; 21(3):273-276.
 - ❖ Croft M. Activation of naïve, memory and effector T cells. *Curr Opin Immunol*. 1994; 6(3):431-437.
 - ❖ Crow MK. Collaboration, Genetic Associations, and Lupus Erythematosus. *N Engl J Med*. 2008; 358(9):956–961.
 - ❖ Cullen MR. Workers with multiple chemical sensitivities: an overview. *Occup Med*. 1987; 2(4):655-661.
 - ❖ D'Alfonso S, Rampi M, Bocchio D, Colombo G, Scorza-Smeraldi R, Momigliano-Richiardi P. Systemic lupus erythematosus candidate genes in the Italian population: Evidence for a significant association with interleukin-10. *Arthritis Rheum*. 2000; 43(1):120–128.
 - ❖ D'Auria L, Bonifati C, Mussi A, D'Agosto G, De Simone C, Giacalone B, Ferraro C, Ameglio F. Cytokines in the sera of patients with pemphigus vulgaris: interleukin-6 and tumour necrosis factor-alpha levels are significantly increased as compared to healthy subjects and correlate with disease activity. *Eur Cytokine Netw*. 1997; 8(4):383–387.
 - ❖ D'Cruz D. Autoimmune diseases associated with drugs, chemicals, and environmental factors. *Toxicol Lett*. 2000; 112-113:421–432.
 - ❖ Dalkilic E, Dilek K, Gullulu M, Yavuz M, Karakocy Y, Yurtkuran M. Lymphocyte phenotypes in systemic sclerosis. *Ann Rheum Dis*. 1999; 58(11):719-720.
 - ❖ Daly AK. Molecular basis of polymorphic drug metabolism. *Mol Med*. 1995; 73(11):539–553.

-
- ❖ Daly MJ, Rioux JD, Schaffner SF, Hudson TJ, Lander ES. High-resolution haplotype structure in the human genome. *Nature Genet.* 2001; 29(2):229–232.
 - ❖ Danchenko N, Satia JA, Anthony MS. Epidemiology of systemic lupus erythematosus: a comparison of worldwide disease burden. *Lupus.* 2006; 15:308-318.
 - ❖ Danis VA, Millington M, Hyland V, Lawford R, Huang Q, Grennan D. Increased frequency of the uncommon allele of a tumor necrosis factor alpha gene polymorphism in rheumatoid arthritis and systemic lupus erythematosus. *Dis Markers.* 1995; 12(2):127–133.
 - ❖ Davis MM, Boniface JJ, Reich Z, Lyons D, Hamp J, Arden B, Chien Y. Ligand recognition by $\alpha\beta$ T cell receptors. *Annu Rev Immunol.* 1998; 16:523-544.
 - ❖ Dawes R, Petrova S, Liu Z, Wraith D, Beverley PC, Tchilian EZ. Combinations of CD45 isoforms are crucial for immune function and disease. *J Immunol.* 2006; 176(6):3417–3425.
 - ❖ De Waal Malefyt R, Abrams J, Bennett B, Figdor CG, de Vries JE. Interleukin 10 (IL-10) inhibits cytokine synthesis by human monocytes: an auto-regulatory role of IL-10 produced by monocytes. *J Exp Med.* 1991a; 174(5):1209-1220.
 - ❖ De Waal Malefyt R, Haanen J, Spits H, Roncarolo MG, te Velde A, Figdor C, Johnson K, Kastelein R, Yssel H, de Vries JE. Interleukin 10 (IL-10) and viral IL-10 strongly reduce antigen-specific human T cell proliferation by diminishing the antigen-presenting capacity of monocytes via downregulation of class II major histocompatibility complex expression. *J Exp Med.* 1991b; 174(4):915-924.
 - ❖ Deapen D, Escalante A, Weinrib L, Horwitz D, Bachman B, Roy- Burman P, Walker A, Mack TM. A revised estimate of twin concordance in systemic lupus erythematosus. *Arthritis Rheum.* 1992; 35(3):311–318.
 - ❖ Degiannis D, Seibold JR, Czarnecki M, Raskova J, Raska K. Soluble Interleukin-2 receptors in patients with systemic sclerosis. *Arthritis Rheum.* 1990; 33(3):375-380.
 - ❖ Dencker L, Hassoun E, d'Argy R, Alm G. Fetal thymus organ culture as an in vitro model for the toxicity of 2,3,7,8- tetrachlorodibenzo-p-dioxin and its congeners. *Mol Pharmacol.* 1985; 27(1):133–140.
 - ❖ Denton CP, Black CM, Abraham DJ. Mechanisms and consequences of fibrosis in systemic sclerosis. *Nat Clin Pract Rheumatol.* 2006; 2(3):134–144.

-
- ❖ Desai MM, Shah DM. Rheumatic and collagen diseases: A review of scientific papers published during 25 years 1944-69. *J Assoc Physicians India*. 1970; 18(4):409-410.
 - ❖ Dick SE, Werth VP. Pemphigus: a treatment update. *Autoimmunity*. 2006; 39(7):591-599.
 - ❖ Ding L, Linsley PS, Huang LY, Germain RN, Shevach EM. IL-10 inhibits macrophage co-stimulatory activity by selectively inhibiting the up-regulation of B7 expression. *J Immunol*. 1993; 151(3):1224-1234.
 - ❖ Dinges MM, Orwin PM, Schlievert PM. Exotoxins of *Staphylococcus aureus*. *Clin Microbiol Rev*. 2000; 13(1):16-34.
 - ❖ Diot E, Lesire V, Guilmot JL, Metzger MD, Pilore R, Rogier S, Stadler M, Diot P, Lemarie E, Lasfargues G. Systemic sclerosis and occupational risk factors: a case-control study. *Occup Environ Med*. 2002; 59(8):545-549.
 - ❖ Dooley MA, Hogan SL. Environmental epidemiology and risk factors for autoimmune disease. *Curr Opin Rheumatol*. 2003; 15(2):99-103.
 - ❖ Drappa J, Vaishnav AK, Sullivan KE, Chu JL, Elkon KB. Fas gene mutations in the Canale-Smith syndrome, an inherited lymphoproliferative disorder associated with autoimmunity. *N Engl J Med*. 1996; 335(22):1643-1649.
 - ❖ Dudda JC, Perdue N, Bachtanian E, Campbell DJ. Foxp3+ regulatory T cells maintain immune homeostasis in the skin. *J Exp Med*. 2008; 205(7):1559-1565.
 - ❖ Ellmeier W, Sawada S, Littman DER. The regulation of CD4 and CD8 coreceptor gene expression during T-cell development. *Annu Rev Immunol*. 1999; 17:523-524.
 - ❖ Encinas JA, Kuchroo VK. Mapping and identification of autoimmunity genes. *Curr Opin Immunol*. 2000; 12(6):691-697.
 - ❖ Erb KJ, Ruger B, von Brevern M, Ryffel B, Schimpl A, Rivett K. Constitutive expression of interleukin (IL)-4 in vivo causes autoimmune- type disorders in mice. *J Exp Med*. 1997; 185(2):329-339.
 - ❖ Eskdale J, Keijsers V, Huizinga T, Gallagher G. Microsatellite alleles and single nucleotide polymorphisms (SNP) combine to form four major haplotype families at the human interleukin-10 (IL-10) locus. *Genes Immun*. 1999; 1(2):151-155.
 - ❖ Esquivel-Sentíes MS, Barrera I, Ortega A, Vega L. Organophosphorous pesticide metabolite (DEDTP) induces changes in the activation status of human lymphocytes

- by modulating IL-2 receptor signal transduction pathway. *Toxicol Appl Pharmacol.* 2010; 248(2):122–133.
- ❖ Evans AS. E.B. virus antibody in systemic lupus erythematosus. *Lancet.* 1971; 1: 1023–1024.
 - ❖ Fadok VA, Bratton DL, Konowal A, Freed PW, Westcott JY, Henson PM. Macrophages that have ingested apoptotic cells in vitro inhibit proinflammatory cytokine production through autocrine/paracrine mechanisms involving TGF-beta, PGE2, and PAF. *J Clin Invest.* 1998; 101(4):890-898.
 - ❖ Fae KC, Oshiro SE, Toubert A, Charron D, Kalil J, Guilherme L. How an autoimmune reaction triggered by molecular mimicry between streptococcal M protein and cardiac tissue proteins leads to heart lesions in rheumatic heart disease. *J Autoimmun.* 2005; 24(2):101-109.
 - ❖ Fahlen L, Read S, Gorelik L, Hurst SD, Coffman RL, Flavell RA, Powrie F. T cells that cannot respond to TGF-beta escape control by CD4⁺CD25⁺ regulatory T cells. *J Exp Med.* 2005; 201(5):737–746.
 - ❖ Fairweather D, Afanasyeva M, Rose NR. Cellular immunity: a role for cytokines. In: Doria A, Pauletto P, editors. *Handbook of systemic autoimmune diseases, Vol I.* Amsterdam: Elsevier, 2004; p. 3-17.
 - ❖ Fairweather D, Kaya Z, Shellam GR, Lawson CM, Rose NR. From infection to autoimmunity. *J Autoimmun.* 2001; 16(3):175-186.
 - ❖ Famularo G, Procopio A, Giacomelli R, Danese C, Sacchetti S, Perego MA, Santoni A, Tonietti G. Soluble interleukin-2 receptor, interleukin-2 and interleukin-4 in sera and supernatants from patients with progressive systemic sclerosis. *Clin Exp Immunol.* 1990; 81(3):368-372.
 - ❖ Farber DL. Cutting edge commentary: differential TCR signaling and the generation of memory T cells. *J Immunol.* 1998; 160(2):535–539.
 - ❖ Fassihi H, Wong T, Wessagowit V, McGrath JA, Mellerio JE. Target proteins in inherited and acquired blistering skin disorders. *Clin Exp Dermatol.* 2006; 31(2):252–259.
 - ❖ Feldman M. TNF blockade in rheumatoid arthritis. *Prog AAI, CIS, AAI Meeting, California.* 1997.

- ❖ Feliciani C, Toto P, Amerio P, Pour SM, Coscione G, Shivji G, Wang B, Sauder DN. In vitro and in vivo expression of interleukin-1 α and tumor necrosis factor- α mRNA in pemphigus vulgaris: interleukin-1 α and tumor necrosis factor- α are involved in acantholysis. *J Invest Dermatol.* 2000; 114(1):71–77.
- ❖ Ferri C, Zignego AL. Relation between infection and autoimmunity in mixed cryoglobulinemia. *Curr Opin Rheumatol.* 2000; 12(1):53–60.
- ❖ Ferry T, Thomas D, Perpoint T, Lina G, Monneret G, Mohammedi I, Chidiac C, Peyramond D, Vandenesch F, Etienne J. Analysis of superantigenic toxin V β T-cell signatures produced during cases of staphylococcal toxic shock syndrome and septic shock. *Clin Microbiol Infect.* 2008; 14(6):546–554.
- ❖ Fey PD, Said-Salim B, Rupp ME, Hinrichs SH, Boxrud DJ, Davis CC, Kreiswirth BN, Schlievert PM. Comparative molecular analysis of community- or hospital-acquired methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother.* 2003; 47(1):196–203.
- ❖ Filippi C, Herrath MV. How viral infections affect the autoimmune process leading to type 1 diabetes. *Cell Immunol.* 2005; 233(2):125–132.
- ❖ Fine JS, Gasiewicz TA, Fiore NC, Silverstone AE. Pro-thymocyte activity is reduced by perinatal 2, 3, 7, 8- tetrachlorodibenzo-p-dioxin exposure. *J Pharmacol Exp Ther.* 1990; 255(1):128–132.
- ❖ Fiorentino DF, Zlotnik A, Mosmann TR, Howard M, O'Garra A. IL-10 inhibits cytokine production by activated macrophages. *J Immunol.* 1991a; 147(11):3815-3822.
- ❖ Fiorentino DF, Zlotnik A, Vieira P, Mosmann TR, Howard M, Moore KW, O'Garra A. IL-10 acts on antigen-presenting cell to inhibit cytokine production by Th1 cells. *J Immunol.* 1991b; 146(10):3444-3451.
- ❖ Fishman D, Faulds G, Jeffery R, Mohamed-Ali V, Yudkin JS, Humphries S, Woo P. The effect of novel polymorphisms in the interleukin-6 (IL-6) gene on IL-6 transcription and plasma IL-6 levels, and an association with systemic-onset juvenile chronic arthritis. *J Clin Invest.* 1998; 102(7):1369–1376.
- ❖ Fleming A, Crown JM, Corbett M. Early rheumatoid arthritis disease. *Ann Rheum Dis.* 1976; 35:357-363.

- ❖ Florquin S, Amraoui Z, Goldman M. Persistent production of TH2- type cytokines and polyclonal B cell activation after chronic administration of staphylococcal enterotoxin B in mice. *J. Autoimmun.* 1996; 9(5):609-615.
- ❖ Fontenot JD, Gavin MA, Rudensky AY. Foxp3 programs the development and function of CD4+CD25+ regulatory T cells. *Nature Immunol.* 2003; 4:330–336.
- ❖ Fontenot JD, Gavin MA, Rudensky AY. Foxp3 programs the development and function of CD4+CD25+ regulatory T cells. *Nat Immunol.* 2003; 4(4):330-336.
- ❖ Fraser JD, Straus D, Weiss A. Signal transduction events leading to T-cell lymphokine gene expression. *Immunol Today.* 1993; 14(7):357–362.
- ❖ Fraser PA, Ding WZ, Mohseni M, Treadwell EL, Dooley MA, St Clair EW, Gilkeson GS, Cooper GS. Glutathione S-transferase M null homozygosity and risk of systemic lupus erythematosus associated with sun exposure: a possible gene–environmental interaction for autoimmunity. *J Rheumatol.* 2003; 30(2):276–282.
- ❖ Friedman SM, Posnett DN, Tumang JR, Cole BC, Crow MK. A potential role for microbial superantigens in the pathogenesis of systemic autoimmune disease. *Arthritis Rheum.* 1991; 34(4):468-480.
- ❖ Friedman SM, Tumang JR, Crow MK. Microbial superantigens as etiopathogenic agents in autoimmunity. *Rheum Dis Clin North Am.* 1993; 19(1):207–222.
- ❖ Fritsch RD, Shen X, Illei GG, Yarboro CH, Prussin C, Hathcock KS, Hodes RJ, Lipsky PE. Abnormal differentiation of memory T-cells in systemic lupus erythematosus. *Arthritis Rheum.* 2006; 54(7):2184–2197.
- ❖ Fuleihan R, Mourad W, Geha RS, Chatila T. Engagement of MHC-class II molecules by staphylococcal exotoxins delivers a co-mitogenic signal to human B cells. *J Immunol.* 1991; 146(5):1661-1666.
- ❖ Gabriel SB, Schaffner SF, Nguyen H, Moore JM, Roy J, Blumenstiel B, Higgins J, DeFelice M, Lochner A, Faggart M, Liu-Cordero SN, Rotimi C, Adeyemo A, Cooper R, Ward R, Lander ES, Daly MJ, Altshuler D. The structure of haplotype blocks in the human genome. *Science.* 2002; 296(5576):2225–2229.
- ❖ Gale CA, Bendel CM, McClellan M, Hauser M, Becker JM, Berman J, Hostetter MK. Linkage of adhesion, filamentous growth, and virulence in *Candida albicans* to a single gene, INT1. *Science.* 1998; 279(5355):1355–1358.

-
- ❖ Garcia KC, Teyton L, Wilson IA. Structural basis of T cell recognition. *Annu Rev Immunol.* 1999; 17:369-397.
 - ❖ Gatenby PA, Kotzin BL, Kansa GS, Engleman EG. Immunoglobulin secretion in the human autologous mixed leukocyte reaction: definition of a suppressor-amplifier circuit using monoclonal antibodies. *J Exp Med.* 1982; 156(1):55-67.
 - ❖ Gazzinelli RT, Oswald IP, James SL, Sher A. IL-10 inhibits parasite killing and nitrogen oxide production by IFN γ - activated macrophages. *J Immunol.* 1992; 148:1792–1796.
 - ❖ Gerli R, Nocentini G, Alunno A, Bocci EB, Bianchini R, Bistoni O, Riccardi C. Identification of regulatory T cells in systemic lupus erythematosus. *Autoimmun Rev.* 2009; 8(5):426–430.
 - ❖ Germain RN. T-cell development and the CD4-CD8 lineage decision. *Nature.* 2002; 2(5):309–322.
 - ❖ Giacomelli R, Cipriani P, Danese C, Pizzuto F, Lattanzio R, Parzanese I, Passacantando A, Perego MA, Tonietti G. Peripheral blood mononuclear cells of patients with systemic sclerosis produce increased amounts of interleukin 6, but not transforming growth factor beta 1. *J Rheumatol.* 1996; 23(2):291-296.
 - ❖ Giacomelli R, Cipriani P, Lattanzio R, Di FM, Locanto M, Parzanese I, Passacantando A, Ciocci A, Tonietti G. Circulating levels of soluble CD30 are increased in patients with systemic sclerosis (SSc) and correlate with serological and clinical features of the disease. *Clin Exp Immunol.* 1997; 108(1):42–46.
 - ❖ Gibson AW, Edberg JC, Wu J, Westendorp RG, Huizinga TW, Kimberly RP. Novel single nucleotide polymorphisms in the distal IL-10 promoter affect IL-10 production and enhance the risk of systemic lupus erythematosus. *J Immunol.* 2001; 166(6):3915-3922.
 - ❖ Gilbert KM, Griffin JM, Pumford NR. Trichloroethylene activates CD4⁺ T cells: potential role in an autoimmune response. *Drug Metab Rev.* 1999; 31(4):901-916.
 - ❖ Ginn LR, Lin JP, Plotz PH, Bale SJ, Wilder RL, Mbauya A, Miller FW. Familial autoimmunity in pedigrees of idiopathic inflammatory myopathy patients suggests common genetic risk factors for many autoimmune diseases. *Arthritis Rheum.* 1998; 41(3):400-405.

-
- ❖ Giurgea R, Witterberger C, Frecus G, Manciulea S, Borsa M, Coprean D, Ilyes S. Effects of some organochlorine pesticides on the immunological reactivity of white rats. *Archiv Fur Exp Vet.* 1978; 32(5):769–774.
 - ❖ Glick B. Antibody-mediated immunity in the presence of mirex and DDT. *Poultry Sci.* 1974; 53(4):1476–1485.
 - ❖ Go NF, Castle BE, Barret R, Kastelein R, Dang W, Mosmann TR, Moore KW, Howard M. Interleukin-10, a novel B cell stimulatory factor: unresponsiveness of X-chromosome-linked immunodeficiency B cells. *J Exp Med.* 1990; 172(6):1625–1631.
 - ❖ Goddard GZ, Shoenfeld Y. Infections and SLE. *Autoimmunity.* 2005; 38(7):473–485.
 - ❖ Goodnow CC, Sprent J, Fazekas de St Groth B, Vinuesa CG. Cellular and genetic mechanisms of self tolerance and autoimmunity. *Nature.* 2005; 435:590-597.
 - ❖ Goodnow CC. Balancing immunity and tolerance: deleting and tuning lymphocyte repertoires. *Proc Nat Acad Sci USA.* 1996; 93(6):2264-2271.
 - ❖ Gorla R, Airo P, Malagoli A, Carella G, Prati E, Brugnoli D, Franceschini F, Cattaneo R. CD4⁺ and CD8⁺ subsets: Naive and memory cells in the peripheral blood of patients with systemic sclerosis. *Clin Rheumatol.* 1994; 13(1):83-87.
 - ❖ Grainger DJ, Heathcote K, Chiano M, Snieder H, Kemp PR, Metcalfe JC, Carter ND, Spector TD. Genetic control of the circulating concentration of transforming growth factor type beta 1. *Hum Mol Genet.* 1999; 8(1):93–97.
 - ❖ Grando SA. Autoimmunity to keratinocyte acetylcholine receptors in pemphigus. *Dermatology.* 2000; 201(4):290-295.
 - ❖ Granel B, Chevillard C, Allanore Y, Arnaud V, Cabantous S, Marquet S, Weiller PJ, Durand JM, Harlé JR, Grange C, Frances Y, Berbis P, Gaudart J, de Micco P, Kahan A, Dessein A. Evaluation of interleukin 13 polymorphisms in systemic sclerosis. *Immunogenetics.* 2006; 58(8):693–699.
 - ❖ Gregersen PK. Genetic analysis of rheumatic diseases. In: Kelley WN, Harris ED Jr, Ruddy S, Sledge CB, editors. *Textbook of rheumatology*, 5th ed. Philadelphia: WB Saunders, 1997; p. 209-227. v.1.
 - ❖ Grewal IS. Overview of TNF superfamily: a chest full of potential therapeutic targets. *Adv Exp Med Biol.* 2009; 647:1-7.

-
- ❖ Griem P, Wulferink M, Sachs B, González JB, Gleichmann E. Allergic and autoimmune reactions to xenobiotics: how do they arise? *Immunol Today*. 1998; 19(3):133–141.
 - ❖ Grondal G, Gunnarsson I, Ronnelid J, Rogberg S, Klareskog L, Lundberg I. Cytokine production, serum levels and disease activity in systemic lupus erythematosus. *Clin Exp Rheumatol*. 2000; 18(5):565–570.
 - ❖ Grossman C, Dovrish Z, Shoenfeld Y, Amital H. Do infections facilitate the emergence of systemic sclerosis? *Autoimmun Rev*. 2011; 10(5):244-247.
 - ❖ Grossman Z, Paul WE. Self-tolerance: context-dependent tuning of T cell antigen recognition. *Semin Immunol*. 2000; 12(3):197–203.
 - ❖ Grunwald M, Katz I, Friedman-Birnbaum R. Association of pemphigus vulgaris and herpes simplex virus infection. *Int J Dermatol*. 1986; 25(6):392-393.
 - ❖ Guarneri F, Guarneri C, Benvenga S. Helicobacter pylori and autoimmune pancreatitis: Role of carbonic anhydrase via molecular mimicry? *J Cell Mol Med*. 2005; 9(3):741-744.
 - ❖ Guarnizo-Zuccardi P, Lopez Y, Giraldo M, Garcia N, Rodriguez L, Ramirez L, Uribe O, Garcia L, Vasquez G. Cytokine gene polymorphisms in Colombian patients with systemic lupus erythematosus. *Tissue Antigens*. 2007; 70(5):376–382.
 - ❖ Guiducci R et al. Infection and systemic sclerosis. In: Shoenfeld Y, Rose NR, editors. *Infection and Autoimmunity*. Amsterdam, San Diego, United Kingdom: Elsevier, 2004; p. 613–622
 - ❖ Guilherme L, Faé KC, Oshiro SE, Tanaka AC, Pomerantzeff PM, Kalil J. T cell response in rheumatic fever: crossreactivity between streptococcal M protein peptides and heart tissue proteins. *Curr Protein Pept Sci*. 2007; 8(1):39-44.
 - ❖ Guilherme L, Kalil J. Rheumatic fever and rheumatic heart disease: cellular mechanisms leading autoimmune reactivity and disease. *J Clin Immunol*. 2010; 30(1):17-23.
 - ❖ Gustafsson R, Totterman TH, Klareskog L, Hällgren R. Increase in activated T-cells and reduction in suppressor inducer T-cells in systemic sclerosis. *Ann Rheum Dis*. 1990; 49(1):40-45.

- ❖ Haahr S, Hollsberg P. Multiple sclerosis is linked to Epstein-Barr virus infection. *Rev Med Virol*. 2006; 16(5):297–310.
- ❖ Hagiwara E, Gourley MF, Lee S, Klinman DK. Disease severity in patients with SLE correlates with an increased ratio of IL-10: IFN γ -secreting cells in the peripheral blood. *Arthritis Rheum*. 1996; 39(3):379–385.
- ❖ Hajeer AH, Lazarus M, Turner D, Mageed RA, Vencovsky J, Sinnott P, Hutchinson IV, Ollier WE. IL-10 gene promoter polymorphism in rheumatoid arthritis. *Scand J Rheumatol*. 1998; 27(2):142–145.
- ❖ Hajeer AH, Worthington J, Silman AJ, Ollier WER. Association of tumor necrosis factor microsatellite polymorphisms with HLA-DRB1*04-bearing haplotypes in rheumatoid arthritis patients. *Arthritis Rheum*. 1996; 39(7):1109–1114.
- ❖ Hall SK, Perregaux DG, Gabel CA, Woodworth T, Durham LK, Huizinga TW, Breedveld FC, Seymour AB. Correlation of polymorphic variation in the promoter region of the interleukin-1 β gene with secretion of interleukin-1 β protein. *Arthritis Rheum*. 2004; 50(6):1976–1983.
- ❖ Hamad AR, Herman A, Marrack P, Kappler JW. Monoclonal antibodies defining functional sites on the toxin superantigen staphylococcal enterotoxin B. *J Exp Med*. 1994; 180:615–621.
- ❖ Hamamdžić D, Harley RA, Hazen-Martin D, LeRoy EC. MCMV induces neointima in IFN-gamma R^{-/-} mice: intimal cell apoptosis and persistent proliferation of myofibroblasts. *BMC Musculoskelet Disord*. 2001; 2:3.
- ❖ Harley JB, Harley IT, Guthridge JM, James JA. The curiously suspicious: a role for Epstein-Barr virus in lupus. *Lupus*. 2006; 15(11):768–777.
- ❖ Harrison's Principles of Internal Medicine, 18th ed. Chapter 319. Systemic Lupus Erythematosus [Accessed on 5th August, 2012]. Available from <http://www.accessmedicine.com/resourceTOC.aspx?resourceID=4#18top>.
- ❖ Hart HH, Grigor RR, Caughey DE. Ethnic difference in the prevalence of systemic lupus erythematosus. *Ann Rheum Dis*. 1983; 429(5):529-532.
- ❖ Hasegawa M, Fujimoto M, Kikuchi K, Takehara K. Elevated serum tumor necrosis factor-alpha levels in patients with systemic sclerosis: association with pulmonary fibrosis. *J Rheumatol*. 1997; 24(4):663-665.

-
- ❖ He B, Xu C, Yang B, Landtblom AM, Fredrikson S, Hillert J. Linkage and association analysis of genes encoding cytokines and myelin proteins in multiple sclerosis. *J Neuroimmunol.* 1998; 86(1):13–19.
 - ❖ He D, Wu L, Kim HK, Li H, Elmetts CA, Xu H. CD8 + IL-17-producing T cells are important in effector functions for the elicitation of contact hypersensitivity responses. *J Immunol.* 2006; 177(10):6852–6858.
 - ❖ Healy JI, Goodnow CC. Positive versus negative signaling by lymphocyte antigen receptors. *Annu Rev Immunol.* 1998; 16:645–670.
 - ❖ Hendel L, Hage E, Hendel J, Stentoft P. Omeprazole in the long-term treatment of severe gastro-oesophageal reflux disease in patients with systemic sclerosis. *Aliment Pharmacol Ther.* 1992; 6(5):565–577.
 - ❖ Henderson RD, Bain CJ, Pender MP. The occurrence of autoimmune diseases in patients with multiple sclerosis and their families. *J Clin Neurosci.* 2000; 7(5):434-437.
 - ❖ Hermiston ML, Xu Z, Weiss A. CD45: a critical regulator of signaling thresholds in immune cells. *Annu Rev Immunol.* 2003; 21:107–137.
 - ❖ Hershey GKH, Friedrich MF, Esswein LA, Thomas ML, Chatila TA. The association of atopy with a gain-of-function mutation in the α subunit of the interleukin-4 receptor. *N Engl J Med.* 1997; 337(24):1720–1725.
 - ❖ Hertl M, Veldman C. Pemphigus: paradigm of autoantibody-mediated autoimmunity. *Skin Pharmacol Appl Skin Physiol.* 2001; 14(6):408–418.
 - ❖ Hertl M. Humoral and cellular autoimmunity in autoimmune bullous skin disorders. *Int Arch Allergy Immunol.* 2000; 122(2):91–100.
 - ❖ Hess EV. Are there environmental forms of systemic autoimmune diseases? *Environ Health Perspectives.* 1999; 107(suppl 5):709-711.
 - ❖ Heufelder AE. T-cell restriction in thyroid eye disease. *Thyroid.* 1998; 8(5):419-422.
 - ❖ Hochberg MC. The incidence of systemic lupus erythematosus in Baltimore, Maryland, 1970-1977. *Arthritis Rheum.* 1985; 28(1):80-86.
 - ❖ Hoffmann SC, Stanley EM, Darrin Cox E, Craighead N, DiMercurio BS, Koziol DE, Harlan DM, Kirk AD, Blair PJ. Association of cytokine polymorphic inheritance and

- in vitro cytokine production in anti-CD3/CD28-stimulated peripheral blood lymphocytes. *Transplantation*. 2001; 72(8):1444–1450.
- ❖ Hohlfeld R, Toyka KV, Heininger K, Grosse-Wilde H, Kalies I. Autoimmune human T lymphocytes specific for acetylcholine receptor. *Nature*. 1984; 310(5974):244–246.
 - ❖ Holladay SD, Lindstorm P, Blaylock BL, Comment CE, Germolec DR, Heindell JJ, Luster MI. Perinatal thymocyte antigen expression and postnatal immune development altered by gestational exposure to tetrachlorodibenzo-p-dioxin (TCDD). *Teratology*. 1991; 44(4):385–393.
 - ❖ Hollegaard MV, Bidwell JL. Cytokine gene polymorphisms in human disease: on-line databases, Supplement 3. *Genes Immun*. 2006; 7(4):269–276.
 - ❖ Holsapple MP. Autoimmunity by pesticides: A critical review of the state of the science. *Toxicol Lett*. 2002; 127(1-3):101–109.
 - ❖ Holweg CT, Baan CC, Niesters HGM, Vantrimpont PJMJ, Mulder PGH, Maat APWM, Weimar W, Balk AHMM. TGF-beta1 gene polymorphisms in patients with end stage heart failure. *J Heart Lung Transplant*. 2001; 20(9):979–984.
 - ❖ Hom G, Graham RR, Modrek B, Taylor KE, Ortmann W, Garnier S, Lee AT, Chung SA, Ferreira RC, Pant PV, Ballinger DG, Kosoy R, Demirci FY, Kamboh MI, Kao AH, Tian C, Gunnarsson I, Bengtsson AA, Rantapää-Dahlqvist S, Petri M, Manzi S, Seldin MF, Rönnblom L, Syvänen AC, Criswell LA, Gregersen PK, Behrens TW Association of Systemic Lupus Erythematosus with C8orf13–BLK and ITGAM–ITGAX. *N Engl J Med*. 2008; 358(9):900–909.
 - ❖ Hori S, Nomura T, Sakaguchi S. Control of regulatory T cell development by the transcription factor Foxp3. *Science*. 2003; 299(5609):1057–1061.
 - ❖ Horiuchi T, Washio M, Kiyohara C, Tsukamoto H, Tada Y, Asami T, Ide S, Kobashi G, Takahashi H, the Kyushu Sapporo SLE Study Group. Combination of TNF-RII, CYP1A1 and GSTM1 polymorphisms and the risk of Japanese SLE: Findings from the KYSS study. *Rheumatology*. 2009; 48(9):1045–1049.
 - ❖ Horwitz DA. Regulatory T cells in systemic lupus erythematosus: past, present and future. *Arthritis Res Ther*. 2008; 10(6):227.
 - ❖ Hoyles RK, Ellis RW, Wellsbury J, Lees B, Newlands P, Goh NS, Roberts C, Desai S, Herrick AL, McHugh NJ, Foley NM, Pearson SB, Emery P, Veale DJ, Denton CP,

- Wells AU, Black CM, du Bois RM. A multicenter, prospective, randomized, double-blind, placebo-controlled trial of corticosteroids and intravenous cyclophosphamide followed by oral azathioprine for the treatment of pulmonary fibrosis in scleroderma. *Arthritis Rheum.* 2006; 54(12):3962–3970.
- ❖ Hsieh AH, Zhou YJ, Liang CT, Chang M, Wang SL. Fragment of tegument protein pp65 of human cytomegalovirus induces autoantibodies in BALB/c mice. *Arthritis Res Ther.* 2011; 13:R162.
 - ❖ Huang CM, Wu MC, Wu JY, Tsai FJ. Lack of association of interleukin-1 β gene polymorphisms in Chinese patients with systemic lupus erythematosus. *Rheumatol Int.* 2002; 21(5):173-175.
 - ❖ Huber H C, Huber W, Ritter U. Simple in vitro test systems for determining the toxicity of environmental chemicals: Microcultures of human lymphocytes and monoxenic ciliate cultures. *Zentral Hyg Umweltmed.* 1990; 189(6):511–526.
 - ❖ Hueser G. Diagnostic markers in clinical immunotoxicology and neurotoxicology. *J Occup Med Toxicol.* 1992; 1:5-9.
 - ❖ Huggins ML, Todd I, Cavers MA, Pavuluri SR, Tighe PJ, Powell RJ. Antibodies from systemic lupus erythematosus (SLE) sera define differential release of autoantigens from cell lines undergoing apoptosis. *Clin Exp Immunol.* 1999; 118(2):322-328.
 - ❖ Hughes MM, Yassai M, Sedy JR, Wehrly TD, Huang CY, Kanagawa O, Gorski J, Sleckman BP. T cell receptor CDR3 loop length repertoire is determined primarily by features of the V(D)J recombination reaction. *Eur J Immunol.* 2003; 33(6):1568-1575.
 - ❖ Huizinga TW, Westendorp RG, Bollen EL, Keijsers V, Brinkman BM, Langermans JA, Breedveld FC, Verweij CL, van de Gaer L, Dams L, Crusius JB, García-Gonzalez A, van Oosten BW, Polman CH, Peña AS. TNF-alpha promoter polymorphisms, production, and susceptibility to multiple sclerosis in different groups of patients. *J Neuroimmunol.* 1997; 72(2):149–153.
 - ❖ Hulkkonen J, Pertovaara M, Anttonen J, Pasternack A, Hurme M. IL-1 gene family haplotypes and Raynaud's phenomenon in primary Sjogren's syndrome. *Rheumatology.* 2002; 41(10):1206–1208.
 - ❖ Hurme M, Lahdenpohja N, Santtila S. Gene polymorphisms of interleukins 1 and 10 in infectious and autoimmune diseases. *Ann Med.* 1998; 30(5):469-473.

-
- ❖ Hutyrova B, Lukac J, Bosak V, Buc M, du Bois R, Petrek M. Interleukin 1_ single nucleotide polymorphism associated with systemic sclerosis. *J Rheumatol.* 2004; 31(1):81–84.
 - ❖ Hwang IR, Kodama T, Kikuchi S, Sakai K, Peterson LE, Graham DY, Yamaoka Y. Effect of interleukin 1 polymorphisms on gastric mucosal interleukin 1beta production in *Helicobacter pylori* infection. *Gastroenterology.* 2002; 123(6):1793-1803.
 - ❖ Iacoviello L, Di Castelnuovo A, Gattone M, Pezzini A, Assanelli D, Lorenzet R, Del Zotto E, Colombo M, Napoleone E, Amore C, D'Orazio A, Padovani A, de Gaetano G, Giannuzzi P, Donati MB; IGIGI Investigators. Polymorphisms of the interleukin-1beta gene affect the risk of myocardial infarction and ischemic stroke at young age and the response of mononuclear cells to stimulation in vitro. *Arterioscler Thromb Vasc Biol* 2005, 25(1):222-227.
 - ❖ Ignatowicz L, Kappler J, Marrack P. The repertoire of T cells shaped by a single MHC/peptide ligand. *Cell.* 1996; 84(4):521–529.
 - ❖ International Human Genome Sequencing Consortium. Finishing the euchromatic sequence of the human genome. *Nature.* 2004; 431(7011):931–945.
 - ❖ Isomaki P, Clark JM, Panesar M, Cope AP. Pathways of T-cell activation and terminal differentiation in chronic inflammation. *Curr Drug Targets Inflamm Allergy.* 2005; 4(3):287–293.
 - ❖ Jacobson D L, Gange SJ, Rose NR, Graham NM. Epidemiology and estimated population burden of selected autoimmune diseases in the United States. *Clin Immunol Immunopathol.* 1997; 84(3):223–243.
 - ❖ Janeway CA Jr. Approaching the asymptote? Evolution and revolution in immunology. *Cold Spring Harb Symp Quant Biol.* 1989; 54:1–13.
 - ❖ Jardetzky TS, Brown JH, Gorga JC, Stern LJ, Urban RG, Chi YI, Stauffacher C, Strominger JL, Wiley DC. Threedimensional structure of a human class II histocompatibility molecule complexed with superantigen. *Nature.* 1994; 368(6473):711–718.
 - ❖ Jarvinen P, Kaprio J, Makitalo R, Koskenvuo M, Aho K. Systemic lupus erythematosus and related systemic diseases in a nationwide twin cohort: An

- increased prevalence of disease in MZ twins and concordance of disease features. *J Intern Med.* 1992; 231(1):67–72.
- ❖ Javor J, Chmurova N, Parnicka Z, Ferencik S, Grosse-Wilde H, Buc M, Svecova D. TNF- α and IL-10 gene polymorphisms show a weak association with pemphigus vulgaris in the Slovak population. *JEADV.* 2010; 24(1):65–68.
 - ❖ Jeffrey AJ, Kauppi L, Neumann R. Intensely punctate meiotic recombination in the class II region of the major histocompatibility complex. *Nature Genet.* 2001; 29:217–222.
 - ❖ Johnson G, Wu TT. Kabat database and its applications: 30 years after the first variability plot. *Nucleic Acids Res.* 2000; 28(1):214-218.
 - ❖ Johnson HM, Magazine HI. Potent mitogenic activity of staphylococcal enterotoxin A requires induction of interleukin 2. *Int Arch Allergy Appl Immunol.* 1988; 87(1):87–90.
 - ❖ Jorgensen JL, Esser U, Fazekas de St. Groth B, Reay PA, Davis MM. Mapping T-cell receptor-peptide contacts by variant peptide immunization of single-chain transgenics. *Nature.* 1992; 355(6357):224-230.
 - ❖ Kahaleh B. Raynaud's phenomenon and the vascular disease in scleroderma. *Curr Opin Rheumatol.* 1995; 7(6):529-534.
 - ❖ Kalinkovich A, Weisman Z, Greenberg Z, Nahmias J, Eitan S, Stein M, Bentwich ZR. Decreased CD4 and increased CD8 counts with T-cell activation is associated with chronic helminth infection. *Clin Exp Immunol.* 1998; 114(3):414-421.
 - ❖ Kammer GM, Stein RL. T-Lymphocyte immune dysfunction in systemic lupus erythematosus. *J Lab Clin Med.* 1990; 115(3):273–282.
 - ❖ Kantor TV, Frieberg D, Medsger TA Jr, Buckingham RB, Whiteside TL. Cytokine production and serum levels in systemic sclerosis. *Clin Immunol Immunopathol.* 1992; 65(3):278-285.
 - ❖ Kanwar AJ, Ajith AC, Narang T. Pemphigus in North India. *J Cutan Med Surg.* 2006; 10(1):21–25.
 - ❖ Katsumoto TR, Whitfield ML, Connolly MK. The pathogenesis of systemic sclerosis. *Ann Rev Pathol.* 2011; 6:509-537.

-
- ❖ Kerker BJ, Morison WL. The photoaggravated dermatoses. *Semin Dermatol.* 1990; 9(1):70-77.
 - ❖ Keskin DB, Stern JN, Fridkis-Hareli M, Razzaque AA. Cytokine profiles in pemphigus vulgaris patients treated with intravenous immunoglobulins as compared to conventional immunosuppressive therapy. *Cytokine.* 2008; 41(3):315–321.
 - ❖ Khattri R, Cox T, Yasayko SA, Ramsdell F. An essential role for Scurfin in CD4+CD25+ T regulatory cells. *Nature Immunol.* 2003; 4(4):337–342.
 - ❖ Khoa PD, Sugiyama T, Yokochi T. Polymorphism of interleukin-10 promoter and tumor necrosis factor receptor II in Vietnamese patients with systemic lupus erythematosus. *Clin Rheumatol.* 2005; 24(1):11–13.
 - ❖ Kimura A, Naka T, Kishimoto T. IL-6-dependent and -independent pathways in the development of interleukin 17-producing T helper cells. *Proc Natl Acad Sci US.* 2007; 104(29):12099–12104.
 - ❖ Kirchner H, Kruse A, Neustock P, Rink L. Cytokine. In: Kirchner H, editors. *Cytokine und Interferone.* Heidelberg-Berlin-Oxford: Spektrum Akademischer Verlag, 1993; p. 30–34.
 - ❖ Kisand K, Lilic D, Casanova JL, Peterson P, Meager A, Willcox N. Mucocutaneous candidiasis and autoimmunity against cytokines in APECED and thymoma patients: Clinical and pathogenetic implications. *Eur J Immunol.* 2011; 41(6):1517–1527.
 - ❖ Kjer-Nielsen L, Clements CS, Brooks AG, Purcell AW, McCluskey J, Rossjohn J. The 1.5 Å crystal structure of a highly selected antiviral T cell receptor provides evidence for a structural basis of immunodominance. *Structure.* 2002; 10(11):1521-1532.
 - ❖ Klein J, Sato A. The HLA system. Second of two parts. *N Engl J Med.* 2000; 343(11):782-786.
 - ❖ Komai A, Amagai M, Ishii K, Nishikawa T, Chorzelski T, Matsuo I, Hashimoto T. The clinical transition between pemphigus foliaceus and pemphigus vulgaris correlates well with the changes in autoantibody profile assessed by an enzyme-linked immunosorbent assay. *Br J Dermatol.* 2001; 144(6):1177–1182.
 - ❖ Kotb M, Fraser JD, editors. *Superantigens: molecular basis for their role in human diseases.* Washington: ASM Press, 2008; p. 263.

-
- ❖ Kotzin BL, Leung DY, Kappler J, Marrack P. Superantigens and their potential role in human disease. *Adv Immunol.* 1993; 54:99–166.
 - ❖ Kouki T, Sawai Y, Gardine CA, Fisfalen ME, Alegre ML, DeGroot LJ. CTLA-4 gene polymorphism at position 49 in exon 1 reduces the inhibitory function of CTLA-4 and contributes to the pathogenesis of Graves' disease. *J Immunol.* 2000; 165(11): 6606-6611.
 - ❖ Krain LS. Pemphigus. Epidemiologic and survival characteristics of 59 patients, 1955-1973. *Arch Dermatol.* 1974; 110(6):862-865.
 - ❖ Kronenberg M. Self tolerance and autoimmunity. *Cell.* 1991; 65(4):537-542.
 - ❖ Kukreja A, Maclaren NK. Autoimmunity and diabetes. *J Clin Endocrinol Metab.* 1999; 84(12):4371-4378.
 - ❖ Kukreja A, Maclaren NK. Current cases in which epitope mimicry is considered as a component cause of autoimmune disease: immunemediated (type 1) diabetes. *Cell Mol Life Sci.* 2000; 57(4):534-541.
 - ❖ Kumar KA. Incidence of pemphigus in Thrissur district, south India. *Indian J Dermatol Venereol Leprol.* 2008; 74(12):349–351.
 - ❖ Kurreeman FA, Schonkeren JJ, Heijmans BT, Toes RE, Huizinga TW. Transcription of the IL10 gene reveals allele-specific regulation at the mRNA level. *Hum Mol Genet.* 2004; 13(16):1755–1762.
 - ❖ Lahita RG. The connective tissue diseases and the overall influence of gender. *Int J Fertil Menopausal Stud.* 1996; 41(2):156–165.
 - ❖ Lambert J, Schepea S, Jansseas J, Dockx P. Skin lesions as a sign of subacute pentachlorophenol intoxication. *Acta Derm Venerol.* 1986; 66(2):170-172.
 - ❖ Larche M. Determining MHC restriction of T-cell responses. *Methods Mol Med.* 2008; 138:57-72.
 - ❖ Larsen CE, Alper CA. The genetics of HLA associated disease. *Curr Opin Immunol.* 2004; 16(5):660–667.
 - ❖ Larsson EL, Gullberg M, Bandeira A, Coutinho A. Activation and growth requirements for cytotoxic and noncytotoxic T-lymphocytes. *Cell Immunol.* 1984; 89(1):223-231.

- ❖ Laufer TM, DeKoning J, Markowitz JS, Lo D, Glimcher LH. Unopposed positive selection and autoreactivity in mice expressing class II MHC only on thymic cortex. *Nature*. 1996; 383(6595):81–85.
- ❖ Lazarus M, Hajeer AH, Turner D, Sinnott P, Worthington J, Ollier WE, Hutchinson IV. Genetic variation in the interleukin 10 gene promoter and systemic lupus erythematosus. *J Rheumatol*. 1997; 24(12):2314–2317.
- ❖ Lee E, Lendas KA, Chow S, Pirani Y, Gordon D, Dionisio R, Nguyen D, Spizuoco A, Fotino M, Zhang Y, Sinha AA. Disease relevant HLA class II alleles isolated by genotypic, haplotypic, and sequence analysis in North American Caucasians with pemphigus vulgaris. *Hum Immunol*. 2006; 67(1-2):125–139.
- ❖ Lee IH, Li WP, Hisert KB, Ivashkiv LB. Inhibition of IL-2 signaling and signal transducer and activator of transcription (STAT) 5 activation during T-cell receptor-mediated feedback inhibition of T-cell expansion. *J Exp Med*. 1999; 190(9):1263–1274.
- ❖ Lee JH, Wang LC, Lin YT, Yang YH, Lin DT, Chiang BL. Inverse correlation between CD4+ regulatory T-cell population and autoantibody levels in paediatric patients with systemic lupus erythematosus. *Immunology*. 2006; 117(2):280–286.
- ❖ Lee TP, Moscati R, Park BH. Effects of pesticides on human leukocyte functions. *Res Commun Chem Pathol Pharmacol*. 1979; 23(3):597–609.
- ❖ Lehmann PV, Forsthuber T, Miller A, Sercarz EE. Spreading of T-cell autoimmunity to cryptic determinants of an autoantigen. *Nature*. 1992; 358(6382):155–157.
- ❖ Leitenberg D, Boutin Y, Lu DD, Bottomly K. Biochemical association of CD45 with the T cell receptor complex: regulation by CD45 isoform and during T cell activation. *Immunity*. 1999; 10(6):701–711.
- ❖ LeRoy EC, Black C, Fleischmajer R, Jablonska S, Krieg T, Medsger TA Jr, Rowell N, Wollheim F. Scleroderma (systemic sclerosis): classification, subsets and pathogenesis. *J Rheumatol*. 1988; 15(2):202-205.
- ❖ Letterio J, Roberts A. Regulation of immune responses by TGF-beta. *Annu Rev Immunol*. 1998; 16:137–161.
- ❖ Li H, Llera A, Malchiodi EL, Mariuzza RA. The structural basis of T cell activation by superantigens. *Annu Rev Immunol*. 1999; 17:435–466.

- ❖ Li YL, Li HM, Dimasi N, McCormick JK, Martin R, Schuck P, Schlievert PM, Mariuzza RA. Crystal structure of a superantigen bound to the high-affinity, zinc-dependent site on MHC class II. *Immunity*. 2001; 14(1):93–103.
- ❖ Lin MS, Fu CL, Aoki V, Hans-Filho G, Rivitti EA, Moraes JR, Moraes ME, Lazaro AM, Giudice GJ, Stastny P, Diaz LA. Desmoglein-1-specific T lymphocytes from patients with endemic Pemphigus foliaceus (fogo selvagem). *J Clin Invest*. 2000; 105(2):207–213.
- ❖ Lin MS, Liu Z, Drolet BA, Diaz LA. Cutaneous autoimmune diseases. In: Rose NR, Mackays IR, editors. *The Autoimmune Diseases*. San Diego: Academic Press, 1998; p. 545–570.
- ❖ Lin MS, Swartz SJ, Lopez A, Ding X, Fernandez-Vina MA, Stastny P, Fairley JA, Diaz LA. Development and characterization of desmoglein- 3 specific T cells from patients with pemphigus vulgaris. *J Clin Invest*. 1997; 99(1):31–40.
- ❖ Lin SC, Chen KH, Lin CH, Kuo CC, Ling QD, Chan CH. The quantitative analysis of peripheral blood FOXP3-expressing T cells in systemic lupus erythematosus and rheumatoid arthritis patients. *Eur J Clin Invest*. 2007; 37(12):987–996.
- ❖ Lindqvist KJ, Makena WJ, Shaba JD, Nantulya V. Immunofluorescence and electron microscopic studies of kidney biopsies from patients with nephritic syndrome, possibly induced by skin lightening creams containing mercury. *E Afr Med*. 1974; 51:168.
- ❖ Linker-Israeli M, Bakke AC, Kifridou RC, Gendler KS, Gillis S, Horwitz DA. Defective production of IL-1 and IL-2 in patients with systemic lupus erythematosus. *J Immunol*. 1983; 130(6):2651–2655.
- ❖ Liston A, Lesage S, Wilson J, Peltonen L, Goodnow CC. Aire regulates negative selection of organ-specific T cells. *Nature Immunol*. 2003; 4:350–354.
- ❖ Liu MF, Wang CR, Fung LL, Wu CR. Decreased CD4⁺CD25⁺ T cells in peripheral blood of patients with systemic lupus erythematosus. *Scand J Immunol*. 2004; 59(2):198–202.
- ❖ Llorente L, Richaud-Patin Y, Wijdenes J, Alcocer-Varela J, Maillot MC, Durand-Gasselin I, Fourrier BM, Galanaud P, Emilie D. Spontaneous production of IL-10 by

- B-lymphocytes and monocytes in systemic lupus erythematosus. *Eur Cytokine Netw.* 1993; 4(6):421–427.
- ❖ Llorente L, Zou W, Levy Y, Richaud-Patin Y, Wijdenes J, Alcocer- Varela J, Morel-Fourrier B, Brouet JC, Alarcon-Segovia D, Galanaud P, Emilie D. Role of interleukin-10 in the B-lymphocyte hyperactivity and autoantibody production of human systemic lupus erythematosus. *J Exp Med.* 1995; 181(3):839–844.
 - ❖ Lloyd TM, Panush RS. Cell-mediated immunity in rheumatoid arthritis: Impaired lymphocyte responsiveness, humoral immunosuppressants and correlation with clinical status in patients off drug therapy. *J Rheumatol.* 1977; 4(3):231-44.
 - ❖ Lo D, Reilly CR, Burkly LC, DeKoning J, Laufer TM, Glimcher LH. Thymic stromal cell specialization and the T-cell receptor repertoire. *Immunol Res.* 1997; 16(1):3-14.
 - ❖ Lopez-Robles E, Avalos-Dí'az E, Vega-Memije E, Hojyo-Tomoka T, Villalobos R, Fraire S, Domí'guez-Soto L, Herrera-Esparza R. TNF- α and IL-6 are mediators in the blistering process of pemphigus. *Int J Dermatol.* 2001; 40(3):185–188.
 - ❖ Lockshin MD. Sex differences in autoimmune disease. *Lupus.* 2006; 15(11):753-756.
 - ❖ Lord SJ, Rajotte RV, Korbitt GS, Bleackley RC. Granzyme B: a natural born killer. *Immunol Rev.* 2003; 193:31–38.
 - ❖ Lu LF, Rudensky A. Molecular orchestration of differentiation and function of regulatory T cells. *Genes Dev.* 2009; 23(11):1270–1282.
 - ❖ Lu LY, Cheng HH, Sung PK, Yeh JJ, Shiue YL, Chen A. Single-nucleotide polymorphisms of transforming growth factor-beta1 gene in Taiwanese patients with systemic lupus erythematosus. *J Microbiol Immunol Infect.* 2004; 37(3):145–152.
 - ❖ Lunardi C, Bason C, Navone R, Millo E, Damonte G, Corrocher R, Puccetti A. Systemic sclerosis immunoglobulin G autoantibodies bind the human cytomegalovirus late protein UL94 and induce apoptosis in human endothelial cells. *Nat Med.* 2000; 6(10):1183-1186.
 - ❖ Lunardi C, Dolcino M, Peterlana D, Bason C, Navone R, Tamassia N, Beri R, Corrocher R, Puccetti A. Antibodies against human cytomegalovirus in the pathogenesis of systemic sclerosis: a gene array approach. *PLoS Med.* 2006; 3(1):e2.

- ❖ Lupus Foundation of America, Inc. [www document]. URL http://www.lupus.org/webmodules/webarticlesnet/templates/new_newsroomreporters.aspx?articleid=247&zoneid=60 [accessed on 5th August, 2012].
- ❖ Lupus: Symptoms [www document]. URL <http://www.mayoclinic.com/health/lupus/DS00115/DSECTION=symptoms> [accessed on 10th August, 2012].
- ❖ Luster MI, Portier C, Pait GG, White KL, Gennings C, Munson AE, Rosenthal G. Risk assessment in immunotoxicology. I: Sensitivity and predictability of immune tests. *Fundam Appl Toxicol.* 1992; 18(2):200–210.
- ❖ Magalhaes DAR, Silveira ELV, Junta EM, Sandrin-Garcia P, Fachin AL, Donadi EA, Sakamoto-Hojo ET, Passos GAS. Promiscuous gene expression in the thymus: the root of central tolerance. *Clin Dev Immuno.* 2006; 13(2-4):81-99.
- ❖ Magnusson V, Nakken B, Bolstad A, Alarcon-Riquelme M. Cytokine polymorphisms in systemic lupus erythematosus and Sjogren's syndrome. *Scand J Immunol.* 2001; 54(1-2):55–61.
- ❖ Magro CM, Crowson AN, Ferri C. Cytomegalovirus-associated cutaneous vasculopathy and scleroderma sans inclusion body change. *Hum Pathol.* 2007; 38(1):42-49.
- ❖ Malek TR, Bayer AL. Tolerance, not immunity, crucially depends on IL-2. *Nat Rev Immunol.* 2004; 4(9):665–674.
- ❖ Mandrup-Poulsen T, Pociot F, Molvig J, Shapiro L, Nilsson P, Emdal T, Røder M, Kjems LL, Dinarello CA, Nerup J. Monokine antagonism is reduced in IDDM. *Diabetes.* 1994; 43(10):1242–1247.
- ❖ Mansfield JC, Holden H, Tarlow JK, Di Giovine FS, McDowell TL, Wilson AG, Holdsworth CD, Duff GW. Novel genetic association between ulcerative colitis and the anti-inflammatory cytokine interleukin-1 receptor antagonist. *Gastroenterology.* 1994; 106(3):637– 642.
- ❖ Marrack P, Kappler J. The staphylococcal enterotoxins and their relatives. *Science.* 1990; 248(4956):705–711.
- ❖ Martinez A, Fernández-Arquero M, Pascual-Salcedo D, Conejero L, Alves H, Balsa A, de la Concha EG. Primary association of tumor necrosis factor-region genetic

- markers with susceptibility to rheumatoid arthritis. *Arthritis Rheum.* 2000; 43(6):1366–1370.
- ❖ Marusic-Galesic S, Pavelic K. Dynamics of positive and negative selection in the thymus: review and hypothesis. *Immunol Lett.* 1990; 24(3):149-154.
 - ❖ Masi AT. Classification of systemic sclerosis (scleroderma): relationship of cutaneous subgroups in early disease to outcome and serologic reactivity. *J Rheumatol.* 1988; 15(6):894–898.
 - ❖ Mathai SK, Gulati M, Peng X, Russell TR, Shaw AC, Rubinowitz AN, Murray LA, Siner JM, Antin-Ozerkis DE, Montgomery RR, Reilkoff RA, Bucala RJ, Herzog EL. Circulating monocytes from systemic sclerosis patients with interstitial lung disease show an enhanced profibrotic phenotype. *Lab Invest.* 2010; 90(6):812-823.
 - ❖ Mathis D, Benoist C. Back to central tolerance. *Immunity.* 2004; 20(5):509–516.
 - ❖ Matis LA. The molecular basis of T-cell specificity. *Annu Rev Immunol.* 1990; 8:65-82.
 - ❖ Matsushita T, Hasegawa M, Hamaguchi Y, Takehara K, Sato S. Longitudinal analysis of serum cytokine concentrations in systemic sclerosis: association of interleukin 12 elevation with spontaneous regression of skin sclerosis. *J Rheumatol.* 2006; 33(2):275–284.
 - ❖ Mattuzzi S, Barbi S, Carletto A, Ravagnani V, Moore PS, Bambara LM, Scarpa A. Association of polymorphisms in the IL1B and IL2 genes with susceptibility and severity of systemic sclerosis. *J Rheumatol.* 2007; 34(5):997–1004.
 - ❖ Maul GG, Jimenez SA, Riggs E, Ziemnicka-Kotula D. Determination of an epitope of the diffuse systemic sclerosis marker antigen DNA topoisomerase I: sequence similarity with retroviral p30gag protein suggests a possible cause for autoimmunity in systemic sclerosis. *Proc Natl Acad Sci USA.* 1989; 86(21):8492.
 - ❖ Mavalia C, Scaletti C, Romagnani P, Carossino AM, Pignone A, Emmi L, Pupilli C, Pizzolo G, Maggi E, Romagnani S. Type 2 helper T-cell predominance and high CD30 expression in systemic sclerosis. *Am J Pathol.* 1997; 151(6):1751–1758.
 - ❖ Mayes MD, Lacey JV, Beebe-Dimmer J, Gillespie BW, Cooper B, Laing TJ, Schottenfeld D. Prevalence, incidence, survival, and disease characteristics of systemic sclerosis in a large US population. *Arthritis Rheum.* 2003; 48(8):2246-2255.

-
- ❖ Mayes MD. Classification and epidemiology of scleroderma. *Semin Cutan Med Surg.* 1998; 17(1):22-26.
 - ❖ Mayes MD. Epidemiologic studies of environmental agents and systemic autoimmune diseases. *Environ Health Perspect.* 1999; 107(Suppl 5):743-748.
 - ❖ McConnachie PR, Zahalsky AC. Immune alterations in humans exposed to the termiticide technical chlordane. *Arch Environ Health.* 1992; 47(4):295–301.
 - ❖ McDowell TL, Symons JA, Ploski R, Førre O, Duff GW. A genetic association between juvenile rheumatoid arthritis and a novel interleukin-1 α polymorphism. *Arthritis Rheum.* 1995; 38(2):221–228.
 - ❖ McGaha TL, Bona CA. Role of profibrogenic cytokines secreted by T cells in fibrotic processes in scleroderma. *Autoimmun Rev.* 2002; 1(13):174–181.
 - ❖ McVean GAT, Myers SR, Hunt S, Deloukas P, Bentley DR, Donnelly P. The fine-scale structure of recombination rate variation in the human genome. *Science.* 2004; 304(5670):581–584.
 - ❖ Medsger TA Jr. Epidemiology of systemic sclerosis. *Clin Dermatol.* 1994; 12(2):207-216.
 - ❖ Medsger TA. Systemic sclerosis (scleroderma): clinical aspects. In: Koopman WJ, editor. *Arthritis and Allied Conditions*, 14th ed. Philadelphia: Lippincott Williams & Wilkins, 2001; p. 1590-1624.
 - ❖ Melendro EI, Saldate C, Rivero SJ, Alarcon-Segovia D. T-cell subpopulations in the peripheral blood of patients with connective tissue diseases as determined by flowcytometry using monoclonal antibodies. *Clin Immunol Immunopathol.* 1983; 27(3):340-347.
 - ❖ Mellor-Pita S, Citores MJ, Castejon R, Tutor-Ureta P, Yebra-Bango M, Andreu JL, Vargas JA. Decrease of regulatory T cells in patients with systemic lupus erythematosus. *Ann Rheum Dis.* 2006; 65(4):553–554.
 - ❖ Meyer MF, Hellmich B, Kotterba S, Schatz H. Cytomegalovirus infection in systemic necrotizing vasculitis: causative agent or opportunistic infection? *Rheumatol Int.* 2000; 20(1):35-38.

-
- ❖ Miethke T, Duschek K, Wahl C, Heeg K, Wagner H. Pathogenesis of the toxic shock syndrome: T cell mediated lethal shock caused by the superantigen TSST-1. *Eur J Immunol.* 1993; 23(7):1494–1500.
 - ❖ Mignogna MD, Lo Muzio L, Ruocco E. Pemphigus induction by influenza vaccination (letter). *Int J Dermatol.* 2000; 39(10):800.
 - ❖ Miller A, Lider O, Weiner HL. Antigen-driven bystander suppression following oral administration of antigens. *J Exp Med.* 1991; 174(4):791-798.
 - ❖ Mills KH, McGuirk P. Antigen-specific regulatory T cells—their induction and role in infection. *Semin Immunol.* 2004; 16(2):107-117.
 - ❖ Miyara M, Amoura Z, Parizot C, Badoual C, Dorgham K, Trad S, Nochy D, Debré P, Piette JC, Gorochov G. Global natural regulatory T cell depletion in active systemic lupus erythematosus. *J Immunol.* 2005; 175(12):8392-8400.
 - ❖ Miyara M, Yoshioka Y, Kitoh A, Shima T, Wing K, Niwa A, Parizot C, Taflin C, Heike T, Valeyre D, Mathian A, Nakahata T, Yamaguchi T, Nomura T, Ono M, Amoura Z, Gorochov G, Sakaguchi S. Functional delineation and differentiation dynamics of human CD4⁺ T cells expressing the FoxP3 transcription factor. *Immunity.* 2009; 30(6):899–911.
 - ❖ Mok CC, Lau CS. Pathogenesis of systemic lupus erythematosus. *J Clin Pathol.* 2003; 56(7):481–490.
 - ❖ Moore KW, de Waal Malefyt R, Coffman RL, O’Garra A. Interleukin-10 and the interleukin-10 receptor. *Annu Rev Immunol.* 2001; 19:683–765.
 - ❖ Moore KW, O’Garra A, de Waal Malefyt R, Vieira P, Mossman TR. Interleukin-10. *Annu Rev Immunol.* 1993; 11:165–190.
 - ❖ Moore KW, Vieira P, Fiorentino DF, Trounstein ML, Khan TA, Mossman TR. Homology of cytokine synthesis inhibitory factor (IL-10) to the Epstein–Barr virus gene BCRFI. *Science.* 1990; 248(4960):1230–1234.
 - ❖ Morshed S, Mercadante MT, Lombroso PJ. Genetics of childhood disorders XXVIII. Auto immunity part I. *J Am Acad adolesc psychiatry.* 2000; 40(7):855-858.
 - ❖ Mosmann TR, Coffman RL. Th1 and Th2 cell: different patterns of lymphokine secretion lead to different functional properties. *Annu Rev Immunol.* 1989; 7:145–173.

- ❖ Mudd PA, Teague BN, Farris AD. Regulatory T-cells and systemic lupus erythematosus. *Scand J Immunol.* 2006; 64(3):211–218.
- ❖ Muller W, Kuhn R, Rajewsky K. Major histocompatibility complex class II hyperexpression on B cells in interleukin 4-transgenic mice does not lead to B cell proliferation and hypergammaglobulinemia. *Eur J Immunol.* 1991; 21(4):921–925.
- ❖ Mulloy KB. Silica exposure and systemic vasculitis. *Environ Health Perspect.* 2003; 111(16): 1933–1938.
- ❖ Munoz Fernandez MA, Fernandez MA, Fresno M. Synergism between tumor necrosis factor- α and interferon- γ on macrophage activation for the killing of intracellular *Trypanosoma cruzi* through a nitric oxide-dependent mechanism. *Eur J Immunol.* 1992; 22(2):301-307.
- ❖ Muraki Y, Tsutsumi A, Takahashi R, Suzuki E, Hayashi T, Chino Y, Goto D, Matsumoto I, Murata H, Noguchi E, Sumida T. Polymorphisms of the IL-1 β gene in Japanese patients with Sjogren's syndrome and systemic lupus erythematosus. *J Rheumatol.* 2004; 31(4):720–725.
- ❖ Murali-Krishna K, Altman JD, Suresh M, Sourdive DJ, Zajac AJ, Miller JD, Slansky J, Ahmed R. Counting antigen-specific CD8 T cells: a reevaluation of bystander activation during viral infection. *Immunity.* 1998; 8(2):177–187.
- ❖ Murphy KM, Reiner SL. The lineage decisions of helper T-cells. *Nat Rev Immunol.* 2002; 2(12):933-944.
- ❖ Mydlarski PR, Ho V, Shear NH. Canadian consensus statement on the use of intravenous immunoglobulin therapy in dermatology. *J Cutan Med Surg.* 2006; 10(5):205-21.
- ❖ Nakamura K, Kitani A, Strober W. Cell contact-dependent immunosuppression by CD4⁺CD25⁺ regulatory T cells is mediated by cell surface-bound transforming growth factor beta. *J Exp Med.* 2001; 194(5):629–644.
- ❖ Narbutt J, Lukamowicz J, Bogaczewicz J, Sysa-Jedrzejowska A, Torzecka JD, Lesiak A. Serum concentration of interleukin-6 is increased both in active and remission stages of pemphigus vulgaris. *Mediators Inflamm.* 2008; 2008:875394.
- ❖ Nemazee D, Hogquist KA. Antigen receptor selection by editing or downregulation of V (D) J recombination. *Curr Opin Immunol.* 2003; 15(2):182–189.

-
- ❖ Nerheim PL, Meier JL, Vasef MA, Li WG, Hu L, Rice JB, Gavrilu D, Richenbacher WE, Weintraub NL. Enhanced cytomegalovirus infection in atherosclerotic human blood vessels. *Am J Pathol.* 2004; 164(2):589–600.
 - ❖ Nishizuka Y. The role of protein kinase C in cell surface signal transduction and tumor promotion. *Nature.* 1984; 308:693–698.
 - ❖ O’Garra A, Murphy K. T-cell subsets in autoimmunity. *Curr Opin Immunol.* 1993; 5(6):880-886.
 - ❖ Ohashi PS. Negative selection and autoimmunity. *Curr Opin Immunol.* 2003; 15(6):668–676.
 - ❖ Ohkura N, Hamaguchi M, Sakaguchi S. FOXP3⁺ regulatory T cells: control of FOXP3 expression by pharmacological agents. *Trends Pharmacol Sci.* 2011; 32(3):158–166.
 - ❖ Ohtsuka K, Gray JD, Stimmler MM, Horwitz DA. The relationship between defects in lymphocyte production of transforming growth factor-beta 1 in systemic lupus erythematosus and disease activity or severity. *Lupus.* 1999; 8(2):90–94.
 - ❖ Ohtsuka T, Yamakage A, Yamazaki S. The polymorphism of transforming growth factor-beta1 gene in Japanese patients with systemic sclerosis. *Br J Dermatol.* 2002; 147(3):458-463.
 - ❖ Oliver GF, Winkelmann RK. The current treatment of scleroderma. *Drugs.* 1989; 37(1):87–96.
 - ❖ Olsen NJ, King LE, Park JH. Muscle abnormalities in scleroderma. *Rheum Dis Clin North Am.* 1996; 22(4):783–796.
 - ❖ Oltz EM. Regulation of antigen receptor gene assembly in lymphocytes. *Immunol Res.* 2001; 23(2-3):121-133.
 - ❖ Ortonne JP. Recent developments in the understanding of the pathogenesis of psoriasis. *Br J Dermatol.* 1999; 140(Suppl 54):1-7.
 - ❖ Palma-Carlos AG, Palma-Carlos ML, da Silva SL. Natural killer (NK) cells in mucocutaneous candidiasis. *Allerg Immunol (Paris).* 2002; 34(6):208-212.
 - ❖ Parel Y, Chizzolini C. CD4⁺CD8⁺ double positive (DP) cells in health and disease. *Autoimmun Rev.* 2004; 3(3):215–220.

-
- ❖ Park SG, Chang JY, Cho YH, Kim SC, Lee MG. Transition from pemphigus foliaceus to pemphigus vulgaris: case report with literature review. *Yonsei Med J.* 2006; 47(2):278–281.
 - ❖ Parks CG, Cooper GS, Dooley MA, Treadwell EL, St Clair EW, Gilkeson GS, Pandey JP. Systemic lupus erythematosus and genetic variation in the interleukin 1 gene cluster: a population based study in the southeastern United States. *Ann Rheum Dis.* 2004; 63(1):91-94.
 - ❖ Parks CG, Pandey JP, Dooley MA, Treadwell EL, St Clair EW, Gilkeson GS, Feghali-Bostwick CA, Cooper GS. Genetic polymorphism in tumor necrosis factor (TNF)-alpha and TNF-beta in a population-based study of systemic lupus erythematosus: associations and interaction with the interleukin-1alpha-889 C/T polymorphism. *Hum Immunol.* 2004; 65(6):622–631.
 - ❖ Parronchi P, de Carli M, Manetti R, Simonelli C, Sampognaro S, Piccinni MP, Macchia D, Maggi E, Del Prete G, Romagnani S. IL-4 and IFN (alpha and gamma) exert opposite regulatory effects on the development of cytolytic potential by Th1 or Th2 human T cell clones. *J Immunol.* 1992; 149(9):2977–2983.
 - ❖ Patil N, Berno AJ, Hinds DA, Barrett WA, Doshi JM, Hacker CR, Kautzer CR, Lee DH, Marjoribanks C, McDonough DP, Nguyen BT, Norris MC, Sheehan JB, Shen N, Stern D, Stokowski RP, Thomas DJ, Trulson MO, Vyas KR, Frazer KA, Fodor SPA, Cox DR. Blocks of limited haplotype diversity revealed by high-resolution scanning of human chromosome 21. *Science.* 2001; 294(5547):1719–1723.
 - ❖ Pender MP. Infection of autoreactive B lymphocytes with EBV, causing chronic autoimmune diseases. *Trends Immunol.* 2003; 24(11):584–588.
 - ❖ Perez-Maldonado IN, Diaz-Barriga F, de la Fuente H, Gonzalez-Amaro R, Calderon J, Yanez L. DDT induces apoptosis in human mononuclear cells in vitro and is associated with increased apoptosis in exposed children. *Environ Res.* 2004; 94(1):38–46.
 - ❖ Perrey C, Pravica V, Sinnott PJ, Hutchinson IV. Genotyping for polymorphisms in interferon- γ , interleukin-10, transforming growth factor- β 1 and tumor necrosis factor- α genes: a technical report. *Transpl Immunol.* 1998; 6(3):193–197.

-
- ❖ Petri M, Robinson C. Oral contraceptives and systemic lupus erythematosus. *Arthritis Rheumatol.* 1997; 40(5):797-803.
 - ❖ Petri M. Epidemiology of systemic lupus erythematosus. *Best Pract Res Clin Rheumatol.* 2002; 16(5):847-858.
 - ❖ Petrovic R. Sistemski eritemski lupus. In: Pilipovic N, editor. *Reumatologija Beograd: Zavod za udzbenike i nastavna sredstva*, 2000; p. 384-387.
 - ❖ Pillai V, Ortega SB, Wang CK, Karandikar NJ. Transient regulatory T-cells: A state attained by all activated human T-cells. *Clin Immunol.* 2007; 123(1):18–29.
 - ❖ Ploski R, McDowell TL, Symons JA, Flatø B, Duff GW, Thorsby E, Førre O. Interaction between HLA-DR and HLA-DP, and between HLA and interleukin 1a in juvenile rheumatoid arthritis indicates heterogeneity of pathogenic mechanisms of the disease. *Hum Immunol.* 1995; 42(4):343–347.
 - ❖ Plumb R, Doolittle J. Paraneoplastic pemphigus in a patient with non- Hodgkin's lymphoma. *Am J Hematol.* 1996; 52(1):58-59.
 - ❖ Pociot F, Briant L, Jongeneel CV, Mølviq J, Worsaae H, Abbal M, Thomsen M, Nerup J, Cambon-Thomsen A. Association of tumor necrosis factor (TNF) and class II major histocompatibility complex alleles with the secretion of TNF-alpha and TNF-beta by human mononuclear cells: a possible link to insulin-dependent diabetes mellitus. *Eur J Immunol.* 1993; 23(1):224–231.
 - ❖ Pociot F, Mølviq J, Wogensen L, Worsaae H, Nerup J. A TaqI polymorphism in the human interleukin-1b (IL-1b) gene correlates with IL-1 beta secretion in vitro. *Eur J Clin Invest.* 1992; 22(6):396–402.
 - ❖ Pociot F, Ronningen KS, Bergholdt R, Lorenzen T, Johannesen J, Ye K, Dinarello CA, Nerup J. Genetic susceptibility in Danish patients with type 1 (insulin dependent) diabetes: evidence for polygenicity in man. Danish Study Group of Diabetes in Childhood. *Autoimmunity.* 1994; 19(3):169–178.
 - ❖ Pollard KM, Pearson DL, Hultman P, Hildebrandt B, Kono DH. Lupus-prone mice as models to study xenobiotic-induced acceleration of systemic autoimmunity. *Environ Health Perspect.* 1999; 107(Suppl 5):729–735.
 - ❖ Poole BD, Scofield RH, Harley JB, James JA. Epstein-Barr virus and molecular mimicry in systemic lupus erythematosus. *Autoimmunity.* 2006; 39(1):63–70.

-
- ❖ Povey A, Guppy MJ, Wood M, Knight C, Black CM, Silman AJ. Cytochrome P2 polymorphism and susceptibility to scleroderma following exposure to organic solvents. *Arthritis Rheum.* 2001; 44(3):662–665.
 - ❖ Pravica V, Asderakis A, Perrey C, Hajeer A, Sinnott PJ, Hutchinson IV. In vitro production of IFN-gamma correlates with CA repeat polymorphism in the human IFN-gamma gene. *Eur J Immunogenet.* 1999; 26(1):1–3.
 - ❖ Pravica V, Perrey C, Stevens A, Lee JH, Hutchinson IV. A single nucleotide polymorphism in the first intron of the human IFN-gamma gene: Absolute correlation with a polymorphic CA microsatellite marker of high IFN-gamma production. *Hum Immunol.* 2000; 61(9):863–866.
 - ❖ Proft T, Fraser JD. Bacterial superantigens. *Clin Exp Immunol.* 2003; 133(3):299–306.
 - ❖ Qiu H, Straat K, Rahbar A, Wan M, Soderberg-Naucler C, Haeggstrom JZ. Human CMV infection induces 5-lipoxygenase expression and leukotriene B4 production in vascular smooth muscle cells. *J Exp Med.* 2008; 205(1):19–24.
 - ❖ Rahman A, David A. Systemic Lupus Erythematosus. *N Engl J Med.* 2008; 358(9):929–939.
 - ❖ Raman K, Mohan C. Genetic underpinnings of autoimmunity-lessons from studies in arthritis, diabetes, lupus and multiple sclerosis. *Curr Opin Immunol.* 2003; 15(6):651–659.
 - ❖ Raziuddin S, Nur MA, Al-Wabel AA. Increased circulating HLA-DR+ CD4+ T-cell in systemic lupus erythematosus: Alterations associated with prednisolone therapy. *Scand J Immunol.* 1990; 31(2):139–145.
 - ❖ Rea WJ, Johnson AR, Youdim S, Fenyves EJ, Samadi N. T- and B-Lymphocyte parameters measured in chemically sensitive patients and controls. *Clin Ecol.* 1986; 4:11–14.
 - ❖ Reich DE, Schaffner SF, Daly MJ, McVean G, Mullikin JC, Higgins JM, Richter DJ, Lander ES, Altshuler D. Human genome sequence variation and the influence of gene history, mutation and recombination. *Nature Genet.* 2002; 32(1):135–142.

-
- ❖ Reinherz EL, Schlossman SF. Current concepts in immunology: regulation of the immune response—inducer and suppressor T lymphocyte subsets in human beings. *N Engl J Med.* 1980; 303(7):370-373.
 - ❖ Renno T, Acha-Orbea H. Superantigens in autoimmune diseases: still more shades of gray. *Immunol Rev.* 1996; 154:175–191.
 - ❖ Rider LG, Artlett CM, Foster CB, Ahmed A, Neeman T, Chanock SJ, Jimenez SA, Miller FW. Polymorphisms in the IL-1 receptor antagonist gene VNTR are possible risk factors for juvenile idiopathic inflammatory myopathies. *Clin Exp Immunol.* 2000; 121(1):47–52.
 - ❖ Roberts BW, Cech I. Association of type 2 diabetes mellitus and seroprevalence for cytomegalovirus. *South Med J.* 2005; 98(7):686–692.
 - ❖ Robinson ND, Hashimoto T, Amagai M, Chan LS. The new pemphigus variants. *J Am Acad Dermatol.* 1999; 40(5 Pt 1):649-667.
 - ❖ Romagnani S. Biology of human TH1 and TH2 cells. *J Clin Immunol.* 1995; 15(3):121-129.
 - ❖ Rood MJ, van Krugten MV, Zanelli E, van der Linden MW, Keijsers V, Schneider GMT, Verduyn W, Westendorp RGJ, de Vries RRP, Breedveld FC, Verweij CL, Huizinga TWJ. TNF-308A and HLA-DR3 alleles contribute independently to susceptibility to systemic lupus erythematosus. *Arthritis Rheum.* 2000; 43(1):129–134.
 - ❖ Rose S, Young MA, Reynolds JC. Gastrointestinal manifestations of scleroderma. *Gastroenterol Clin North Am.* 1998; 27(3):563–594.
 - ❖ Rosenwasser LJ, Klemm DJ, Dresback JK, Inamura H, Mascali JJ, Klinnert M, Borish L. Promotor polymorphisms in the chromosome 5 gene cluster in asthma and atopy. *Clin Exp Allergy.* 1995; 25(Suppl 2):74–78.
 - ❖ Rothfeild NF. Cutaneous Manifestations of Multisystem Diseases: Systemic Lupus Erythematosus. In: Fitzpatrick TB, Eisen AZ, Wolff K, Freedberg IM, Austen KF editors. *Dermatology in General Medicine*, 4th ed. New York: McGraw-Hill, 1993; p. 2143-2148. v.2.

- ❖ Rothfield N. Clinical features of systemic lupus erythematosus. In: Kelley WN, Harris ED, Ruddy S, Sledge CB, editors. *Textbook of Rheumatology*. Philadelphia: WB Saunders, 1981.
- ❖ Roubinian JR, Talal N, Greenspan JS, Goodman JR, Siiteri PK. Effect of castration and sex hormone treatment on survival, anti-nucleic acid antibodies, and glomerulonephritis in NZB/NZW F1 mice. *J Exp Med*. 1978; 147(6):1568–1583.
- ❖ Rousset F, Garcia E, Defrance T, Peronne C, Hsu DH, Kastelein R, Moore KW, Banchereau J. Interleukin-10 is a potent growth and differentiation factor for activated human B-lymphocytes. *Proc Natl Acad Sci USA*. 1992; 89(5):1890–1893.
- ❖ Roux F, Treich I, Brun C, Desoize B, Fournier E. Effect of lindane on human lymphocyte responses to phytohemagglutinin. *Biochem Pharmacol*. 1979; 28(16):2419–2426.
- ❖ Rowen L, Koop BF, Hood L. The complete 685-kilobase DNA sequence of the human β T cell receptor locus. *Science*. 1996; 272(5269):1755-1762.
- ❖ Ruach M, Ohel G, Rahav D. Pemphigus vulgaris and pregnancy. *Obstet Gynecol Surv*. 1995; 50(10):755-760.
- ❖ Ruangjutipopan S, Kasitanon N, Louthrenoo W, Sukitawut W, Wichainun R. Causes of death and poor survival prognostic factors in Thai patients with systemic sclerosis. *J Med Assoc Thai*. 2002; 85(11):1204–1209.
- ❖ Ruocco V, Wolf R, Ruocco E, Baroni A. Viruses in pemphigus: A casual or causal relationship? *Int J Dermatol*. 1996; 35(11):782-784.
- ❖ Rus V, Maury EE, Hochberg MC. The epidemiology of systemic lupus erythematosus. In: Wallace DJ, Hahn BH, editors. *Dubois' Lupus Erythematosus*. Philadelphia: Lippincott Williams and Wilkins, 2002.
- ❖ Sabry A, Sheashaa H, El-Husseini A, Mahmoud K, Eldahshan KF, George SK, Abdel-Khalek E, El-Shafey EM, Abo-Zenah H. Proinflammatory cytokines (TNF-alpha and IL-6) in Egyptian patients with SLE: its correlation with disease activity. *Cytokine*. 2006; 35(3-4):148–153.
- ❖ Sachidanandam R, Weissman D, Schmidt SC, Kakol JM, Stein LD, Marth G, Sherry S, Mullikin JC, Mortimore BJ, Willey DL, Hunt SE, Cole CG, Coggill PC, Rice CM, Ning Z, Rogers J, Bentley DR, Kwok PY, Mardis ER, Yeh RT, Schultz B, Cook L,

- Davenport R, Dante M, Fulton L, Hillier L, Waterston RH, McPherson JD, Gilman B, Schaffner S, Van Etten WJ, Reich D, Higgins J, Daly MJ, Blumenstiel B, Baldwin J, Stange-Thomann N, Zody MC, Linton L, Lander ES, Altshuler D; International SNP Map Working Group. A map of human genome sequence variation containing 1.42 million single nucleotide polymorphisms. *Nature*. 2001; 409(6822):928–933.
- ❖ Sad S, Marcotte R, Mosmann TR. Cytokine-induced differentiation of precursor mouse CD8⁺ T cells into cytotoxic CD8⁺ T cells secreting Th1 or Th2 cytokines. *Immunity*. 1995; 2(3):271–279.
 - ❖ Sagi L, Baum S, Agmon-Levin N, Sherer Y, Katz BSP, Barzilai O, Ram M, Bizzaro N, SanMarco M, Trau H, Shoenfeld Y. Autoimmune bullous diseases: The spectrum of infectious agent antibodies and review of the literature. *Autoimmun Rev*. 2011; 10(9):527–535.
 - ❖ Sagi L, Sherer Y, Trau H, Shoenfeld Y. Pemphigus and infectious agents. *Autoimmun Rev*. 2008; 8(1):33-35.
 - ❖ Sakaguchi S, Ono M, Setoguchi R, Yagi H, Hori S, Fehervari Z, Shimizu J, Takahashi T, Nomura T. Foxp3⁺CD25⁺CD4⁺ natural regulatory T cells in dominant self tolerance and autoimmune disease. *Immunol Rev*. 2006; 212:8-27.
 - ❖ Sakaguchi S, Wing K, Onishi Y, Prieto-Martin P, Yamaguchi T. Regulatory T cells: how do they suppress immune responses? *Int Immunol*. 2009; 21(10):1105–1111.
 - ❖ Sakaguchi S. Naturally arising CD4⁺ regulatory T cells for immunologic self-tolerance and negative control of immune responses. *Annu Rev Immunol*. 2004; 22:531–562.
 - ❖ Sakaguchi S. Naturally arising Foxp3-expressing CD25⁺CD4⁺ regulatory T cells in immunological tolerance to self and non-self. *Nat Immunol*. 2005; 6:345–352.
 - ❖ Saline M, Rodstrom KE, Fischer G, Orekhov VY, Karlsson BG, Lindkvist-Petersson K. The structure of superantigen complexed with TCR and MHC reveals novel insights into superantigenic T cell activation. *Nat Commun*. 2010; 1:119.
 - ❖ Sallam H, McNearney TA, Chen JD. Systematic review: pathophysiology and management of gastrointestinal dysmotility in systemic sclerosis (scleroderma). *Aliment Pharmacol Ther*. 2006; 23(6):691–712.

-
- ❖ Salmon M, Pilling D, Borthwick NJ, Viner N, Janossy G, Bacon PA, Akbar, AN. The progressive differentiation of primed T cells is associated with an increasing susceptibility to apoptosis. *Eur J Immunol.* 1994; 24(4):892-899.
 - ❖ Salomon B, Bluestone JA. Complexities of CD28/B7: CTLA-4 costimulatory pathways in autoimmunity and transplantation. *Annu Rev Immunol.* 2001; 19:225–252.
 - ❖ Sant AJ. Endogenous antigen presentation by MHC class II molecules. *Immunol Res.* 1994; 13(4):253-267.
 - ❖ Santiago ML, Fossati L, Jacquet C, Muller W, Izui S, Reininger L. Interleukin-4 protects against a genetically linked lupus-like autoimmune syndrome. *J Exp Med.* 1997; 185(1):65–70.
 - ❖ Sato S, Hasegawa M, Takehara K. Serum levels of interleukin- 6 and interleukin-10 correlate with total skin thickness score in patients with systemic sclerosis. *J Dermatol Sci.* 2001; 27(2):140–146.
 - ❖ Saunus JM, Kazoullis A, Farah CS. Cellular and molecular mechanisms of resistance to oral *Candida albicans* infections. *Front Biosci.* 2008; 13:5345-5358.
 - ❖ Schaller J. Lupus in childhood. *Clin Rheum Dis.* 1982; 8(1):219-228.
 - ❖ Scheffold A, Huhn J, Hofer T. Regulation of CD4 +CD25 + regulatory T-cell activity: It takes (IL-) two to tango. *Eur J Immunol.* 2005; 35(5):1336–1341.
 - ❖ Scherer HU, Burmester GR, Riemekasten G. Targeting activated T cells: successful use of anti-CD25 monoclonal antibody basiliximab in a patient with systemic sclerosis. *Ann Rheum Dis.* 2006; 65(9):1245–1247.
 - ❖ Scherer MT, Ignatowicz L, Winslow GM, Kappler JW, Marrack P. Superantigens: bacterial and viral proteins that manipulate the immune system. *Annu Rev Cell Biol.* 1993; 9:101–128.
 - ❖ Schiffenbauer J, Soos JM, Johnson HM. The possible role of bacterial superantigens in the pathogenesis of autoimmune disorders. *Immunol Today.* 1998; 19(3):117-120.
 - ❖ Schumacher TN. T-cell-receptor gene therapy. *Nat Rev Immunol.* 2002; 2(7):512-519.
 - ❖ Schwartz RH. T cell anergy. *Annu Rev Immunol.* 2003; 21:305–334.

-
- ❖ Scussel-Lonzetti L, Joyal F, Raynauld JP, Roussin A, Rich E, Goulet JR, Raymond Y, Sénécal JL. Predicting mortality in systemic sclerosis: Analysis of a cohort of 309 French Canadian patients with emphasis on features at diagnosis as predictive factors for survival. *Medicine (Baltimore)*. 2002; 81(2):154-67.
 - ❖ Seibold JR, McCloskey DA. Skin involvement as a relevant outcome measure in clinical trials of systemic sclerosis. *Curr Opin Rheumatol*. 1997; 9(6):571–575.
 - ❖ Sekigawa I, Nawata M, Seta N, Yamada M, Iida N, Hashimoto H. Cytomegalovirus infection in patients with systemic lupus erythematosus. *Clin Exp Rheumatol*. 2002; 20(4):559–564.
 - ❖ Serdula MK, Rhoads GG. Frequency of systemic lupus erythematosus in different ethnic groups in Hawaii. *Arthritis Rheum*. 1979; 22(4):328-333.
 - ❖ Shand L, Lunt M, Nihtyanova S, Hoseini M, Silman A, Black CM, Denton CP. Relationship between change in skin score and disease outcome in diffuse cutaneous systemic sclerosis: Application of a latent linear trajectory model. *Arthritis Rheum*. 2007; 56(7):2422-2431.
 - ❖ Shevach EM. Mechanisms of foxp3+ T regulatory cell-mediated suppression. *Immunity*. 2009; 30(5):636–645.
 - ❖ Shinkai Y, Rathbun G, Lam KP, Oltz EM, Stewart V, Mendelsohn M, Charron J, Datta M, Young F, Stall AM, et al. RAG-2 deficient mice lack mature lymphocytes owing to inability to initiate V (D) J rearrangement. *Cell*. 1992; 68(5):885-867.
 - ❖ Shlomchik MJ, Cooke A, Weigert M. Autoimmunity: The genes and phenotypes of autoimmunity. *Curr Opin Immunol*. 2004; 16(6):738–740.
 - ❖ Siegmund W, Franke G, Biebler KE, Donner I, Kawellis R, Kairies M, Scherber A, Huller H. The influence of the acetylator phenotype on the clinical use of dihydralazine. *Int J Clin Pharmacol Ther Toxicol*. 1985; 23(Suppl 1):S74–78.
 - ❖ Silkworth JB, Loose LD. Environmental chemical-induced modification of cell-mediated immune responses. *Adv Exper Med Biol*. 1979; 121(A):499–522.
 - ❖ Sim BC, Aftahi N, Reilly C, Bogen B, Schwartz RH, Gascoigne NR, Lo D. Thymic skewing of the CD4/CD8 ratio maps with the T-cell receptor alpha-chain locus. *Curr Biol*. 1998; 8(12):701-704.

-
- ❖ Singh B, Read S, Asseman C, Malmström V, Mottet C, Stephens LA, Stepankova R, Tlaskalova H, Powrie F. Control of intestinal inflammation by regulatory T cells. *Immunol Rev.* 2001; 182:190–200.
 - ❖ Slinger E, Maussang D, Schreiber A, Siderius M, Rahbar A, Fraile-Ramos A, Lira SA, Soderberg-Naucler C, Smit MJ. HCMV-encoded chemokine receptor US28 mediates proliferative signaling through the IL-6–STAT3 axis. *Sci Signal.* 2010; 3(133):ra58.
 - ❖ Smith SH, Brown MH, Rowe D, Callard RE, Beverley PC. Functional subsets of human help-inducer cells defined by a new monoclonal antibody, UCHL1. *Immunology.* 1986; 58(1):63–70.
 - ❖ Sobel ES, Gianini J, Butfiloski EJ, Croker BP, Schiffenbauer J, Roberts SM. Acceleration of autoimmunity by organochlorine pesticides in (NZB x NZW) F1 mice. *Environ Health Perspect.* 2005; 113(3):323–328.
 - ❖ Soderberg-Naucler C. Autoimmunity induced by human cytomegalovirus in patients with systemic lupus erythematosus. *Arthritis Research & Therapy.* 2012; 14(1):101.
 - ❖ Solomou EE, Juang YT, Gourley MF, Kammer GM, Tsokos GC. Molecular basis of deficient IL-2 production in T cells from patients with systemic lupus erythematosus. *J Immunol.* 2001; 166(6):4216–4222.
 - ❖ Sontheimer RD, Stastny P, Gilliam JN. Human histocompatibility antigen associations in subacute cutaneous lupus erythematosus. *J Clin Invest.* 1981; 67(1):312-316.
 - ❖ Soos JM, Schiffenbauer J, Torres BA, Johnson HM. Superantigens as virulence factors in autoimmunity and immunodeficiency diseases. *Med Hypotheses.* 1997; 48(3): 253–259.
 - ❖ Sozeri B, Gulez N, Aksu G, Kutukculer N, Akalin T, Kandiloglu G. Pesticide-induced scleroderma and early intensive immunosuppressive treatment. *Arch Environ Occup Health.* 2012; 67(1):43-47.
 - ❖ Stanley JR. Pemphigus. In: Freedberg IM, Eisen AZ, Wolff K, et al, editors. *Dermatology in General Medicine*, 5th ed. New York: McGraw-Hill, 1999; p. 654-666.

-
- ❖ Steen VD, Medsger TA. Epidemiology and natural history of systemic sclerosis. *Rheum Dis Clin North Am.* 1990; 16(1):1-10.
 - ❖ Steen VD. The lung in systemic sclerosis. *J Clin Rheumatol.* 2005; 11(1):40–46.
 - ❖ Stefanova I, Dorfman JR, Germain RN. Self-recognition promotes the foreign antigen sensitivity of naïve T lymphocytes. *Nature.* 2002; 420(6914):429-434.
 - ❖ Stekman IL, Blasini AM, Leon-Ponte M, Baroja ML, Abadi I, Rodriguez MA. Enhanced CD3-mediated T-lymphocyte proliferation in patients with systemic lupus erythematosus. *Arthritis Rheum.* 1991; 34(4):459–467.
 - ❖ Stiles BG, Garza AR, Ulrich RG, Boles JW. Mucosal vaccination with recombinantly attenuated staphylococcal enterotoxin B and protection in a murine model. *Infect Immun.* 2001; 69(4):2031–2036.
 - ❖ Stohl W, Elliott JE, Linsley PS. Human T cell-dependent B cell differentiation induced by staphylococcal superantigens. *J Immunol.* 1994; 153(1):117-127.
 - ❖ Street JC, Sharma RP. Alteration of induced cellular and humoral immune responses by pesticides and chemicals of environmental concern: Quantitative studies of immunosuppression by DDT, Arochlor 1254, Carbaryl, Carbofuran, methylparathion. *Toxicol Appl Pharmacol.* 1975; 32(3):587–602.
 - ❖ Sturfelt G, Hellmer G, Truedsson L. TNF microsatellites in systemic lupus erythematosus - a high frequency of the TNFabc 2-3-1 haplotype in multicase SLE families. *Lupus.* 1996; 5(6):618–622.
 - ❖ Suarez A, Castro P, Alonso R, Mozo L, Gutierrez C. Inter-individual variations in constitutive interleukin-10 messenger RNA and protein levels and their association with genetic polymorphisms. *Transplantation.* 2003; 75(5):711–717.
 - ❖ Subcommittee for Scleroderma Criteria of the American Rheumatism Association Diagnostic and Therapeutic Criteria Committee. Preliminary criteria for the classification of systemic sclerosis (scleroderma). *Arthritis Rheum.* 1980; 23:581-590.
 - ❖ Suen JL, Li HT, Jong YJ, Chiang BL, Yen JH. Altered homeostasis of CD4 (+) FoxP3 (+) regulatory T-cell subpopulations in systemic lupus erythematosus. *Immunology.* 2009; 127(2):196–205.

-
- ❖ Sugiura Y, Banno S, Matsumoto Y, Niimi T, Yoshinouchi T, Hayami Y, Naniwa T, Ueda R. Transforming growth factor beta1 gene polymorphism in patients with systemic sclerosis. *J Rheumatol.* 2003; 30(7):1520–1523.
 - ❖ Sugiura Y, Niimi T, Sato S, Yoshinouchi T, Banno S, Naniwa T, Maeda H, Shimizu S, Ueda R. Transforming growth factor beta1 gene polymorphism in rheumatoid arthritis. *Ann Rheum Dis.* 2002; 61(9):826-828.
 - ❖ Sugiyama H, Matsue H, Nagasaka A, Nakamura Y, Tsukamoto K, Shibagaki N, Kawamura T, Kitamura R, Ando N, Shimada S. CD4+CD25high regulatory T cells are markedly decreased in blood of patients with pemphigus vulgaris. *Dermatology.* 2007; 214(3):210-220.
 - ❖ Sumida T, Matsumoto I, Maeda T. TCR repertoire of autoreactive T cells in autoimmune disorders. *Nippon Rinsho.* 1997; 55(6):1410-1418.
 - ❖ Sung YK, Park BL, Shin HD, Kim LH, Kim SY, Bae SC. Interleukin-10 gene polymorphisms are associated with the SLICC/ACR Damage Index in systemic lupus erythematosus. *Rheumatology (Oxford).* 2006; 45(4):400-404.
 - ❖ Susol E, Rands AL, Herrick A, McHugh N, Barrett JH, Ollier WER, Worthington J. Association of markers for TGFβ3, TGFβ2 and TIMP1 with systemic sclerosis. *Rheumatology.* 2000; 39:1332–1336.
 - ❖ Suttmuller M, Baelde HJ, Ouellette S, De Heer E, Bruijn JA. T-cell receptor V beta gene expression in experimental lupus nephritis. *Immunology.* 1998; 95(1):18-25.
 - ❖ Swain SL, Croft M, Dubey C, Haynes L, Rogers P, Zhang X, Bradley LM. From naive to memory T cells. *Immunol Rev.* 1996; 150:143–167.
 - ❖ Symmons DP. Frequency of lupus in people of African origin. *Lupus.* 1995; 4(3):176-178.
 - ❖ Takatsu K. Role of interleukin-5 in immune regulation and inflammation. *Nippon Rinsho.* 2004; 62(10):1941-1951.
 - ❖ Tan EM, Feltkamp TEW, Smolen JS, Butcher B, Dawkins R, Fritzler MJ, Gordon T, Hardin JA, Kalden JR, Lahita RG, Maini RN, McDougal JS, Rothfield NF, Smeenk RJ, Takasaki E, Whik A, Wilson MR, Koziol JA. Range of antinuclear antibodies in “healthy” individuals. *Arthritis Rheum.* 1997; 40(9):1601–1611.

-
- ❖ Tashkin DP, Elashoff R, Clements PJ, Goldin J, Roth MD, Furst DE, Arriola E, Silver R, Strange C, Bolster M, Seibold JR, Riley DJ, Hsu VM, Varga J, Schraufnagel DE, Theodore A, Simms R, Wise R, Wigley F, White B, Steen V, Read C, Mayes M, Parsley E, Mubarak K, Connolly MK, Golden J, Olman M, Fessler B, Rothfield N, Metersky M. Cyclophosphamide versus placebo in scleroderma lung disease. *N Engl J Med.* 2006; 354(25):2655–2666.
 - ❖ Tatic V, Dimitrijevic J, Mitrovic D, Popovic M. Znaej histopatoloskih ispitivanja u dijagnozi reumaticnih bolesti. In: Popovic M, editor. *Reumaticne i slicne bolesti.* Beograd: Vojno-izdavacki zavod, 2000; p. 842-856.
 - ❖ Tenbrock K, Tsokos GC. Transcriptional regulation of interleukin 2 in SLE T-cells. *Int Rev Immunol.* 2004; 23(3-4):333–345.
 - ❖ Terry C, Loukaci V, Green F. Cooperative influence of genetic polymorphisms on interleukin 6 transcriptional regulation. *J Biol Chem.* 2000; 275(24):18138–18144.
 - ❖ Teyton L, Peterson DA. Assembly and transport of MHC class II molecules. *New Biol.* 1992; 4(5):441-447.
 - ❖ Theofilopoulos AN. The basis of autoimmunity: mechanism of aberrant self recognition. *Immunol Today.* 1995; 16(2):90-98.
 - ❖ Tulusso B, Fabris M, Caporali R, Cuomo G, Isola M, Soldano F, Montecucco C, Valentini G, Ferraccioli G. -238 and +489 TNF-alpha along with TNF-RII gene polymorphisms associate with the diffuse phenotype in patients with systemic sclerosis. *Immunol Lett.* 2005; 96(1):103–108.
 - ❖ Tomura M, Honda T, Tanizaki H, Otsuka A, Egawa G, Tokura Y, Waldmann H, Hori S, Cyster JG, Watanabe T, Miyachi Y, Kanagawa O, Kabashima K. Activated regulatory T cells are the major T cell type emigrating from the skin during a cutaneous immune response in mice. *J Clin Invest.* 2010; 120(3):883–893.
 - ❖ Torres BA, Soos JM, Perrin GQ, Johnson HM. Microbial superantigens and immunological deregulation. In: Nataro JP, Blaser MJ, Cunningham-Rundles S, Eds. *Persistent bacterial infections.* Washington, DC: ASM Press, 2000; p. 183–197.
 - ❖ Totterman TH, Karlsson FA, Bengtsson M, Harlwig IM. Induction of circulating activated suppressor like T cells by melhimazole therapy for Graves' disease. *N Engl J Med.* 1987; 316(1):15-22.

- ❖ Treiner E, Duban L, Bahram S, Radosavljevic M, Wanner V, Tilloy F, Affaticati P, Gilfillan S, Lantz O. Selection of evolutionarily conserved mucosal-associated invariant T cells by MR1. *Nature*. 2003; 422(6928):164–169.
- ❖ Trimeche M, Ziadi S, Amara K, Khelifa M, Bahri F, Mestiri S, Braham H, Hachana M, Sriha B, Mokni M, Korbi S. Prevalence of Epstein-Barr virus in Sjogren's syndrome in Tunisia. *Rev Med Interne*. 2006; 27(7):519–523.
- ❖ Tsankov N, Kazandjieva J, Gantcheva M. Contact pemphigus induced by dihydrodiphenyltrichlorethane. *Eur J Dermatol*. 1998; 8(6):442–443.
- ❖ Tufano MA, Baroni A, Buommino E, Ruocco E, Lombard ML, Ruocco V. Detection of herpesvirus DNA in peripheral blood mononuclear cells and skin lesions of patients with pemphigus by polymerase chain reaction. *Br J Dermatol*. 1999; 141(6):1033–1039.
- ❖ Tumang JR, Zhou JL, Gietl D, Crow MK, Elkon KB, Friedman SM. T helper cell-dependent, microbial superantigen-mediated B cell activation in vivo. *Autoimmunity*. 1996; 24(4):247–255.
- ❖ Turner DM, Williams DM, Sankaran D, Lazarus M, Sinnott PJ, Hutchinson IV. An investigation of polymorphism in the interleukin-10 gene promoter. *Eur J Immunogenet*. 1997; 24(1):1–8.
- ❖ Tzankov VN, Stransky L, Kostowa M, Mitrowa T, Obreschkowa E. Induced pemphigus caused by occupational contact with basochrom. *Derm Beruf Umwelt*. 1990; 38(3):91–93.
- ❖ Ueda H, Howson JM, Esposito L, Heward J, Snook H, Chamberlain G, Rainbow DB, Hunter KM, Smith AN, Di Genova G, Herr MH, Dahlman I, Payne F, Smyth D, Lowe C, Twells RC, Howlett S, Healy B, Nutland S, Rance HE, Everett V, Smink LJ, Lam AC, Cordell HJ, Walker NM, Bordin C, Hulme J, Motzo C, Cucca F, Hess JF, Metzker ML, Rogers J, Gregory S, Allahabadia A, Nithiyananthan R, Tuomilehto-Wolf E, Tuomilehto J, Bingley P, Gillespie KM, Undlien DE, Rønningen KS, Guja C, Ionescu-Tîrgoviște C, Savage DA, Maxwell AP, Carson DJ, Patterson CC, Franklyn JA, Clayton DG, Peterson LB, Wicker LS, Todd JA, Gough SC. Association of the T-cell regulatory gene CTLA4 with susceptibility to autoimmune disease. *Nature*. 2003; 423(6939):506–511.

-
- ❖ Uramoto KM, Michet CJ Jr, Thumboo J, Sunku J, O'Fallon WM, Gabriel SE. Trends in the incidence and mortality of systemic lupus erythematosus, 1950-1992. *Arthritis Rheum.* 1999; 42(1):46-50.
 - ❖ US Environment Protection Agency. Method 3620B- Florisil clean up by column chromatography. [www document] URL <http://www.epa.gov/SW-846/pdfs/3620b.pdf> (accessed on 02 July 2011).
 - ❖ Uthman I, Tabbarah Z, Gharavi AE. Hughes syndrome associated with cytomegalovirus infection. *Lupus.* 1999; 8(9):775-777.
 - ❖ Valentini G, Black C. Systemic sclerosis. Best practice & research. *Clin rheumatol.* 2002; 16(5):807-816.
 - ❖ Valikhani M, Kavusi S, Chams-Davatchi C, Daneshpazhooh M, Barzegari M, Ghiasi M, Abedini R. Pemphigus and associated environmental factors: a case-control study. *Clin Exp Dermatol.* 2007; 32(3):256-260.
 - ❖ Valle A, Zuber CE, Defrance T, Djossou O, De Rie M, Banchereau J. Activation of human B lymphocytes through CD40 and interleukin 4. *Eur J Immunol.* 1989; 19(8):1463-1467.
 - ❖ Varani S, Landini MP. Cytomegalovirus-induced immunopathology and its clinical consequences. *Herpesviridae.* 2011; 2:6.
 - ❖ Vargas-Rojas MI, Crispín JC, Richaud-Patin Y, Alcocer-Varela J. Quantitative and qualitative normal regulatory T cells are not capable of inducing suppression in SLE patients due to T-cell resistance. *Lupus.* 2008; 17(4):289-294.
 - ❖ Venigalla RK, Tretter T, Krienke S, Max R, Eckstein V, Blank N, Fiehn C, Ho AD, Lorenz HM. Reduced CD4+,CD25- T cell sensitivity to the suppressive function of CD4+,CD25high,CD127 -/low regulatory T cells in patients with active systemic lupus erythematosus. *Arthritis Rheum.* 2008; 58(7):2120-2130.
 - ❖ Vento S, Guella L, Mirandola F, Cainelli F, Di Perri G, Solbiati M, Ferraro T, Concia E. Epstein-Barr virus as a trigger for autoimmune hepatitis in susceptible individuals. *Lancet.* 1995; 346(8975):608-609.
 - ❖ Vial T, Nicolas B, Descotes J. Clinical immunotoxicity of pesticides. *J Toxicol Environ Health.* 1996; 48(3):215-229.

- ❖ Villadangos JA. Presentation of antigens by MHC class II molecules: getting the most out of them. *Mol Immunol.* 2001; 38(5):329-346.
- ❖ Vine MF, Stein L, Weigle K, Schroeder J, Degnan D, Tse CK, Backer L. Plasma 1,1-dichloro-2,2-bis(p-chlorophenyl) ethylene (DDE) levels and immune response. *Am J Epidemiol.* 2001; 153(1):53–63.
- ❖ von Schmiedeberg S, Fritsche E, Ronnau AC, Specker C, Golka K, Richter-Hintz D, Schuppe HC, Lehmann P, Ruzicka T, Esser C, Abel J, Gleichmann E. Polymorphisms of the xenobiotic-metabolizing enzymes CYP1A1 and NAT-2 in systemic sclerosis and lupus erythematosus. *Adv Exp Med Biol.* 1999; 455:147–152.
- ❖ Vrbikova J, Janatkova I, Zamrazil V, Tomiska F, Fucikova T. Epstein-Barr virus serology in patients with autoimmune thyroiditis. *Exp Clin Endocrinol Diabetes.* 1996; 104(1): 89–92.
- ❖ Walker LS, Abbas AK. The enemy within: keeping self-reactive T cells at bay in the periphery. *Nature Rev Immunol.* 2002; 2(1):11–19.
- ❖ Walsh P, Norris DA, Abe J, Martin DK, Giorno R, Leung DY. *Candida albicans* induces selective expansion of human T lymphocytes expressing the T-cell receptor variable region V β 5.1. *J Dermatol Sci.* 1996; 12(2):140–146.
- ❖ Wang B, Morinobu A, Kanagawa S, Nakamura T, Kawano S, Koshiba M, Hashimoto H, Kumagai S. Transforming Growth Factor Beta 1 Gene Polymorphism in Japanese Patients with Systemic Lupus Erythematosus. *Kobe J Med Sci.* 2007; 53(1-2):15-23.
- ❖ Wang DG, Fan JB, Siao CJ, Berno A, Young P, Sapolsky R, Ghandour G, Perkins N, Winchester E, Spencer J, Kruglyak L, Stein L, Hsie L, Topaloglou T, Hubbell E, Robinson E, Mittmann M, Morris MS, Shen N, Kilburn D, Rioux J, Nusbaum C, Rozen S, Hudson TJ, Lipshutz R, Chee M, Lander ES. Large-scale identification, mapping, and genotyping of single-nucleotide polymorphisms in the human genome. *Science.* 1998; 280(5366):1077–1082.
- ❖ Wang J, Ioan-Facsinay A, van der Voort EI, Huizinga TW, Toes RE. Transient expression of FOXP3 in human activated non-regulatory CD4⁺ T-cells. *Eur J Immunol.* 2007; 37(1):129–138.

- ❖ Wardemann H, Yurasov S, Schaefer A, Young JW, Meffre E, Nussenzweig MC. Predominant autoantibody production by early human B cell precursors. *Science*. 2003; 301(5638):1374–1377.
- ❖ Waxman J, Locksmith MD, Schnapp JJ, Doneson IN. Cellular immunity in rheumatic disease. *Arthritis Rheumtol*. 1973; 16(4):499-506.
- ❖ Weiner HL. Oral tolerance: immune mechanisms and the generation of Th3-type TGF-beta-secreting regulatory cells. *Micorobes Infect*. 2001; 3(11):947-954.
- ❖ Weiss A, Imboden J, Hardy K, Manager B, Terhorst C, Stobo J. The role of the T3/antigen receptor complex in T-cell activation. *Annu Rev Immunol*. 1986; 4:593–619.
- ❖ Wen AQ, Gu W, Wang J, Feng K, Qin L, Ying C, Zhu PF, Wang ZG, Jiang JX. Clinical relevance of IL- 1 β promoter polymorphisms (-1470,-511, and-31) in patients with major trauma. *Shock*. 2010; 33(6):576–582.
- ❖ Weninger W, Manjunath N, von Andrian UH. Migration and differentiation of CD8+ T cells. *Immunol Rev*. 2002; 186:221-233.
- ❖ White J, Herman A, Pullen A, Kubo R, Kappler JW, Marrack P. The V β -specific superantigen staphylococcal enterotoxin B: Stimulation of mature T-cells and clonal deletion in neonatal mice. *Cell*. 1989; 56(1):27-35.
- ❖ Whiteside TL, Kumagai Y, Roumm AD, Almendinger R, Rodnan GP. Suppressor cell function and T lymphocyte subpopulations in peripheral blood of patients with progressive systemic sclerosis. *Arthritis Rheum*. 1983; 26(7):841-847.
- ❖ Williams A, Peh CA, Elliott T. The cell biology of MHC class I antigen presentation. *Tissue Antigens*. 2002; 59(1):3-17.
- ❖ Wohl Y, Brenner S. Pemphigus in Israel: An Epidemiologic Analysis of Cases in Search of Risk Factors. *IMAJ*. 2003; 5(6):410-412.
- ❖ Wohl Y, Goldberg I, Shirazi I, Brenner S. Chlorpyrifos exacerbating pemphigus vulgaris: a preliminary report and suggested in-vitro immunologic evaluation model. *SKINmed*. 2006; 5(3):111–113.
- ❖ Woosley RL, Drayer DE, Reidenberg MM, Nies AS, Carr K, Oates JA. Effect of acetylator phenotype on the rate at which procainamide induces antinuclear antibodies and the lupus syndrome. *N Engl J Med*. 1978; 298(21):1157–1159.

-
- ❖ Wu K, Bi Y, Sun K, Wang C. IL-10-producing type 1 regulatory T cells and allergy. *Cell Mol Immunol*. 2007; 4(4):269–275.
 - ❖ Wucherpfennig KW, Weiner HL, Hafler DA. T-cell recognition of myelin basic protein. *Immunol Today*. 1991; 12(8):277–282.
 - ❖ Wucherpfennig KW. Mechanisms for the induction of autoimmunity by infectious agents. *J Clin Invest*. 2008; 108(8):1097–1104.
 - ❖ Wucherpfennig KW. Structural basis of molecular mimicry. *J Autoimmun*. 2001; 16(3):293–302.
 - ❖ Wynn TA. Fibrotic disease and the T (H) 1 / T (H) 2 paradigm. *Nat Rev Immunol*. 2004; 4(8):583–594.
 - ❖ Xaubet A, Marin-Arguedas A, Lario S, Transforming growth factor beta 1 gene polymorphisms are associated with disease progression in idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med*. 2003; 168(4):431–435.
 - ❖ Yanez L, Borja-Aburto VH, Rojas E, de la Fuente H, Gonzalez-Amaro R, Gomez H, Jongitud AA, Diaz-Barriga F. DDT induces DNA damage in blood cells. Studies in vitro and in women chronically exposed to this insecticide. *Environ Res*. 2004; 94(1):18–24.
 - ❖ Yang X, Buonpane RA, Moza B, Rahman AK, Wang N, Schlievert PM, McCormick JK, Sundberg EJ, Kranz DM. Neutralization of multiple staphylococcal superantigens by a single-chain protein consisting of affinity-matured, variable domain repeats. *J Infect Dis*. 2008; 198(3):344–348.
 - ❖ Yeh SW, Sami N, Ahmed AR. Treatment of pemphigus vulgaris: current and emerging options. *Am J Clin Dermatol*. 2005; 6(5):327-342.
 - ❖ Yewdell J, Antón LC, Bacik I, Schubert U, Snyder HL, Bennink JR. Generating MHC class I ligands from viral gene products. *Immunol Rev*. 1999; 172:97-108.
 - ❖ Young V, Ho M, Vosper H, Belch JJ, Palmer CN. Elevated expression of the genes encoding TNF-alpha and thromboxane synthase in leukocytes from patients with systemic sclerosis. *Rheumatology (Oxford)*. 2002; 41(8):869-875.
 - ❖ Yu CR, Mahdi RM, Ebong S, Vistica BP, Gery I, Egwuagu CE. Suppressor of cytokine signaling 3 regulates proliferation and activation of T-helper cells. *J Biol Chem*. 2003; 278(32):29752–29759.

-
- ❖ Yu J, Zhang M, Wang JL, Xu WL, Ou JP, Wang FR, Lai YY, Liu F, Ren HY, Huang XJ, Song YX, An YZ, Li RY. *Candida* species distribution in the patients with high risk of deep fungal infections and relevant risk factors: a prospective cohort study. *Zhonghua Yi Xue Za Zhi*. 2007; 87(42):2991-2993.
 - ❖ Zamoyska R. CD4 and CD8: modulators of T cell receptor recognition of antigen and of immune responses? *Curr Opin Immunol*. 1998; 10(1):82-87.
 - ❖ Zandman-Goddard G, Tweezer-Zaks N, Shoenfeld Y. New therapeutic strategies for systemic sclerosis - a critical analysis of the literature. *Clin Dev Immunol*. 2005; 12(3):165-173.
 - ❖ Zerrahn J, Held W, Raulet DH. The MHC reactivity of the T cell repertoire prior to positive and negative selection. *Cell*. 1997; 88(5):627-636.
 - ❖ Zhang B, Zhang X, Tang F, Zhu L, Liu Y. Reduction of forkhead box P3 levels in CD4⁺CD25^{high} T cells in patients with new-onset systemic lupus erythematosus. *Clin Exp Immunol*. 2008; 153(2):182-187.
 - ❖ Zheng SG, Wang JH, Gray JD, Soucier H, Horwitz DA. Natural and induced CD4⁺CD25⁺ cells educate CD4⁺CD25⁻ cells to develop suppressive activity: The role of IL-2, TGF-beta, and IL-10. *J Immunol*. 2004; 172(9):5213-5221.
 - ❖ Zhou X, Tan FK, Stivers DN, Arnett FC. Microsatellites and intragenic polymorphisms of transforming growth factor beta and platelet-derived growth factor and their receptor genes in native Americans with systemic sclerosis (scleroderma). *Arthritis Rheum*. 2000; 43(5):1068-1073.
 - ❖ Zhou Y, Gsicombe R, Huang D, Lefvert AK. Novel genetic association of Wegener's granulomatosis with the interleukin-10 gene. *J Rheumatol*. 2002; 29(2):317-320.
 - ❖ Zhu H, Cong JP, Yu D, Bresnahan WA, Shenk TE. Inhibition of cyclooxygenase 2 blocks human cytomegalovirus replication. *Proc Natl Acad Sci USA*. 2002; 99(6):3932-3937.
 - ❖ Zhu J, Paul WE. CD4 T cells: fates, functions, and faults. *Blood*. 2008; 112(5):1557-1569.
 - ❖ Zinkernagel RM, Doherty PC. Restriction of in vitro T cell-mediated cytotoxicity in lymphocytic choriomeningitis within a syngeneic or semiallogeneic system. *Nature*. 1974; 248:701-702.