
REFERENCES

REFERENCES

Aghajari, N., Feller, G., Gerday, C. and Haser, R. (1998b). Structures of the psychrophilic *Alteromonas haloplanctis* alpha-amylase give insights into cold adaptation at a molecular level. *Structure* 6, 1503–1516.

Aghajari, N., Feller, G., Gerday, C., Haser, R., (1998a). Crystal structures of the psychrophilic alpha-amylase from *Alteromonas* alpha-amylases in its native form and (Erratum Prot. Sci. 1998 Jun 7(6), p. 1481). *Prot. Sci.* 7, 564–572.

Alain Ferrer, Johan Hoebeke and Daniel Bout.(1999). Purification and characterization of two α -Amylases from *Toxoplasma gondii*. *Eperimental Parasitology*. 92: 64-72.

Arikan Burhan, Unaldi Nisa, Coral Gokhan, Colak Omer, Aygan Ashabil and Gulnaz Osman. (2003). Enzymatic properties of a novel thermostable, thermophilic, alkaline and chelator resistant amylase from an alkaliphilic *Bacillus sp.* isolate ANT-6. *Process Biochemistry*. 38: 1397-1403.

Armstrong, W.P. (1982). Beware of the Jumping Cholla. *Environment Southwest*. 498: 3-5.

Armstrong, W.P. (1992). Devil Claws. *Pacific Horticulture*. 53: 19-23.

Bairoch, A. and Apweiler, R. (2000). The SWISS-PROT protein sequence database and its Supplement TrEMBL in 2000. *Nucleic Acids Res.* 28: 45–48.

Banks, W. and Greenwood, C.T. (1975). Starch degrading enzymes. In: *Starch and its Components* Edinburger, University Press. 191-241.

Barbera, G., Carimi, F. and Inglese, P. (1992). Past and present role of the Indian-fig prickly pear (*Opuntia ficus-indica* (L.) Miller, Cactaceae) in the agriculture of Sicily. *Economic Botany.* 46: 10-22.

Barthlott, W. (1977). (English Trans. L. Glass). *Cacti.* Cheltenham: Stanley Thornes Ltd.

Ben, A.M., Mezghani, M. and Bejar, S.A. (1999). Thermostable alpha-amylase producing maltohexaose from a new isolated *Bacillus* sp. US100: study of activity and molecular cloning of the corresponding gene. *Enzyme Microb Technol.* 24: 584-9.

Benson, D.A., Karsch-Mizrachi, I., Lipman, D.J., Ostell, J., Rapp, B.A. and Wheeler, D.L. (2002). GenBank. *Nucleic Acids Res.* 30: 17–20.

Benson, L. (1982). *The Cacti of the United States and Canada.* Stanford, Ca: Stanford University Press.

Bergquist, P.L. and Morgan, H.W. (1992). The molecular genetics and biotechnological application of enzymes from extremely thermophilic eubacteria. In R.A. Herbert and R.J. Sharp (ed.), *Molecular biology and biotechnology of extremophiles.* Chapman & Hall, New York, N. Y. 44-75.

Berman, H.M., Battistuz, T., Bhat, T.N., Bluhm, W.F., Bourne, P.E., Burkhardt, K., Feng, Z., Gilliland, G.L., Iype, L., Jain, S., Fagan, P., Marvin, J., Padilla, D., Ravichandran, V., Schneider, B., Thanki, N.,

Weissig, H., Westbrook, J.D. and Zardecki, C. (2000). The protein data bank. *Acta Crystallogr.* D58, 899-907.

Bertoldo, C. and Antranikian, G. (2001). Amylolytic enzymes from hyperthermophiles. *Methods Enzymol.* 330: 269-289.

Boel, E., Brady, L., Brzozowski, A.M., Derewenda, Z., Dodson, G.G., Jensen, V.J., Petersen, S.B., Swift, H., Thim, L. and Woldike, H.F. (1990). Calcium binding in α -amylases: an X-ray diffraction study at 2.1-Å resolution of two enzymes from *Aspergillus*. *Biochemistry.* 29: 6244-6249.

Boissenvain Charles, H. and Davidson Carol. (1940). Colorado Cacti. Abbey Garden Press, Pasadena, California.

Bragger, J.M., Daniel, R.M., Coolbear, T. and Morgan, H.W. (1989). Very stable enzymes from extremely thermophilic archae bacteria and eubacteria. *Appl. Microbiol. Biotechnol.* 31: 556-561.

Bronnenmeir, K., Kern, A., Liebl, W. and Standenbauer, W.L. (1995). Purification of *Thermotoga maritima* enzymes for the degradation of cellulosic materials. *Appl. Environ. Microbiol.* 61: 1399-1407.

Brown, S.H. and Kelly, R.M. (1993). Characterization of amylolytic enzymes, having both α -1,4 and α -1,6 hydrolytic activity, from the thermophilic archaea *Pyrococcus furiosus* and *Thermococcus litoralis*. *Appl. Environ. Microbiol.* 59: 2614-2621.

Brutsche, M.O. (1984). Prickly pear (*Opuntia ficus-indica*) cultivation in southern Africa. In: Symposium of agricultural use of the Cactaceae. Prospects and problems. International organization for succulent plant study, Frankfurt.

Buchanan, C.L., Connaris, H., Danson, M.J., Reeve, C.D. and Hough, D.W. (1999). An extremely thermostable aldolase from *Sulfolobus solfataricus* with specificity for non-phosphorylated substrates. *Biochem. J.* 343: 563-570.

Campus, B.G., Priest, F.G. and Stark, J.R. (1992). Hydrolysis of starch granules by the amylase from *Bacillus Stearothermophilus* NCA 26. *Proc. Biochem* 27: 17-21.

Canganella, F., Andrade, C.M. and Antranikian, G. (1994). Characterization of amylolytic and pullulytic enzymes from thermophilic archaea and from a new Fevidobacterium species. *Appl. Microbiol. Biotechnol.* 42: 239-245.

Chen, W., Avison, M.J., Zhu, X.H. and Shulman, R.G. (1993). *Biochemistry.* 32: 11483-11487.

Chidamian, C. (1958). The Book of Cacti and Other Succulents. New York: The American Garden Guild and Doubleday Company Inc.

Chomczynski, P. and Sacchi, N. (1987). Single step method of RNA isolation by acid guanidium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* 162: 156-159.

Clairf Vieille, Gregory I. and Zeikus. (2001). Hyperthermophilic Enzymes: Sources, Uses and Molecular Mechanisms for Thermostability. *Microbiology and Molecular Biology.* 65: 1-43.

Clovis de Andrade, J. (1990). As palmas forrageiras em Alagoas, Maceio, Brazil. 181.

Colombo, S., D' Auria, S., Fusi, P., Zecca, L., Raia, C.A. and Tortora, P. (1992). Purification and characterization of a thermostable carboxypeptidase from the extreme thermophilic archaeobacterium *Sulfolobus solfataricus*. *Eur. J. Biochem.* 206: 349-357.

Crueger, W. and Crueger, A. (eds.) (1989). Industrial Microbiology, Sinauer Associates, Sunderland, MA. pp. 189–218

Cullmann, W., Gotz, D. and Groner, G. (1986). The Encyclopedia of Cacti. Portland, OR: Timber Press.

Curtis, R.J. (1979). Prickly pear farming in the Santa Clara Valley, California. *Economic Botany*.

Da Lage, J.-L., Maczkowiak, F. and Cariou, M.-L. (2000). Molecular characterization and evolution of the amylase multigene family of *Drosophila ananassae*. *J. Mol. Evol.* 51: 391-403.

Daniel Y. Sellos and Alain Van Wormhoudt. (2002). Structure of the of alpha amylase genes in crustaceans and molluscs: evolution of the exon/intron organization. *Biologia*. Bratislava. 11: 191-196.

Davies, G.J., Gamblin, S.J., Littlechild, J.A. and Watson, H.C. (1993). The structure of a thermally stable 3-phosphoglycerate kinase and a comparison with its mesophilic equivalent. *Proteins: Struct. Funct. Genet.*, 15: 283–289

De Kock, G.C. (1965). Manejo utilizacion del nopal sin espinas. En: Congreso Internacional de Pasturas. 9o Anales Sao Paulo, Brazil, 2: 1471.

De Mot, R. and Verachtert, H. (1986). Secretion of alpha-amylase and multiple forms of glucomylase by the yeast *Trichosporon pullulans*. *Can. J. Microbiol.* 32: 47-51.

De Mot, R. and Verachtert, H. (1987). Purification and Characterization of extracellular alpha-amylase and glucoamylase from the yeast *Candida antarctica* CBS 6678. *Eur. J. Biochem.* 164: 643-654.

Dong, G. and Zeikus, J.G. (1997). Purification and characterization of alkaline phosphatase from *Thermotoga neapolitana*. *Enzyme Microb. Technol.* 21: 335-340.

Egelseer, E., Schocher, I., Sleytr, U.B. and Sara, M. (1996). Evidence that an N-terminal S-layer protein fragment triggers the release of a cell associated high-molecular-weight-amylase in *Bacillus stearothermophilus* ATCC 12980. *J. Bacteriol.* 178: 5602-9.

Farez-Vidal, M.E., Fernandez-Vivas, A., Gonzaler, F. and Arias, J. M. (1995). *Appl. Bacteriol.* 78: 14-19.

Fensome, R.A. and Williams, G.L. (2001). The Last Billion Years. Atlantic Geoscience Society: Halifax, Nova Scotia: Nimbus.

Fitter, J., Herrmann, R., HauB, T., Lechner, R.E. and Dencher, N.A. (2001). Dynamical properties of α -amylase in the folded and unfolded state: the role of thermal equilibrium fluctuations for conformational entropy and protein stabilization. *Physica B* 301: 1-2.

French, D. (1973). *J. Animal Sci.* 37: 1048-1061.

French, D. (1975). International review of Biochemistry, (Whelan, W.J., ed), Univ. Park Press, Baltimore, Maryland (Series one). 5: 269.

Friedrich, A.B., and Antranik, G. (1996). Keratin degradation by *Fervidobacterium pennavorans*, a novel thermophilic anaerobic species of the order Thermotogales. *Appl. Environ. Microbiol.* 62: 2875-2882.

Gabelsberger, J., Liebl, W. and Schleifer, K.H. (1993). Cloning and characterization of β -glycoside hydrolyzing enzymes of *Thermotoga maritime*. *FEMS Microbiol. Lett.* 109: 131-138.

Gashaw Mamo and Amare Gessesse. (1999). Purification and characterization of two raw-starch-digesting thermostable α -amylases from a thermophilic *Bacillus*. *Enzyme and Microbial Technology*. 25: 433-438.

Gomes, J., and Steiner, W. (1998). Production of a high activity of an extremely thermostable β -mannanase by the thermophilic eubacterium *Rhodothermus marinus*, grown on locust bean gum. *Biotechnol. Lett.* 20: 729-733.

Grossman, G.L., Campos, Y., Severson, D.W. and James, A.A. (1997). *Insect Biochem. Mol. Biol.* 27: 769-781.

Galley, J.A. and Edelman, G.M (1964). *Biopolymers Symp.* 1: 367-381.

Henry, R. J., Chiamori, N. (1960). *Clin. Chem.* 6: 434.

Hoffman, G.O. and Walker, A.H. (1912). Prickly pear – Good or bad? Texas Agriculture Experiment Station. 806: 3-7.

Horikoshi, K. (1998). Alkaliphiles. In *Extremophiles – Microbial Life in Extreme Environments*, Edited by Horikoshi, K., Grant, W.D. New York: Wiley-Liss. 155-179.

Hsiu, J., Fischer, E. H. and Stein, E. A. (1964). *Biochemistry.* 3: 61–66

Hybl, A., Rundle, R.E. and Williams, D.E. (1965). *J. Am. Chem. Soc.* 87: 2779-2788.

Hyun, H.H. and Zeikus, J.G. (1985). General Biochemical characterization of thermostable extracellular beta-amylase from *Clostridium thermosulfurogenes*. *Appl. Environ. Microbiol.* 49: 1162-1167.

Ikeda, M., and Clark, D.S. (1998). Molecular cloning of extremely thermostable esterase gene from hyperthermophilic archaeon *Pyrococcus furiosus* in *Escherichia coli*. *Biotechnol. Bioeng.* 57: 624-629.

Ishikawa, K., Nakamura, H., Morikawa, K. and Kanaya, S. (1993). Stabilization of *Escherichia coli* ribonuclease HI by cavity-filling mutations within a hydrophobic core. *Biochemistry.* 32: 6171–6178.

Ito, S., Kobayashi, T., Ara, K., Ozaki, K., Kawai, S. and Hatada, Y. (1998). Alkaline detergent enzymes from alkaliphiles: Enzymatic properties, genetics and structures. *Extremophilies.* 2: 185-190.

Janecek, S. (1994b). Sequences similarities and evolutionary relationships of microbial, plant and animal α -amylases. *Eur. J. Biochem.* 224: 519-524.

Jean-Luc Da Lage, Alain Van Wormhoudt and Marie-Louise Cariou (2002). Diversity and evolution of the alpha-amylase genes in animals. *Biologia. Bratislava.* 11: 181-189.

Jean-Luc Da Lage, Alain Van Wormhoudt and Marie-Louise Cariou. (2002). Diversity and evolution of the amylase genes in animals. *Biologia. Bratislava.* 11: 181-189.

Jei-Fu Shaw, Fu-Pang Lin, Su-Chiu Chen and Hsing-Chen Chen. (1995). Purification and properties of an extra cellular α -amylase from *Thermus sp.* *Bot. Bull. Acad. Sin.* 36: 195-200.

Jensen, G. (1990). Tenside Superf. Deterg. 27- 30.

Jacobsen, H. (1954c). A Handbook of Succulent Plants. Poole, Vol. Dorset: Blandford Press

Karshikoff, A. and Ladenstein, R. (1998). Proteins from thermophilic and mesophilic organisms essentially do not differ in packing. *Protein Eng.* 11: 867–872.

Kashiwabara, S., Ogawa, S., Miyoshi, N., Oda, M. and Suzuki, Y. (1999). Three domains comprised in thermostable molecular weight 54,000 pullulanase of type I from *Bacillus flavocaldarius* KP1228. *Biosci. Biotechnol. Biochem.* 63: 1736-1748.

Kelly, C.T., McTigue, M.A., Doyle, E.M. and Fogarty, W.M. (1995). The raw starch-degrading alkaline amylases of *Bacillus sp* IMD 370. *J Ind. Microbiol.* 15: 446-8.

Kengen, A.W., Luesink, E.V., Stams A.J. and Zehnder, A.J. (1993). Purification and characterization of an extremely thermostable β -glucosidase from the hyperthermophilic archaeon *Pyrococcus furiosus*. *Eur. J. Biochem.* 213: 305-12.

Kenneth, A., Laderman Bradley., Devie, R., Henry C. Krutzach., Mars, S., Lewis, Y.V. Griko., Peter L. Privalov and Christian B. Anfineen. (1993). The Purification and Characterization of an Extremely Thermostable α -Amylase from the Hyperthermophilic *Archaeobacterium Pyrococcus furiosus*. *The Journal of Biological Chemistry.* 32: 24398-24401.

Kenya J. Wanderlay, Fernando A.G. Torres, Lidia M.P. Moraes, and Cirano J. Ulhoa. (2004). Biochemical characterization of α -amylase from the yeast *Cryptococcus flavus*. *FEMS. Microbiology Letters.* 231: 165-169.

Kim, Y.O., Kim, H.-K., Bae, K.-S., Yu, J.-H. and Oh, T.K. (1998). Purification and properties of a thermostable phytase from *Bacillus sp.* DS11. *Enzyme Microb. Technol.* 22: 2-7.

Koch, R., Zabłowski, P., Spreinat, A. and Antranikian, G. (1990). Extremely thermostable amyolytic enzymes from archaebacterium *Pyrococcus furiosus*. *FEMS Microbiol. Lett.* 71: 21-26.

Koh, S., Shin, H.J., Kim, J.S., Lee, D.S. and Lee, S.Y. (1998). Trehalose synthesis from maltose by a thermostable trehalose synthase from *Thermus caldophilus*. *Biotechnol. Lett.* 20: 757-761.

Kozianowski, G., Canganella, F., Rainey, F.A., Hippe, H. and Antranikian, G. (1997). Purification and characterization of thermostable pectate-lyases from a newly isolated thermophilic bacterium, *Thermoanaerobacter italicus* sp. Nov. *Extremophiles* 1: 171-182.

Kristjanson, K.J. (1989). Thermophilic organism as source of thermostable enzymes. *Trends in Biotechnol.* 7: 49-53.

Kristjansson, J.K. and Hreggvidsson, G.O. (1995). Ecology and habitats of Extremophiles. *World J Microbiol Biotechnol.* 11: 17-25.

Krystyna Zoltowska. (2001). Purification and characterization of α -amylases from the intestine and muscle of *Ascaris suum* (Nematoda). *Acta. Biochimica. Polonica.* 48: 763-774.

Laemmli, U.K. (1970). Cleavage of structural protein during the assembly of the head of bacteriophage T4. *Nature.* 227: 680-685.

Larson, S.B., Greenwood, A., Cascio, D., Day, J. and McPherson, A. (1994). Refined molecular structure of pig pancreatic alpha-amylase at 2.1 Å^o resolution. *J. Mol. Biol.* 235: 1560-1584.

Le Houerou, H.N. (1979). Resources and potential of the native flora for fodder and sown pastures production in the arid and semi-arid zones of North Africa. In Goodin JR and Worthington DK, eds, *Arid Land plant Resources*. ICASALS, Texas Tech Univ. Lubbock, Texas. 384-401.

Lee, C.Y., and Zeikus, J.G. (1991). Purification and characterization of thermostable glucose isomerase from *Clostridium thermosulfurogens* and *Thermoanaerobacter strain B6A*. *Biochem. J.* 273: 565-571.

Lee, S.G., Lee, D.C., Hong, S.P., Sung, M.H. and Kim., H.S. (1995). Thermostable D-hydantoinase from thermophilic *Bacillus stearothermophilus* DS-1: characteristics of purified enzyme. *Appl. Microbiol. Biotechnol.* 43: 270-276.

Lefuji, H., Chino, M., Kato, M. and Imura, Y. (1996). Raw-starch-digesting and thermostable alpha-amylase from the yeast *Cryptococcus* sp. S-2: purification, characterization, cloning and sequencing. *Biochem. J.* 318: 989-996.

Leibl, W., Ruile, P., Bronnenmeier, K., Riedel, K., Lottspeich, F. and Greif, L. (1996). Analysis of a *Thermotoga maritime* DNA fragment encoding two similar thermostable cellulases, CetA and CetB, and characterization of the recombinant enzymes. *Microbiology.* 142: 2533-2542.

Leveque, E., Janecek, S., Haye, B. and Belarbi, A. (2000). Thermophilic archaeal amylolytic enzymes. *Enzyme Microb. Technol.* 26: 3-14.

Liebl, W., Wangner, B. and Schellhase, J. (1998). Properties of an α -galactosidase, and structure of its gene gal4, within an α and β -galactoside utilization gene cluster of the hyperthermophilic bacterium *Thermotoga maritime*. *Syst. Appl. Microbiol.* 21: 1-11.

Linke, B., Rudiger, A., Wittenberg, G., Jorgensen, P.L. and Antraikian, G. (1992). Production of heat-stable pullulanase and alpha-glucosidase from the extremely thermophilic archaeon *Pyrococcus woesei*. *DEHEMA Biotechnol. Conf.* 161-163.

Lomako, J., Lomako, W.M., Whelan, W.J., Domdro, R.S., Neary, J.T. and Norenberg, M.D. (1993). Glycogen synthesis in the astrocyte: from glycogenin to proglycogen to glycogen. *FASEB J.* 7: 1386-1393.

Long-Liu Lin, Charng-Cherng Chyau and Wen-Hwei Hsu. (1998). Production and properties of a rawstarch-degrading amylase from the thermophilic and alkaliphilic *Bacillus sp* TS- 23. *Biotechnol Appl. Biochem.* 28: 62-68.

Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J. (1951). Protein measurement with the folin-phenol reagent. *J. Biol. Chem.* 193: 265-275.

Lozano, G.M. (1958). Contribucion al estudio e industrializacion del nopal (*Opuntia sp.*). Tesis profesional. Saltillo, Esc. Sup. de Agric. Antonio Narro.

Lohf, Lin L.L., Chen, H.L., Hsu, H.H. and Chang, C.T. (2001). Enzymatic properties of a SDS-resistant *Bacillus sp.* TS-23 α - amylase produced by recombinant *E.Coli*. *Process Biochem.* 36: 743-50.

Merkler, D. J., Srikumar, k., Marchese Ragona, S .P. and Wedler, F .C. (1988). Aggregation of thermo-inactivation of glutamine synthetase from an extreme thermophile, *Bacillus caldolyticus*. *Biochimica et Biophysica Acta.* 952: 101-114.

Madeleine, M.L., Libemart, N., Bellanger, F., Delrue, B., D' Hulst, C., Van den Koorhuysse, N., Fontaine, T., Wieruszkeski, J.M., Decq, A. and Ball, S. (1994). Toward an understanding of the biogenesis of the starch granule. Determination of granule-bound and soluble starch synthase functions in amylopectin synthesis. *J. Biol. Chem.* 269: 25150-25157.

Madi, E., Antranikian, G., Ohmiya, K. and Gottschalk, G. (1987). Thermostable amyolytic enzymes from a new *Clostridium* isolate. *Appl. Environ. Microbiol.* 53: 1661-1667.

Marchal, L.M., Jonkers, J., Franke, G.T., de Gooijer, C.D. and Tramper, J. (1999). The effect of process conditions on the alpha-amylolytic hydrolysis of amylopectin potato starch: An experimental design approach. *Biotechnol. Bioeng.* 62: 348-357.

Matsuura, Y., Kusunoki, M., Harada, W., Tanaka, N., Iga, Y., Yasuoka, N., Toda, H., Narita, K. and Kakudo, M. (1980). Molecular structure of Taka-amylase A. I. Backbone chain folding at 3-A resolution. *J. Biochem.* 87. 5581.

Matthews, B.W. (1987). Genetic and structural analysis of the protein stability problem. *Biochemistry.* 26: 6885–6888.

Matthews, B.W. (1993). Structural and genetic analysis of protein stability. *Annu. Rev. Biochem.* 62: 139–160.

Mc Gregor, E.A., Janecek, S. and Svensson. B. (2001). Relationship of sequence and structure to specify in the alpha-amylase family. *Biochim. Biophys. Acta.* 1546: 1-20.

McTigue, M.A., Kelly, C.T., Doyle, E.M. and Fogarty, W.M. (1995). The alkaline amylase of the alkaliphilic *Bacillus* sp. IMD 370. *Enzyme. Microb. Technol.* 17: 570-73.

Menéndez-Arias, L and Argos, P. (1989). Engineering protein thermal stability: sequence statistics point to residue substitutions in alpha-helices. *J. Mol. Biol.* 206: 497–406.

Mishra, R.S. and Maheshwari, R. (1996). Amylases of thermophilic fungus *Thermomyces lanuginosus*: their purification, properties, action on starch and response to heat. *J Biosci.* 21: 653-672.

Moal, J., Daniel, J.Y., Sellos, D., Van Wormhoudt, A. and Samain, J.F. (2000). Amylase mRNA expression in *Crassostrea gigas* during feeding cycles. *J. Comp. Physiol.* 170: 21-26.

Monma, M., Kagi, N. and Kainuma, K. (1989). Raw starch digestion by *Chalara paradoxica* immobilized in calcium alginate cells. *Starch*. 41: 355-357.

Moranelli, F., Yaguchi, M., Calleja, B. and Nasim, A. (1982). Purification and characterization of the extracellular alpha-amylase activity of the yeast *Schwanniomyces alluvius*. *Biochem. Cell Biol.* 65: 899-907.

Morita, H. and Fujio, Y. (1997). High specific activity of raw starch digesting glucoamylase producing *Rhizopus sp.* A-11 in liquid culture. *Starch*. 49:293-6.

Morrison, F.B. (1956). Compendio de la alimentacion del Ganado. Mexico. UTEHA. 721.

Murai, T., Ueda, M., Shibasaki Y., Kamasawa, N., Osumi, M., Imanaka, T. and Tanaka, A. (1999). Development of an arming yeast strain for efficient utilization of starch by co-display of sequential amylolytic enzymes on the cell surface. *Appl. Microbiol Biotechnol.* 51: 65-70.

Nagasaka, Y., Kurosawa, K., Yokota, A. and Tomita, F. (1998). Purification and properties of the raw-starch-digesting glucoamylases from *Corticium rolfsii*. *Appl. Microbiol. Biotechnol.* 50: 323-330.

Nakajima, R., Imanaka, T. and Aiba, S. (1986). Comparison of amino acid sequences of eleven different α -amylases. *Appl. Microbiol. Biotechnol.* 23, 355-360.

Nakamura, N., Sashihara, N., Nagayama, H. and Horikoshi. K. (1987). Production of a thermostable pullulanase by a *Thermus sp.* *J. Jpn. Soc. Starch Sci.* 34: 38-44.

Niehaus, F., Bertoldo, C., Kahler, M. and Antranikian, G. (1999). Extremeophiles as a source of novel enzymes for industrial application. *Appl. Microbiol. Biotechnol.* 51: 711-29.

Nobel, P.S. (1989a). A nutrient index quantifying productivity of agaves and cacti. *J. Appl. Ecol.* 26: 635-645.

Nicholas Walsh, J. (1989). Handbook of inductively coupled plasma spectrometry. Blackie, Glasow and London. Published by Chapman and Hall, New York. USA.

Odile Mayzaud. (1985). Purification and kinetic properties of the α -amylase from the copepod *Acartia clause* *Comp. Biochem. Physiol.* 82B: 725-730.

Okolo, B.N., Ezeogu, L.I. and Ebisike, C.O. (1996). Raw starch digesting amylase from *Thermoactinomyces thalpopophilus* F13. *World. J. Microbiol. Biotechnol.* 12: 637-8.

Ouchterlony, O., (1958). *Prog. Allergy* 5:1.

Pace, C.N. and Scholtz, J.M. (1989). T.E. Creighton Ed., Protein structure: A Practical Approach, IRL, Press, Oxford, p. 299.

Peggy Winter. (1980). From the Mailbag. 1: 15-16.

Pimienta-Barrios, E. (1990). El nopal tunero. Universidad de Guadalajara, Mexico.

Plant, A.R., Morgan, H.W. and Daniel, R.M. (1986). A highly stable pullulanase from *Thermus aquaticus* YT-1. *Enzym Microb. Technol.* 8: 668-672.

Prieto, J.A., Bort, B.R., Martinez, J., Randez-Gil F., Buesa, C. and Sanz, P. (1995). Purification and characterization of a new alpha-amylase of

intermediate thermal stability from the yeast *Lipomyces kononenkoae*. *Biochem. Cell. Bio.* 73: 41-49.

Proctor, R.C. and Clair Meyer. (1954). The flowering Cactus. Mc Graw-Hill Book Co., Inc., New York.

Quang D, Nguyen Judit, M., Rezessy-Szabo, Marc Claeysens, Ingeborg Stals and Agoston Hoschke. (2002). Purification and characterization of amylolytic enzymes from thermophilic fungus *Thermomyces lanuginosus* strain ATCC 34626. *Enzyme and Microbiol Technology.* 31: 345-352.

Rao, M.B., Tanksale, A.M., Gathe, M.S. and Deshpande, V.V. (1998). Molecular and biotechnological aspects of microbial proteases. *Microbiol. Mol. Biol. Rev.* 62: 597-635.

Rao, V.S.R. and Sathyanarayana, B.K. (1972). *Biopolymers.* 11: 1379-1394

Ratanakhanokchai, K., Kaneko, J., Kamio, Y. and Izaki, K. (1992). Purification and properties of a maltotetraose and maltotriose producing amylase from *Chloroflexus aurantiacus*. *Appl Environ Microb.* 58: 2490-2.

Riha, J. and Subik, R. (1981). The Illustrated Encyclopedia of Cacti and Other Succulents. Octopus Books Ltd, London.

Rychlik, W., Spencer, W. J. and Rhoads, R.E. (1990). Optimization of the annealing temperature for DNA amplification *in vitro*. *Nucleic Acids Research.* 18: 6409-12.

Rundle, R.E. and French, D. (1943). *J. Am. Chem. Soc.* 65: 1707-1710.

Russell, C.E. and Felker, P. (1987a). The Prickly pear (*Opuntia sp.*, Cactaceae): a source of human and animal food in semi-arid regions. *Econ. Bot.* 41: 433-455.

Ruttersmith, L.D., and Daniel. R.M. (1993). Thermostable β -glucosidase and β -xylosidase from *Thermotoga sp.* strain FjSS3-B.1. *Biochim. Biophys. Acta.* 1156: 167-172.

Saenz, C. (1985). La tuna (*Opuntia ficus-indica*) un cultivo con perspectives. *Alimentos.* 3: 47-49.

Saha, B., and Zeikus. J.G. (1991). Characterization of thermostable α -glucosidase from *Clostridium thermohydrosulfuricum* 39E. *Appl. Microbiol. Biotechnol.* 35: 568-571.

Saha, B.C., Lamed, R., Lee, C.Y., Mathupala, S.P. and Zeikus, G. (1990). Characterization of an endo - acting amylopullulanase from *Thermoanaerobacter Strain B6A.* *Appl. Environ. Microbiol.* 56: 881-886.

Salvino D'Amico, Charles Gerday and Georges Feller. (2000). Structural similarities and evolutionary relationships in chloride-dependent α -amylases. *Gene.* 253: 95-105.

Sandberg, W.S. and Terwillige, T.C. (1989). Influence of interior packing and hydrophobicity on the stability of a protein. *Science.* 245: 54-57.

Saul, D.S., Williams, L.C., Reeves, R.A., Gibbs, M.D. and Bergquist, P.I. (1995). Sequence and expression of a xylanase gene from the hyperthermophile *Thermotoga sp.* strain FjSS3-B.1 and characterization of the recombinant enzyme and its activity on kraft pulp. *Appl. Environm. Microbiol.* 61: 4110-4113.

Schink, B., and Zeikus, J.G. (1983). Characterization of pectinolytic enzymes of *Clostridium thermosulfurogens.* *FEMS Microbiol. Lett.* 17: 295-298.

Schmidt, F.X. (1989). T.E. Creighton Ed., Protein Structure: A Practical Approach, IRL. Press, Oxford, p. 261.

Schultz, J. and Floyd, T. (1979). Desert Survivor: North America's Creosote Bush. *Natural History*. 108: 24.

Schumann, J., Wrba, A., Jaenicke, R. and Stetter, K.O. (1991). Topographical and enzymatic characterization of amylases from the extremely thermophilic eubacterium *Thermotoga maritime*. *FEBS Lett.* 282: 122-126.

Schwantes, D.G. (1957). English Trans. V. Higgins. Flowering Stones and Mid-day Flowers. London: Ernest Benn Ltd.

Sidhu, G.S., Sharma, P., Chakrabarti, T. and Gupta, J.K. (1997). Strain improvement for the production of a thermostable alpha-amylase. *Enzyme Microb. Technol.* 21: 525-30.

Smith, T.A. and Smith, P.W. (1998). Respiratory Symptoms and sensitization in bread and cake bakers. *Occup. Med.* 48: 321-328.

Sonnleitner, B. and Fiechter, A. (1983). Advantages of using thermopiles in biotechnological process: expectations and reality. *Trends in Biotechnol.* 1: 74-80.

Southern, E. (1979). Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.* 98: 513-517.

Spano, L., Medeiros, J. and Mandels, M. (1975). *Enzymatic Hydrolysis of Cellulosic Waste to Glucose*, Pollution Abatement Div., Food Svcs. Labs, US Army Natick, MA, USA.

Specka, U., Mayer, F and Antranikian, G. (1991). Purification and properties of a thermoactive glucoamylase from *Clostridium thermosaccharolyticum*. *Appl. Environ. Microbiol.* 57: 2317-2323.

Srivastava, R.A.K. and Baruah, J.N. (1986). Culture conditions for production of thermostable amylase by *Bacillus stearthermophilus*. *Appl. Environ. Microbiol.* 52: 179-184.

Stetter, K.O. (1992). Life at the upper temperature border, in: J. Van, K.T.T. Van, J.C. Mounolou, J. Schneider, C. McKay Eds., Colloque Interdisciplinaire du Comite National de la Recherche Scientifique, Frontiers of Life, Editions Frontieres, Gif-sur-Yvette. pp. 195-219.

Stetter, K.O. (1996). Hyperthermophilic prokaryotes. *FEMS Microbiol Rev.* 18: 149-158.

Stetter, K.O. (1998). Hyperthermophilic: isolation, classification and properties. In *Extremophiles – Microbial Life in Extreme Environments*. Edited by Horikoshi, K. and Grant, W.D. New York: Wiley-Liss. 1-24.

Stetter, K.O. (1999). Extremophiles and their adaptation to hot environments. *FEBS Lett.* 452: 22-25.

Sunna, A., Moracci, M., Rossi, M. and Antranikian, G. (1996). Glycosyl hydrolases from hyperthermophiles. *Extremophiles.* 1: 2-13.

Suresh, C., Dubey, A.K., Srikanta, S., Kumar, S.U. and Karanth, N.G. (1999). Characterization of a starch-hydrolyzing enzyme of *Aspergillus niger*. *Appl. Microbiol. Biotechnol.* 51: 673-675.

Suzuki, Y., Ito, N., Yuuki, T., Yamagata, H. and Udaka, S. (1989). Amino acid residues stabilizing a *Bacillus* alpha-amylase against irreversible thermoinactivation. *J. Biol. Chem.* 264: 18933-18938.

Suzuki, Y., Oishi, K., Nakano, H. and Nagayama, T. (1987). A strong correlation between the increase in number of praline residues and the rise in thermostability of five *Bacillus* oligo-1,6-glucosidase. *Appl. Microbiol. Biotechnol.* 26: 546-551.

Swamy, M.V., Sai Ram, M. and Seenayya, G. (1994). Beta-amylase from *Clostridium thermocellum* SSA thermophilic anaerobic cellulolytic bacterium. *Letf. Appl. Microbiol.* 18: 301-304.

Tachibana, Y., Kuramura, A., Shirasaka, N., Suzuki, Y., Yamamoto, T., Fujiwara, S., Takagi, M. and Imanaka, T. (1999). Purification and characterization of an extremely thermostable cyclomaltodextrin glucanotransferase from a newly isolated hyperthermophilic archaeon, a *Thermococcus* sp. *Appl. Environ. Microbiol.* 65: 1991-1997.

Teles, F.F.F. (1978). Nutrient analysis of prickly pear (*Opuntia ficus indica*). Ph.D Thesis. Ann Arbor, Michigan.

Tel-Zur, N., Abbo, S., Myslanbodski, D. and Mizarhi, Y. (1999). Modified CTAB Procedure for DNA Isolation from Epiphytic Cacti of the Genera *Hylocereus* and *Selenicereus* (Cactaceae). *Plant Molecular Biology Reporter.* 17: 249-254.

Theriez, M. (1965). Recherches sur la digestibilite de *Opuntia ficus- indica* var. inermis en Tunisie. Inst. Nat. Rech. Agr.

Thompson, J.D., Higgins, D.G. and Gibson, T.J. (1994). CLUSTALW: improving the sensitivity of progressive multiple sequence alignment trough sequence weighting, position specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 22: 4673–4680.

Tolstoguzov, V (2004). Why are polysaccharides necessary? *Food Hydrocolloids.* 18: 873-877.

Tomazic, S.J. and Klibanov, A.M. (1988). Why is one *Bacillus* alpha-amylase more resistant against irreversible thermo inactivation than another? *J. Biol. Chem.* 263: 3092–3096.

Tsujibo, H., Minoura, K., Miyamoto, K., Endo, H., Moriwaki, M. and Inamori, Y. (1993). Purification and properties of a thermostable chitinase from *Streptomyces thermoviolaceus* OPC- 520. *Appl. Environ. Microbiol.* 59: 620-622.

Turner, S.L., Ford, G.C., Mountain, A. and Moir, A. (1992). Selection of a thermostable variant of chloramphenicol acetyltransferase (Cat-86). *Protein Eng.* 5: 535-54.

Vallee, B. L., Stein, E. A., Sumerwell, W. N. & Fischer, E. H. (1959). Metal content of α -amylase of various origins. *J. Biol. Chem.* 234: 2901-2905

Van Wormhoudt, A. and Sellos, D. (1996). *J. Mol. Evol.* 42: 543-551.

Vetriani, C., Maeder, D.L., Tolliday, N., Yip, K.S.P., Stillman, T.J., Britton, K.L., Rice, D.W., Klump, H.H. and Robb, T.T. (1998). Protein thermostability above 100°C: A key role for ionic interactions. *Proc. Natl Acad. Sci. USA.* 95: 12300-12305.

Vieille, C. and Zeikus, G.J. (2001). Hyperthermophilic enzymes: sources, uses and molecular mechanism for thermostability. *Microbiol Mol Biol.* 65: 1-43.

Vihinen, M. and Mantsala, P. (1989). *Critical. Rev. Biochem. Mol. Biol.* 24: 329-418.

Vogt, G. and Argos, P. (1997). Protein thermal stability: hydrogen bonds or internal packing? *Folding Des.* 2: S40-S60.

Watanabe, K and Suzuki, Y. (1998). Protein thermostabilization by proline substitutions *J. Mol. Catal. B.* 4: 167-180.

Watanabe, K., Chishiro, K., Kitamura, K. and Suzuki, Y. (1991). Proline residues responsible for thermostability occur with high frequency in the loop regions of an extremely thermostable oligo-1,6-glucosidase from *Bacillus thermoglucosidasius* KP 1006. *J. Biol. Chem.* 266: 24287-24294.

Watanabe, K., Kitamura, K. and Suzuki, Y. (1996). Analysis of the critical sites for protein thermostabilization by proline substitution in oligo-1,6-glucosidase from *Bacillus coagulans* ATCC 7050 and the evolutionary consideration of proline residues. *Appl. Environ. Microbiol.* 62: 2066-2073.

Watanabe, K., Masuda, T., Ohashi, H., Mihara, H and Suzuki, Y. (1994). Multiple proline substitutions cumulatively thermostabilize *Bacillus cereus* ATCC 7064 oligo-1,6-glucosidase. *Eru. J. Biochem.* 226: 277-283.

Watanabe, T., Yamamoto, A., Nagai, S. and Terabe, S. (1998). Simultaneous measurement of alpha-amylase and glucoamylase activity in sake rice koji by capillary electrophoresis of sodium-protein complex and activity measurement of glucoamylase by in-capillary enzyme reaction method. *Electrophoresis.* 19: 2331-2337.

Wilson, J.J. and Ingledew, W.M. (1982). Isolation and Characterization of *Schwanniomyces alluvius* amyolytic enzymes. *Appl. Environ. Microbiol.* 44: 301-307.

Wind, R.D., Buitelaar, R.M., Eggink, G., Huizing, H.J. and Dijkhuizen, L. (1994). Characterization of a new *Bacillus stearothermophilus* isolate: a highly thermostable alpha-amylase producing strain. *Appl. Microbiol. Biotechnol.* 41: 155-162.

Wanderley, K. J, Torres, F .A., Moraes, L. M. and Ulhoa, C. J (2004). Biochemical characterization of alpha amylase from yeast *Cryptococcus flavus*. *FEMS Microbiol Lett.* 16: 165-9

Woodward, T.E., Turner, W.F. and Griffith, D. (1915). Prickly pear feed for dairy cows. I. *Ag. Res., USDA, Washington, D.C.* 4: 405-449.