

LIST OF TABLES

Table No.	Particulars	Page No.
3.1	Systematic Position of the <i>Centella asiatica</i> (L) Urban	42
3.2	Nutritional components of MS medium	48
4.1	Effect of colchicine concentration on survival rate and response of induction of autotetraploidy in <i>Centella asiatica</i> .	60
4.2	Differences in Morphometric characters of diploid and colchicine treated (0.2% for 4 hours) plants of <i>C. asiatica</i> .	67
4.3	Comparison of stomatal characteristics in diploid and autotetraploid plants of <i>Centella asiatica</i> .	72
4.4	Triterpenoid saponin contents of diploid and autotetraploid plants of <i>Centella asiatica</i> .	74
4.5	HPLC analysis of peak area and retention time for madecassoside + asiaticoside B, asiaticoside, madecassic acid + terminolic acid and asiatic acid in diploid <i>C. asiatica</i> plant.	75
4.6	HPLC Analysis of peak area and retention time for madecassoside + asiaticoside B, asiaticoside, madecassic acid + terminolic acid and asiatic acid in colchicine induced autotetraploid to the <i>Centella asiatica</i>	76
4.7	HPLC Analysis of peak area and retention time for madecassoside + asiaticoside B, asiaticoside and asiatic acid in standard to the <i>Centella asiatica</i> .	77
4.8	Effect of combination of 2,4 D / IAA + BAP on proliferation rate and percent induction of callus in <i>Centella asiatica</i> leaf explants.	79
4.9	Effect of various concentrations and combinations of IBA and Kinetin on mean number and length of shoots produced <i>in vitro</i> from <i>Centella asiatica</i> leaf explants.	82
4.10	Effect of various concentrations and combinations of NAA and IBA on induction and mean length of roots from <i>C. asiatica</i> leaf explants.	83