

LIST OF FIGURES

Figure No.	Particulars	Page No.
1.1	<i>Malassezia furfur</i>	6
1.2	<i>M. furfur</i> exhibiting catalase activity, esculin hydrolysis and tween 80 hydrolysis activity	7
1.3	Pathologies associated with <i>M. furfur</i>	7
1.4	Mechanism of action of <i>Malassezia</i> species in the aetiology of dandruff/ Seborrheic dermatitis	11
3.1	Selected medicinal plants which possessed antimalassezial activity	62
3.2	Antifungal activity of plant extracts on <i>M. furfur</i> by disc diffusion method	63
3.3	Antifungal activity of essential oils on <i>M. furfur</i> by disc diffusion method	71
3.4	Determination of Minimum Inhibitory Concentration (MIC) of plant extracts on <i>M. furfur</i> by microdilution method	73
3.5	Determination of Minimum Inhibitory Concentration (MIC) and Minimum Yeast-cidal Concentration (MYC) of plant extracts on <i>M. furfur</i> by streak plate method	74
3.6	Determination of Minimum Inhibitory Concentration (MIC) of essential oils on <i>M. furfur</i> by microdilution method	75
3.7	Determination of Minimum Inhibitory Concentration (MIC) and Minimum Yeast-cidal Concentration (MYC) of essential oils on <i>M. furfur</i> by streak plate method	76
3.8	Antifungal activity of blended essential oil on <i>M. furfur</i>	77
3.9	HPTLC profile of <i>n</i> - hexane extract of <i>Juglans nigra</i> pericarp	81
3.10	HPTLC profile of aqueous extract of <i>Allium sativum</i> bulb	81
3.11	HPTLC profile of <i>n</i> - hexane extract of <i>Cinnamomum verum</i> bark	82
3.12	HPTLC profile of <i>n</i> - hexane extract of <i>Syzygium aromaticum</i> flower buds	82
3.13	HPTLC profile of <i>Lawsonia inermis</i> leaf extract	83
3.14	TLC of <i>Juglans nigra</i> (<i>n</i> - hexane extract) with bioautography showing inhibition zones on pink background	84
3.15	TLC of <i>Allium sativum</i> (aqueous extract) with bioautography showing inhibition zones on pink background	84
4.1	Flash chromatography, silica 40 g cartridge column and collected fractions	103
4.2	HPTLC fingerprint of <i>Juglans nigra</i> fractions under short wave UV (254 nm)	104

4.3	Antimalassezial activity of <i>Juglans nigra</i> pericarp fractions (1-10) by disc diffusion, microdilution method and streak plate method	105
4.4	LC-MS-MS spectrum of standard Juglone	107
4.5	LC-MS-MS spectrum of Fraction 5-Juglone isolated from <i>Juglans nigra</i> pericarp	108
4.6	Antimalassezial activity of <i>Allium sativum</i> fractions by disc diffusion, microdilution and streak plate method	110
4.7	HPTLC profile of hexane fraction of <i>Allium sativum</i> bulb	111
4.8	Gas chromatogram of <i>n</i> -hexane fraction of <i>A. sativum</i> bulb	111
4.9	Fluorescence imaging of <i>M. furfur</i> under 40 X magnification	114
4.10	Scanning electron micrographs of untreated and treated <i>M. furfur</i>	115
4.11	Cytotoxic effect of the test samples on 3T3 Cell line	117
4.12	Microscopic examination of untreated and treated 3T3 cells	117
5.1	Flow chart shows the synthesis of clove extract (CE) mediated ZnO nanoparticles (NPs)	126
5.2	FTIR spectra of CE mediated ZnO (30 mL) NPs	130
5.3	UV- visible absorption spectra of CE mediated ZnO (10-30 mL) NPs	131
5.4	PXRD patterns of as-formed CE mediated ZnO NPs	132
5.5	Williamson- Hall plots of ZnO NPs synthesised with different concentration of clove extracts (10, 15, 20, 25 and 30 mL)	133
5.6	Rietveld refined structural pattern and Packing diagram obtained from diamond software	134
5.7	SEM micrographs of CE mediated ZnO NPs	135
5.8	Formation process of CE mediated ZnO NPs	136
5.9	TEM, HRTEM micrographs and SAED pattern of CE mediated ZnO NPs	137
5.10	Antifungal activity of CE mediated ZnO NPs on <i>M. furfur</i> by disc diffusion method, microdilution method and streak plate method	139
5.11	Fluorescence imaging of yeast cells	140