

Bibliography

- [1] K. Afsarinejad and A.S. Hedayat, *Repeated measurements designs for a model with self and mixed carryover effects*, Journal of Statistical Planning and Inference 106 (2002), pp. 449–459.
- [2] R.A. Bailey, *Quasi-complete Latin squares: construction and randomization*, Journal of the Royal Statistical Society B 46 (1984), pp. 323-334.
- [3] M. Bose and B. Mukherjee, *Cross-over designs in the presence of higher order carry-overs*, Australian and New Zealand Journal of Statistics 42 (2000), pp. 235-244.
- [4] M. Bose and B. Mukherjee, *Optimal crossover designs under a general model*, Statistics and Probability Letters 62 (2003), pp. 413-418.
- [5] V. Brusasco, E. Crimi, G. Gherson, R. Nardelli, V. Oldani, B. Francucci, G. Cioppa, S. Senn and L. Fabbri, *Actions other than Smooth Muscle Relaxation may Play a Role in the Protective Effects of Formoterol on the Allergen-Induced Late Asthmatic Reaction*, Pulmonary Pharmacology and Therapeutics 15 (2002), pp. 399-406.
- [6] K.C. Carriere, *Optimal two-period repeated measurement designs with two or more treatments*, Biometrika 80 (1993), pp. 924–929.

- [7] C.S. Cheng and C.F. Wu, *Balanced repeated measurements designs*, *Annals of Statistics* 8 (1980), pp. 1272–1283.
- [8] W.G. Cochran and G.M. Cox, *Experimental Designs*, 2nd edition, Wiley, New York, 1957.
- [9] J. Divecha and J. Gondaliya, *Estimation of treatment and carryover effects in optimal cross-over designs for clinical trials*, *Statistics in Biopharmaceutical Research* 7 (2015), pp. 95–105.
- [10] A.F. Ebbutt, *Three-period crossover designs for two treatments*, *Biometrics* 40 (1984), pp. 219 – 224.
- [11] FDA, *Guidance for Industry: Statistical Approaches to Establishing Bioequivalence*, US Department of Health and Human Services, Food and Drug Administration, Center for Drug Evaluation and Research (CDER), 2001.
- [12] W.T. Federer, *Experimental Design: Theory and Application*, Macmillan, New York, 1955.
- [13] D.J. Finney, *Cross-over designs in bioassay*, *Journal of Royal Statistical Society B* 145 (1956), pp. 42–61.
- [14] P.S. Gill, *Balanced change-over designs for autocorrelated observations*, *Australian Journal of Statistics* 34 (1992), pp. 415–420.
- [15] P.S. Gill and G.K. Shukla, *Optimal change-over designs for correlated observations*, *Communication in Statistics-Theory and Methods* 16 (1987), pp. 2243–2261.
- [16] J.E. Grizzle, *The two-period change-over design and its use in clinical trials*, *Biometrics* 21 (1965), pp. 467–480.
- [17] K. Hanford, *Lecture Notes on Mixed Models*, Springer, United States, 2005.

- [18] A.S. Hedayat and K. Afsarinejad, *Repeated measurements designs, I. In a Survey of Statistical Design and Linear Models* (J.N. Srivastava, Ed.), 1975, pp. 229–242 (North-Holland, Amsterdam).
- [19] A.S. Hedayat and K. Afsarinejad, *Repeated measurements designs II*, *Annals of Statistics* 6 (1978), pp. 619–628.
- [20] A. Hedayat and J. Stufken, *Optimal and efficient crossover designs under different assumptions about the carryover effects*, *Journal of Biopharmaceutical Statistics* 13 (2003), pp. 519–528.
- [21] A.S. Hedayat and M. Yang, *Universal optimality of balanced uniform crossover designs*, *Annals of Statistics* 31 (2003), pp. 978–983.
- [22] A.S. Hedayat and M. Yang, *Universal optimality for selected crossover designs*, *Journal of American Statistical Association* 99 (2004), pp. 461–466.
- [23] ICH E14 Guidance, *The Clinical Evaluation of QT/QTc Interval Prolongation and Proarrhythmic Potential for Nonantiarrhythmic Drugs*, International Conference on Harmonization, in Geneva, Switzerland, May 2005.
- [24] R.A. Ipinyomi, *Equineighboured experimental designs*, *Australian Journal of Statistics* 28 (1986), pp. 79–88.
- [25] J. John, K. Russell and D. Whitaker, *CrossOver: an algorithm for the construction of efficient cross-over designs*, *Statistics in Medicine* 23 (2004), pp. 2645–2658.
- [26] B. Jones and M.G. Kenward, *Design and Analysis of Crossover Trials*, Chapman and Hall/CRC, Boca Raton, 2003.
- [27] J.A. Jones and M.H. Quenouille, *Experiments: Design and Analysis*, 2nd Edition, London:Charles Griffin, 1977.

- [28] B. Jones and A.N. Donev, *Modelling and design of cross-over trials*, *Statistics in Medicine* 15 (1996), pp. 1435–1446.
- [29] R. Kempton, S. Ferris and O. David, *Optimal change-over designs when carry-over effects are proportional to direct effects of treatments*, *Biometrika* 88 (2001), pp. 391–399.
- [30] J. Kiefer, *Construction and optimality of generalized Youden designs*, In *A Survey of Statistical Designs and Linear Models* (J. N. Srivastava, Ed.) pp. 333–353, Amsterdam:North-Holland, 1975.
- [31] J. Kiefer and H.P.Wynn, *Optimal balanced block and Latin squares designs for correlated observations*, *The Annals of Statistics* 9 (1981), pp. 737–757.
- [32] G. Koch, I. Amara, B. Brown, T. Colton and D. Gillings, *A two-period crossover design for the comparison of two active treatments and placebo*, *Statistics in Medicine* 8 (1989), pp. 487–504.
- [33] J. Kunert, *Optimality of balanced uniform repeated measurement designs*, *The Annals of Statistics* 12 (1984), pp. 1006–1017.
- [34] J. Kunert, *Crossover designs for two treatments and correlated errors*, *Biometrika* 78 (1991), pp. 315–324.
- [35] J. Kunert and J. Stufken, *Optimal crossover designs in a model with self and mixed carryover effects*, *Journal of American Statistical Association* 97 (2002), pp. 898–906.
- [36] H.B. Kushner, *Optimal repeated measurements designs: the linear optimality equations*, *The Annals of Statistics* 25 (1997), pp. 2328–2344.
- [37] H.B. Kushner, *Optimal and efficient repeated measurements designs for uncorrelated Observations*, *Journal of American Statistical Association* 93 (1998), pp. 1176–1187.

- [38] M.E. Laska and M. Meisner, *A variational approach to optimal two-treatment cross-over designs: application to carry-over effect models*, Journal of American Statistical Association 80 (1985), pp. 704–710.
- [39] N. Laird, J. Skinner and M. Kenward, *An analysis of two-period crossover designs with carry-over effects*, Statistics in Medicine 11 (1992), pp. 1967–1979.
- [40] W. Lehmacher, *Analysis of cross-over trials in the presence of residual effects*, Statistics in Medicine 10 (1991), pp. 891–899.
- [41] H.L. Lucas, *Extra-period Latin square changeover designs*, Journal of Dairy Science 4 (1957), pp. 225 – 239.
- [42] R.J. Martin and J.A. Eccleston, *Variance-balanced change-over designs for dependent observations*, Biometrika 85 (1998), pp. 883–892.
- [43] J.N.S. Matthews, *Modeling and optimality in the design of crossover studies for medical applications*, Journal of Statistical Planning and Inference 42 (1994), pp. 89–108.
- [44] J.P. Morgan, *Balanced polycross designs*, Journal of the Royal Statistical Society B 50 (1988), pp. 93–104.
- [45] M. Ozan and J. Stufken, *Assessing the impact of carryover effects on the variances of estimators of treatment differences in crossover designs*, Statistics in Medicine 29 (2010), pp. 2480–2485.
- [46] A. Ring, B. Walter, A. Larbalestier and D. Chanter, *An efficient crossover design for thorough QT studies*, Biometrie und Epidemiologie 6 (2010), pp. 1–12.
- [47] S. Satpati, V. Gupta, R. Parsad and M. Aggarwal, *Computer-generated change-over designs for correlated observations*, Communications in Statistics-Theory and Methods 41 (2012), pp. 3786–3798.

- [48] S. Senn, *Is the simple carryover model useful?*, *Statistics in Medicine* 11 (1992), pp. 715–726.
- [49] S. Senn, *Cross-over Trials in Clinical Research*, John Wiley and Sons, New York, 1993.
- [50] S. Senn, *Cross-over Trials in Clinical Research*, Second Edition, John Wiley and Sons, New York, 2003.
- [51] S. Senn and D. Lambrou, *Robust and realistic approach to carry-over*, *Statistics in Medicine* 17 (1998), pp. 2849–2864.
- [52] M.T. Taka and P. Armitage, *Autoregressive models in clinical trials*, *Communications in Statistics: Theory and Methods* 12 (1983), pp. 865–867.
- [53] A.N. Tsoy, O.V. Cheltzov, V. Zaseyeva, L.A. Shilinish and L.A. Yashina, *European Respiratory Journal* 3 (1990), pp. 235s.
- [54] E.J. Williams, *Experimental designs balanced for the estimation of residual effects of treatments*, *Australian Journal of Scientific Research* 2 (1949), pp. 149–168.
- [55] M. Yang and J. Stufken, *Optimal and efficient crossover designs for comparing test treatments to a control treatment under various models*, *Journal of Statistical Planning and Inference* 138 (2008), pp. 278–285.