

List of Tables

2.1	The γ -ray energy and absolute intensities in the standard γ - source ^{152}Eu	33
2.2	List of identified ERs produced in $^{20}\text{Ne} + ^{55}\text{Mn}$ system via CF and/or ICF and their spectroscopic data	37
2.3	List of identified ERs produced in $^{20}\text{Ne} + ^{159}\text{Tb}$ system via CF and/or ICF and their spectroscopic data	39
2.4	List of identified ERs produced in $^{16}\text{O} + ^{156}\text{Gd}$ system via CF and/or ICF and their spectroscopic data	42
2.5	Deduced expressions for the extraction of independent cross-sections from the measured cumulative cross-sections for $^{20}\text{Ne} + ^{55}\text{Mn}$ system	50
2.6	The measured cross-section for the production of ERs ^{70}Se , ^{72}As , ^{71}As and ^{70}As in $^{20}\text{Ne} + ^{55}\text{Mn}$ system	51
2.7	The measured cross-section for the production of ERs ^{69}Ge , ^{67}Ge , ^{66}Ge and ^{67}Ga in $^{20}\text{Ne} + ^{55}\text{Mn}$ system	54
2.8	The measured cross-section for the production of ERs ^{66}Ga , ^{65}Ga , ^{63}Zn , ^{61}Cu and ^{60}Cu in $^{20}\text{Ne} + ^{55}\text{Mn}$ system	54
2.9	Deduced expressions for the extraction of independent cross-sections from the measured cumulative cross-sections for $^{20}\text{Ne} + ^{159}\text{Tb}$ system	59
2.10	The measured cross-section for the production of ERs ^{174}W , ^{175}Ta and ^{173}Ta in $^{20}\text{Ne} + ^{159}\text{Tb}$ system	60
2.11	The measured cross-section for the production of ERs ^{172}Ta , ^{173}Hf and ^{164}Yb in $^{20}\text{Ne} + ^{159}\text{Tb}$ system	61
2.12	The measured cross-section for the production of ERs ^{166}Tm , ^{165}Tm , ^{163}Tm and ^{161}Er in $^{20}\text{Ne} + ^{159}\text{Tb}$ system	63
2.13	Deduced expressions for the extraction of independent cross-sections from the measured cumulative cross-sections for $^{16}\text{O} + ^{156}\text{Gd}$ system	68

2.14	The measured cross-section for the production of ERs ^{168}Hf , ^{166}Hf , ^{167}Lu and ^{167}Yb in $^{16}\text{O} + ^{156}\text{Gd}$ system	69
2.15	The measured cross-section for the production of ERs ^{164}Yb , ^{162}Yb , ^{166}Tm and ^{165}Tm in $^{16}\text{O} + ^{156}\text{Gd}$ system	69
2.16	The measured cross-section for the production of ERs ^{163}Tm , $^{162}\text{Tm}^g$, ^{161}Er in $^{16}\text{O} + ^{156}\text{Gd}$ system	70
2.17	The measured cross-section for the production of ERs ^{159}Er , $^{162}\text{Ho}^m$, $^{159}\text{Ho}^g$ and ^{157}Dy in $^{16}\text{O} + ^{156}\text{Gd}$ system	70
3.1	List of Al-catcher foil thicknesses used in FRRD measurements, for $^{20}\text{Ne} + ^{159}\text{Tb}$ and $^{16}\text{O} + ^{156}\text{Gd}$ systems at different projectile energies	92
3.2	Experimentally measured forward recoil range distributions (FRRDs) of the residues ^{174}W , ^{175}Ta , ^{173}Ta , ^{172}Ta and ^{173}Hf produced in $^{20}\text{Ne} + ^{159}\text{Tb}$ reactions at ~ 164 MeV energy	97
3.3	Experimentally measured forward recoil range distributions (FRRDs) of the residues ^{164}Yb , ^{166}Tm , ^{165}Tm and ^{163}Tm produced in $^{20}\text{Ne} + ^{159}\text{Tb}$ reactions at ~ 164 MeV energy	98
3.4	The experimentally measured forward recoil ranges $R_p(\text{exp})$ deduced from RRD curves, and theoretically calculated most probable ranges $R_p(\text{theo})$ for CF and /or ICF component using range energy relation for the reaction products produced in the interaction of ^{20}Ne with ^{159}Tb at energy, $E \sim 164$ MeV	104
3.5	The measured relative contribution of CF and/or ICF of the reaction residues populated in $^{20}\text{Ne} + ^{159}\text{Tb}$ system at energy, $E \sim 164$ MeV	106
3.6	Experimentally measured forward recoil range distributions (FRRDs) of the residues ^{168}Hf , ^{167}Lu , ^{167}Yb , ^{162}Yb , ^{165}Tm and ^{163}Tm produced in $^{16}\text{O} + ^{156}\text{Gd}$ reactions at ~ 72 MeV energy	107
3.7	Experimentally measured forward recoil range distributions (FRRDs) of the residues ^{168}Hf , ^{167}Lu , ^{167}Yb and ^{162}Yb produced in $^{16}\text{O} + ^{156}\text{Gd}$ reactions at ~ 82 MeV energy	108
3.8	Experimentally measured forward recoil range distributions (FRRDs) of the residues ^{165}Tm , ^{163}Tm , $^{159}\text{Ho}^g$ and ^{157}Dy produced in $^{16}\text{O} + ^{156}\text{Gd}$ reactions at ~ 82 MeV energy	111

3.9	Experimentally measured forward recoil range distributions (FRRDs) of the residues ^{168}Hf , ^{167}Lu , ^{167}Yb and ^{162}Yb produced in $^{16}\text{O} + ^{156}\text{Gd}$ reactions at ~ 93 MeV energy	112
3.10	Experimentally measured forward recoil range distributions (FRRDs) of the residues ^{165}Tm , ^{163}Tm , $^{159}\text{Ho}^g$ and ^{157}Dy produced in $^{16}\text{O} + ^{156}\text{Gd}$ reactions at ~ 93 MeV energy	113
3.11	The experimentally measured forward recoil ranges $R_p(\text{exp})$ deduced from RRD curves, and theoretically calculated most probable ranges $R_p(\text{theo})$ for CF and /or ICF component using range energy relation for the reaction products produced in the interaction of ^{16}O with ^{156}Gd at energy, $E \sim 72$ and 82 MeV	123
3.12	The experimentally measured forward recoil ranges $R_p(\text{exp})$ deduced from RRD curves, and theoretically calculated most probable ranges $R_p(\text{theo})$ for CF and/or ICF component using range energy relation for the reaction products produced in the interaction of ^{16}O with ^{156}Gd at energy, $E \sim 93$ MeV	124
3.13	The measured relative contribution of CF and / or ICF of the reaction products populated in $^{16}\text{O} + ^{156}\text{Gd}$ system at three different projectile energies i.e., $E \sim 72$, 82 and 93 MeV	126
4.1	Salient features of Charged Particle Detector Array (CPDA)	134