

LIST OF TABLES

CHAPTERS	TABLE CAPTIONS	PAGE NO.
CHAPTER II	TableII.a Properties of Cyclodextrins	29
CHAPTER IV	Table1 Density (ρ), viscosity (η) and relative permittivity (ϵ) of solvent mixtures at 298.15 K ^a	99
	Table2 The concentration (c) and molar conductance (Λ) of LiAsF ₆ in different mass fraction of EG (w_1) in MeOH at 298.15 K ^a	100
	Table3 Limiting molar conductance (Λ_0), association constant (K_a), co-sphere diameter (R) and standard deviations of experimental Λ (δ) of LiAsF ₆ in different mass fraction of EG (w_1) in MeOH at 298.15 K ^a	102
	Table4 Walden product ($\Lambda_0 \cdot \eta$) and Gibb's energy change (ΔG^0) of LiAsF ₆ in different mass fraction of ethylene glycol (w_1) in methanol at 298.15 K ^a	103
	Table5 Ionic limiting molar conductance (λ_0^\pm), ionic Walden product ($\lambda_0^\pm \eta$), crystallographic radii (r_c) and Stoke's radii (r_s) of Li ⁺ and AsF ₆ ⁻ ion in different mass fraction of EG (w_1) in MeOH at 298.15 K ^a	104
	Table6 Experimental values of densities (ρ), apparent molar volume (ϕ_v), viscosities (η), of LiAsF ₆ in different mass fraction of EG (w_1) in MeOH at 298.15 K ^a	107
	Table7 Limiting apparent molar volume (ϕ_v^0), experimental slopes (S_v^*), A -, B - coefficients of LiAsF ₆ in different mass fraction of EG (w_1) in MeOH at 298.15 K	108
CHAPTER V	Table1 Change in chemical shifts (ppm) of the H3 and H5 protons of cyclodextrin molecules in four different host-guest complexes in D ₂ O at 298.15 K ^a .	118
	Table2 Values of surface tension (γ) and at the break point with corresponding concentration of aqueous α and β -cyclodextrins at 298.15 K ^a	121

List of Tables

CHAPTERS	TABLE CAPTIONS	PAGE NO.
	Table3 Association constants (K_a), Gibb's free energy, enthalpy and entropy of various ionic liquid-cyclodextrin systems	123
	Table4 The observed peaks at different m/z with corresponding ions for the solid inclusion complexes	126
	Table5 Limiting apparent molar volume (φv^0), Experimental slope (Sv^*), Viscosity B and A coefficient of [BMIm]Cl and [HMIm]Cl in different concentration of aqueous α -CD and β -CD solvent mixtures at 298.15K ^a	129
	TableS1 ¹ H-NMR spectra of [BMIm]Cl, [HMIm]Cl, α -CD, β -CD, and [BMIm]Cl+ α -CD, [BMIm]Cl+ β -CD, [HMIm]Cl+ α -CD, [HMIm]Cl+ β -CD complexes	132
	TableS2 Surface Tension (γ) values of aqueous [BMIm]Cl and [HMIm]Cl with α -CD and β -CD at 298.15 K ^a	133
	TableS3 Data for the conductivity study of aqueous [BMIm]Cl+ β -CD and [BMIm]Cl+ β -CD system (concentration of stock solution of [BMIm]Cl and [HMIm]Cl = 10mM, concentration of stock solution of β -CD = 10mM) at 293.15K ^a , 303.15K ^a , 313.15K ^a	134
	TableS4 Data for the conductivity study of aqueous [BMIm]Cl+ α -CD and [HMIm]Cl+ α -CD system (concentration of stock solution of [BMIm]Cl and [HMIm]Cl = 10mM, concentration of stock solution of α -CD = 10mM) at 293.15K ^a , 303.15K ^a , 313.15K ^a	135
	TableS5 Values of conductivity (k) at the break point with corresponding concentration of aqueous α and β -cyclodextrins at 298.15 K ^a , 298.15 K ^a , 298.15 K ^a	135
	TableS6 Data obtained from FT-IR spectroscopic study of α -CD, β -CD, [BMIm]Cl, [HMIm]Cl, α -CD+[BMIm]Cl, β -CD+[BMIm]Cl, α -CD+[HMIm]Cl, β -CD+[HMIm]Cl	136
	TableS7 Experimental values of density (ρ) and viscosity (η) in deferent mass fractions of aq. α -CD and β -CD	137

List of Tables

CHAPTERS	TABLE CAPTIONS	PAGE NO.
	TableS8 Experimental values of densities (ρ) and viscosities (η) with varying concentration in different mass fractions of aq. α -CD and β -CD [BMIm]Cl at 298.15K ^a	137
	TableS9 Experimental values of densities (ρ) and viscosities (η) with varying concentration in different mass fractions of aq. α -CD and β -CD for [HMIm]Cl at 298.15 K ^a	138
CHAPTER VI	Table1 Values of surface tension (γ) and at the break point with corresponding concentration of α -CD , β -CD for [BMPy]PF ₆ at 298.15 K ^a	154
	Table2 Association constants obtained by the Benesi–Hildebrand method (K_a)from UV-Vis study and corresponding thermodynamic parameters and stoichiometry of [BMPy]PF ₆ & α -CD and [BMPy]PF ₆ & β -CD inclusion complexes at 293.15K ^a , 303.15K ^a at 313.15K ^a .	156
	Table3 Association constants obtained by the nonlinear program (K_a^c) from conductance study and corresponding thermodynamic parameters and stoichiometry of [BMPy]PF ₆ & α -CD and [BMPy]PF ₆ & β -CD inclusion complexes at 293.15K ^a , 303.15K ^a and 313.15K ^a .	157
	TableS1 Data for Job's Plot performed by UV-Vis spectroscopy for [BMPy]PF ₆ - β -CD system at 298.15 K ^a	163
	TableS2 Data for Job's Plot performed by UV-Vis spectroscopy for [BMPy]PF ₆ - α -CD system at 298.15 K ^a	163
	TableS3 ¹ H-NMR spectra of [BMPy]PF ₆ , α -CD, β -CD, and [BMPy]PF ₆ + α -CD, [BMPy]PF ₆ + β -CD complexes	164
	TableS4 Change in chemical shifts (ppm) of the H3 and H5 protons of cyclodextrin molecule in two different host-guest complexes in D ₂ O at 298.15 K ^a	164
	TableS5 Surface Tension (γ) values of α -CD& [BMPy]PF ₆ , β -CD & [BMPy]PF ₆ Complex at 298.15 K ^a	165

List of Tables

CHAPTERS	TABLE CAPTIONS	PAGE NO.
	TableS6 Data for the conductivity study of aqueous [BMPy]PF ₆ + α -CD and [BMPy]PF ₆ + β -CD system (concentration of stock solution of IL = 10mM, concentration of stock solution of CD = 10mM at 293.15K ^a , 303.15K ^a , 313.15K ^a)	166
	TableS7 Values of conductivity (κ) at the break point with corresponding concentrations of [BMPy]PF ₆ and CDs at 298.15 K ^a	167
	TableS8 Data for the Benesi-Hildebrand double reciprocal plot performed by UV-VIS spectroscopic study for [BMPy]PF ₆ - α -CD and [BMPy]PF ₆ - β -CD systems at 293.15 K ^a	167
	TableS9 Data for the Benesi-Hildebrand double reciprocal plot performed by UV-VIS spectroscopic study for [BMPy]PF ₆ - α -CD and [BMPy]PF ₆ - β -CD systems at 303.15 K ^a	168
	TableS10 Data for the Benesi-Hildebrand double reciprocal plot performed by UV-VIS spectroscopic study for [BMPy]PF ₆ - α -CD and [BMPy]PF ₆ - β -CD systems at 303.15K ^a	169
	TableS11 Data obtained from FT-IR spectroscopic study of α -CD, β -CD, [BMPy]PF ₆ , α -CD +[BMPy]PF ₆ , β -CD+[BMPy]PF ₆ .	170
	TableS12 The observed peaks at different m/z with corresponding ions for the solid inclusion complexes.	171
CHAPTER VII	Table1 Change in chemical shifts (ppm) of the H3 and H5 protons of cyclodextrin molecules in two different host-guest complexes in D ₂ O at 298.15 K ^a	180
	Table2 Values of surface tension (γ) at the break point with corresponding concentrations of TBPMS and CD at 298.15 K ^a	184
	Table3 Values of conductivity (κ) at the break point with corresponding concentrations of TBPMS and CD at 298.15 K ^a	185
	Table4 The observed peaks at different m/z with corresponding ions for the solid inclusion complexes	188

List of Tables

CHAPTERS	TABLE CAPTIONS	PAGE NO.
	Table5 Association constants (K_a), Gibb's free energy, enthalpy and entropy of various ionic liquid-cyclodextrin systems	189
	TableS1 $^1\text{H-NMR}$ spectra of TBPMS, α -CD, β -CD and TBPMS+ α -CD, TBPMS+ β -CD complexes	192
	TableS2 Surface Tension (γ) values of aqueous TBPMS with α -CD and β -CD at 298.15 K ^a	192
	TableS3 Data for the conductivity study of aqueous TBPMS+ α -CD and TBPMS+ β -CD system (concentration of stock solution of TBPMS = 10mM, concentration of stock solution of CD = 10mM at 293.15K ^a , 303.15K ^a , 313.15K ^a)	193
	TableS4 Data obtained from FT-IR spectroscopic study of α -CD, β -CD, TBPMS and the complexes of α -CD & TBPMS, β -CD & TBPMS	194
CHAPTER VIII	Table1 Values of surface tension (γ) and at the break point with corresponding concentration of aqueous α and β -cyclodextrins at 298.15 K ^a	207
	Table2 Limiting apparent molar volume (ϕ_v^0), Experimental slope (S_v^*), Viscosity B and A coefficient of PTX and 2-PAM in different concentration of aqueous α -CD and β -CD solvent mixtures.	215
	TableS1 Surface Tension (γ) values of aqueous PTX and 2-PAM with α -CD and β -CD at 298.15 K ^a	219
	TableS2 Data for Job's Plot performed by UV-Vis spectroscopy for aqueous Pentoxifylline- α -CD system at 298.15 K ^a	219
	TableS3 Data for Job's Plot performed by UV-Vis spectroscopy for aqueous Pentoxifylline- β -CD system at 298.15 K ^a	220
	TableS4 Data for Job's Plot performed by UV-Vis spectroscopy for aqueous Pentoxifylline- β -CD system at 298.15 K ^a	220
	TableS5 Data for Job's Plot performed by UV-Vis spectroscopy for aqueous Pralidoxim- β -CD system at 298.15 K ^a	221
	TableS6 Data for the Benesi-Hildebrand double reciprocal plot performed by UV-VIS spectroscopic study for aqueous Drug-CD systems at 298.15 K ^a	221

List of Tables

CHAPTERS	TABLE CAPTIONS	PAGE NO.
	TableS7 1H-NMR spectra of PTX, 2-PAM, α -CD, β -CD, and PTX+ α -CD, PTX+ β -CD, 2-PAM+ α -CD, 2-PAM+ β -CD complexes	222
	TableS8 Change in chemical shifts (ppm) of the H3 and H5 protons of cyclodextrin molecules in four different host-guest complexes in D ₂ O at 298.15 K ^a	223
	TableS9 Data obtained from FT-IR spectroscopic study of α -CD, β -CD, PTX, 2-PAM, α -CD+PTX, β -CD+PTX, α -CD+2-PAM, β -CD+2-PAM	223
	TableS10 Experimental values of density (ρ) and viscosity (η) in deferent mass fractions of aq. α -CD and β -CD	224
	TableS11 Experimental values of densities (ρ) and viscosities (η) with varying concentration in different mass fractions of aq. α -CD with drug molecules.	224
CHAPTER IX	Table1 The values of density (ρ_0), viscosity (η_0) and refractive index (n_D) of Li-Ox, Na-Ox, K-Ox at different mass fractions of aqueous UA solution at 293.15K ^a , 303.15 K ^a , 313.15 K ^a	233
	Table2 The values of density (ρ), viscosity (η) of Li-Ox, Na-Ox, K-Ox at different mass fractions of aqueous UA solution at 293.15K ^a , 303.15 K ^a , 313.15 K ^a	234
	Table3 The values of apparent molar volume (φ_V), limiting apparent molar volume (φ_V^0), S_V^* of Li-Ox, Na-Ox, K-Ox at different mass fractions of aqueous UA solution at 293.15K ^a , 303.15 K ^a , 313.15 K ^a	236
	Table4 The values of a_0 , a_1 , a_2 , φ_E^0 , $(\delta\varphi_E^0/\delta T)_P$ Li-Ox, Na-Ox, K-Ox at different mass fractions of aqueous UA solution at 293.15K to 313.15K	237
	Table5 Values of viscosity A and B-coefficient for Li-Oxalate, Na-Oxalate, K-Oxalate against different mass fraction of aqueous UA (w_1) solution	240
	Table6 The values of refractive index (n_D) and molar refractive index (R_M) of Li-Ox, Na-Ox, K-Ox at different mass fractions of aqueous UA solution at 298.15K ^a	242

LIST OF FIGURES

CHAPTERS	FIGURES	PAGE NO.
CHAPTER-I	Fig.I.a. Solvation of a salt.	20
	Fig.I.b. Inclusion complex.	21
CHAPTER-II	Fig.II.a. Pictorial representation of inclusion complex formation.	25
	Fig.II.b. Pictorial diagram of (i) Inclusion complex (ii) Cucurbituril[6], (iii) 18-Crown-6 and (iv) P-tert-butylcalix-4-arene.	26
	Fig.II.c. (a) Torus shape of cyclodextrin with dimensions and molecular structure of (b) α -cyclodextrin(c) β -cyclodextrin (d) γ -cyclodextrin.	28
	Fig.II.d. Different stoichiometric ratios in host-guest inclusion complexes.	30
CHAPTER-IV	Fig.1. Plot of molar conductance (Λ) and the square root of concentration (\sqrt{c}) of LiAsF ₆ in $w_1=0.00$ (-♦-), $w_1=0.25$ (-■-), $w_1=0.50$ (-▲-), $w_1=0.75$ (-●-), and $w_1=1.00$ (-*-) mass fraction of EG in MeOH at T=298.15K.	103
	Fig.2. Plot of limiting molar conductance (Λ_0) (-♦-) and Walden Product ($\Lambda_0\eta$) (-▲-) of LiAsF ₆ in different mass fraction (w_1) of EG in MeOH at T=298.15K.	103
	Fig.3. Plot of limiting ionic conductance (λ_o^\pm) of Li ⁺ (-♦-), AsF ₆ ⁻ (-■-) and ionic Walden Product ($\lambda_o^\pm\eta$) of Li ⁺ (-▲-), AsF ₆ ⁻ (-●-) in different mass fraction (w_1) of EG in MeOH at T=298.15K.	105
	Fig. 4. Plot of association constant (K_a) (-♦-), and Gibb's energy change (ΔG^0) (-▲-) of LiAsF ₆ in different mass fraction (w_1) of EG in MeOH at 298.15K.	105
	Fig.5. Plot of limiting apparent molar volume (ϕ_v^0) (-♦-) and viscosity B-coefficient (-▲-) vs. mass fraction (w_1) of EG in MeOH for LiAsF ₆ at T=298.15K.	109
	Fig.6. The plausible mechanism of interaction between LiAsF ₆ and Solvents (ethylene glycol and methanol).	110
	Fig.7. The plausible diagrams of molecular interaction and solvation of the ions in solvent mixtures.	112

CHAPTERS	FIGURES	PAGE NO.
CHAPTER-V	Fig.1 Representation of Molecular structure of of [BMIm]Cl and [HMIm]Cl and an overall view of cyclodextrin.	115
	Fig.2. ¹ H NMR spectra of (a) pure β -CD(top), [BMIm]Cl (middle) and β-CD&[BMIm]Cl complex (bottom) (b) pure β -CD(top), [HMIm]Cl (middle) and β -CD&[HMIm]Cl complex (bottom).	119
	Fig.3. Plausible stoichiometric inclusion ratio of the host and guest molecule. (Host:Guest).	120
	Fig.4. Linear relationships of logKa vs. 1/T for the interaction of [HMIm]Cl and α-CD (◆), [HMIm]Cl and β-CD (■), [BMIm]Cl and α-CD (▲), [BMIm]Cl and β-CD(●).	123
	Fig.5. FT-IR Spectra of (a) pure β -CD, [BMIm]Cl, and [BMIm]Cl + β -CD inclusion complex (b) pure β -CD, [HMIm]Cl, [HMIm]Cl + β -CD inclusion complex.	125
	Fig.6 (a) Limiting apparent molar volume (φv_0) and Sv^* (b) viscosity A and B-coefficient of [BMIm]Cl and [HMIm]Cl in different mass fractions (w1) of aqueous (■) α-CD and (■) β-CD solution at 298.15K.	128
	Fig.7. Schematic representation of inclusion of (a) [BMIm]Cl and [HMIm]Cl with α-CD and (b) [BMIm]Cl and [HMIm]Cl β-CD	131
	FigS1. ¹ H NMR spectra of (a) pure α-CD(top), [BMIm]Cl (middle) and α-CD&[BMIm]Cl complex (bottom) (b) pure α -CD(top), [HMIm]Cl (middle) and α-CD&[HMIm]Cl complex (bottom).	140
	Fig.S2. Surface tension of (a) [BMIm]Cl with α -CD (b) [BMIm]Cl with β-CD (c) [HMIm]Cl with α -CD (D) [HMIm]Cl with β - CD at 298.15K.	141
	Fig.S3. Plot of conductivity against square root of molar concentration (c) for (a) [BMIm]Cl + α-CD, (b) [BMIm]Cl+β-CD (c)[HMIm]Cl+α-CD and (d) [HMIm]Cl+β-CD complexes at 293.15 K (◆), 303.15K (■) and 313.15K (▲).	141
	FigS4. Schematic representation of fate of the ionic liquid in aqueous medium.	142
	FigS5. FT-IR Spectra of (a) pure β -CD, [BMIm]Cl, and [BMIm]Cl + β -CD inclusion complex (b) pure β -CD, [HMIm]Cl, [HMIm]Cl + β -CD inclusion complex.	143

CHAPTERS	FIGURES	PAGE NO.
	Fig.S6. ESI Mass spectra of (a) [BMIm]Cl- α -CD (b) [BMIm]Cl- β -CD (c) [HMIm]Cl- α -CD (d) [HMIm]Cl- β -CD inclusion complexes.	145
CHAPTER-VI	Fig.1. Job's Plot for (a) [BMPy]PF6 with α -CD and (b) [BMPy]PF6 with β -CD.	149
	Fig.2. ^1H NMR spectra of pure α -CD, β -CD, [BMPy]PF6 and [BMPy]PF6+ α -CD and [BMPy]PF6+ β -CD. (In D2O, 400 MHz)	151
	Fig.3. 2D ROESY spectra of the solid ICs of [BMPy]PF6- α -CD AND [BMPy]PF6- β -CD in D2O. (Cross correlations are indicated by red circles)	153
	Fig.4. Surface tension of [BMPy]PF6 with α -CD, [BMPy]PF6 with β -CD at 298.15 K.	154
	Fig.5. Plot of $\log K_a^c$ vs $1/T$ for the interaction of α -CD with [BMPy]PF6 (\blacklozenge) and β -CD with [BMPy]PF6(\blacksquare).	157
	Fig.6. Plot of $\log K_a$ vs $1/T$ for the interaction of α -CD with [BMPy]PF6 (\blacklozenge) and β -CD with [BMPy]PF6(\blacksquare).	158
	Fig.7: FT-IR spectra of free α -CD, β -CD, [BMPy]PF6 and their 1:1inclusion complexes ([BMPy]PF6+ α -CD) and ([BMPy]PF6+ β -CD) at 298.15K.	161
	Fig.8. Schematic representation of the inclusion of [BMPy]PF6 with Cyclodextrin in mixed solvent media.	162
	Fig.S1. 2D ROESY spectra of the solid ICs of [BMPy]PF6- α -CD in D2O. (Cross correlations are indicated by red circles)	172
	Fig.S2. Plot of conductivity with increasing concentration of α -CD and β -CD with [BMPy]PF6+ α -CD and [BMPy]PF6+ β -CD at (a) 293.15 K, (b) 303.15K and (c) 313.15K.	173
	Fig.S3. Benesi-Hildebrand double reciprocal plots for the effect of β -CD on the absorbance of [BMPy]PF6 + β -CD and [BMPy]PF6+ α -CD (252 nm) at different temperatures(a) and (b) shows at 293015K, (c) and (d) at 303.15K and (e) and (f) at 313.15K.	174
	Fig.S4. ESI mass spectra of [BMPy]PF6- β -CD and [BMPy]PF6- α -CD inclusion complex.	175
CHAPTER-VII	Fig.1. Molecular structures of (a) tetrabutylphosphonium methanesulfonate and (b) cyclodextrin molecule with interior and exterior protons (n = 6, 7 for α -CD and β -CD respectively).	178

CHAPTERS	FIGURES	PAGE NO.
	Fig.2. ¹ H-NMR spectra of (a) pure TBPMS (b) pure α -CD and (c) α -CD & TBPMS complex.	181
	Fig.3. 2D ROESY spectra of solid inclusion complex of 1:1 molar ratio of TBPMS+ α -CD complex and TBPMS + β -CD complex in D ₂ O (correlation signals are marked by green borders).	183
	Fig.4. Variation of surface tension of aqueous TBPMS with increasing concentration of aqueous α -CD and β -CD solution respectively at 298.15 K.	184
	Fig.5. FTIR spectra of free α -CD, TBPMS and their 1:1 inclusion complex (TBPMS: α -CD).	187
	Fig.6. Linear relationships of logK _f vs. 1/T for the interaction of TBPMS+ α -CD (\blacktriangle), TBPMS+ β -CD (\blacklozenge).	189
	Fig.7. Schematic representation of plausible interactions taking place in the inclusion complex.	191
	Fig.S1. ¹ H NMR spectra of (a) pure TBPMS (b) pure β -CD and (c) β -CD & TBPMS complex.	197
	Fig.S2. Plot of conductivity against square root of molar concentration (\sqrt{C}) for TBPMS + α -CD(\blacktriangle), TBPMS+ β -CD(\blacklozenge) at 293.15 K, 303.15K and 313.15K.	198
	Fig.S3. FT-IR spectra of free β -CD, TBPMS and their 1:1 inclusion complex (TBPMS: β -CD).	199
	Fig.S4. ESI mass spectra of TBPMS- β -CD inclusion complex and TBPMS- β -CD inclusion complex.	200
CHAPTER-VIII	Fig.1. Molecular structure of (a) one unit of CD, (b) PTX and (c) 2-PAM.	203
	Fig.2. Surface tension of PTX with α -CD (top left), PTX with β -CD (top right) 2-PAM with α -CD (bottom left) 2-PAM with β - CD (bottom right) at 298.15 K.	206
	Fig.3. Job plot of (a) PTX- α -CD at λ_{\max} =272 nm, (b) PTX- β -CD at λ_{\max} =272 nm, (c) 2-PAM- α -CD at λ_{\max} =292 nm, (d) 2-PAM- β -CD at λ_{\max} =292 nm. $\Delta A = (AIC-ACD)$, $R = [Drug]/([Drug]+[CD])$ (Total concentration of the Soln. =100 μ M).	208
	Fig.4. Schematic diagrams of inclusion (a) 2-PAM with α -CD and β -CD, (b) PTX with 2-PAM with α -CD and β -CD.	210

CHAPTERS	FIGURES	PAGE NO.
	Fig.5 Schematic diagrams of inclusion through the narrower and wider rim of the cyclodextrin molecule.	211
	Fig.6 ¹ H NMR spectra of (a) pure β -CD, PTX and β-CD + PTX (b) 2-PAM, pure β -CD and β-CD + PTX. (Green and blue colored protons are of PTX and 2-PAM respectively while red colored protons are of CD).	212
	Fig.7 FT-IR Spectra of (a) pure β -CD, PTX, and β-CD + PTX and (b) 2-PAM, pure β -CD, β-CD + PTX.	214
	Fig.8 (a) Limiting apparent molar volume (ϕ_{v0}) and (b) viscosity B-coefficient of PTX and 2-PAM in different mole fractions (w_1) of aqueous (■) α-CD and (■)β-CD solution at 298.15K.	217
	Fig.S1. Absorbance vs. wave length curve of α-CD and β-CD with PTX and 2-PAM.	226
	Fig.S2. ¹ H NMR spectra of (a) pureα-CD, PTX andα-CD + PTX; (b) 2-PAM, pure α -CD and α-CD + PTX; (Green and blue colored protons are of PTX and 2-PAM respectively while red colored protons are of CD).	227
	Fig.S3. FT-IR Spectra of (a) pure β -CD, PTX, and β-CD + PTX and (b) 2-PAM, pure β -CD, β-CD + PTX.	228
CHAPTER-IX	Fig.1. Molecular Structure of Uric acid.	231
	Fig.2. Schematic diagram of (a) Potassium oxalate, (b) Sodium oxalate, (c) Lithium oxalate.	231
	Fig.3. Plot of viscosity B-coefficient of Li-Ox (◆), Na-Ox (■), K-Ox (▼) at 293.15K (A), 303.15K (B), 313.15K (C) in aqueous Uric acid (UA) solution.	241
	Fig.4. Plausible solvation consequences of Li ⁺ , Na ⁺ and K ⁺ (hydrated) in aqueous UA medium.	243

