

LIST OF TABLES

Table No.	Title
1	Comparative effect of various concentrations of 2,4-D, TDZ, Dicamba and NAA on callus induction from immature male flower buds of banana cultivars on MS medium
2	Effect of various concentrations of NAA (0.25-2.0mg/l) in combination with 2,4-D (4.0 mg/l), TDZ (1.0mg/l) and Dicamba (3 mg/l) on callus induction from immature male flower buds of banana cultivars on MS medium.
3	Effect of various concentrations of IAA (0.25-2.0mg/l) in combination with 2,4-D (4.0 mg/l), TDZ (1.0mg/l) and Dicamba (3.0 mg/l) on callus induction from immature male flower buds of banana cultivars on MS medium
4	Effect of different concentrations of NAA and IAA (0.25-2.0 mg/l) with 2,4-D (4.0 mg/l) on callus induction from immature male flower buds of banana cultivars on MS medium
5	Effect of position of explant on callus induction on MS medium fortified with 4.0 mg/l of 2,4-D, 1.0 mg/l of IAA and 1.0 mg/l of NAA
6	Effect of position of explant on nature of callus on MS medium supplemented with 4.0 mg/l of 2,4-D, 1.0 mg/l of IAA and 1.0 mg/l of NAA
7	Influence of various concentrations of NAA and IAA (0.25-2.0 mg/l) in combination with 2,4-D (4.0 mg/l) on embryogenic callus induction from immature male flower buds of banana cultivars on MS medium
8	Effect of position of floral hands on embryogenic callus induction on MS medium supplemented with 4.0 mg/l of 2,4-D, 1.0 mg/l of IAA and 1.0 mg/l of NAA
9	Nature of embryogenic callus obtained from male floral hands on MS medium supplemented with 4.0 mg/l of 2,4-D, 1.0 mg/l of IAA and 1.0 mg/l of NAA
10	Effect of photo period on callus induction from immature male flower buds of banana cultivars on MS medium supplemented with 4.0 mg/l of 2,4-D, 1.0 mg/l of IAA and 1.0 mg/l of NAA
11	Effect of photo period on nature of callus obtained from immature male flower buds of banana cultivars on MS medium supplemented with 4.0 mg/l of 2,4-D, 1.0 mg/l of IAA and 1.0 mg/l of NAA
12	Comparison between photo period (6 hrs) and complete darkness on callus induction from male flower buds of banana cultivars on callus induction medium
13	Nature of callus obtained from male floral hands in callus induction medium incubated in 6 hrs light and complete darkness

14	Effect of preculture (liquid medium) and frequent subculture of explant on reduction of explant blackening
15	Effect of preculture of explant on embryogenic response
16	Influence of antioxidants and absorbants (L-Ascorbic acid, Activated charcoal and Polyvenyl pyrrolidone) on reduction of explant blackening.
17	Effect of carbohydrates on embryogenic complex induction from explants of banana cultivars on MS medium supplemented with 4.0 mg/l 2,4-D, 1.0 mg/l IAA and 1.0 mg/l NAA
18	Effect of different media on successful establishment of embryogenic cell suspension from the ideal callus obtained from banana cultivars
19	Effect of different concentration of sucrose on embryogenic response of banana cultivars in MS liquid medium supplemented with 1 mg/l 2,4-D
20	Cell density in embryogenic cell suspension of cvs. Rasthali and Nendran in MS medium with 1 mg/l 2,4-D
21	Change in settled cell volume (SCV) in suspension medium containing with 1 mg/l 2,4-D, with respect to time
22	Influence of various growth regulators (Zeatin, BAP, TDZ and Picloram) on induction of somatic embryos from ECS derived cultivars Rasthali and Nendran in SH medium.
23	Selection of optimum regeneration medium for ECS derived from immature male flower buds banana cultivars Rasthali and Nendran
24	Effect of substrate on somatic embryo formation from ECS of cvs. Rasthali and Nendran in modified regeneration medium (RIII)
25	Effect of cell density on somatic embryo formation from ECS of cvs. Rasthali and Nendran in modified regeneration medium (RIII)
26	Selection of maturation medium for the somatic embryos of two cvs. Rasthali and Nendran
27	Influence of maturation period on germination capacity of somatic embryos of two banana cultivars Rasthali and Nendran in modified regeneration medium (RIII)
28	Influence of age of somatic embryos and regeneration medium on germination efficiency in two commercial cultivars of banana Rasthali and Nendran.
29	Effect of age of ECS on regeneration capacity of banana cultivars Rasthali and Nendran in modified regeneration medium (RIII)
30	Effect of age of ECS on germination capacity of banana cultivars Rasthali and Nendran in MS medium with 0.5 mg/l BAP + 2 mg/l IAA

31	Effect of 2,4-D and BAP on repetitive embryogenesis in banana cultivars Rasthali and Nendran in MS medium
32	Effect of various concentrations of IAA (0.1 – 0.5 mg/l) in combination with 2,4-D (0.75 mg/l) and BAP (2.0 mg/l) on repetitive embryogenesis in banana cultivars Rasthali and Nendran
33	Influence of initial inoculum density on multiplication of somatic embryos in banana cultivars Rasthali and Nendran in MS medium with BAP (2 mg/l) and IAA (0.2 mg/l)
34	Effect of preculture of somatic embryos in liquid medium (20 days) to increase the germination capacity of banana cultivars Rasthali and Nendran
35	Contribution of various ISSR primers to percent polymorphism in cv. Rasthali and Nendran
36	Biometric analysis of ECS derived plants (cvs. Rasthali and Nendran) in field conditions
37	Evaluation of ECS derived plants in field conditions for somaclonal variation in cvs. Rasthali and Nendran